

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07K 5/06, C07D 401/12, 401/04, 401/14, 413/06, 211/56, 211/64, 207/14, 401/06, A61K 31/455, 38/05, C07D 209/44, C07K 5/02</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 96/35713</b>  (43) International Publication Date: 14 November 1996 (14.11.96)</p>
<p>(21) International Application Number: PCT/IB95/00333 (22) International Filing Date: 8 May 1995 (08.05.95)  (71) Applicant (for all designated States except US): PFIZER, INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).  (72) Inventors; and (75) Inventors/Applicants (for US only): CARPINO, Philip, A. [US/US]; 50 Meridian Street #1, Groton, CT 06340 (US). JARDINE, Paul, A., DaSilva [GY/US]; Apartment 21, 6 School Street, Mystic, CT 06355 (US). LEFKER, Bruce, A. [US/US]; 21 Eagle Ridge Drive, Gales Ferry, CT 06335 (US). RAGAN, John, A. [US/US]; 1 Lark Lane, Gales Ferry, CT 06335 (US).  (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., Patent Dept., 235 East 42nd Street, New York, NY 10017 (US).</p>		<p>(81) Designated States: CA, FI, JP, MX, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published With international search report.</p>
<p>(54) Title: DIPEPTIDES WHICH PROMOTE RELEASE OF GROWTH HORMONE</p> <p>(57) Abstract</p> <p>Compounds of formula (I) are growth hormone releasing peptide mimetics which are useful for the treatment and prevention of osteo- porosis.</p> <div style="text-align: center;"> <p>(I)</p> </div>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Larvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

-1-

### Dipeptides Which Promote Release of Growth Hormone

This invention relates to dipeptide compounds which are growth hormone releasing peptide (GHRP) mimetics and are useful for the treatment and prevention of osteoporosis.

5

### Background of the Invention

Growth hormone, which is secreted from the pituitary, stimulates growth of all tissues of the body that are capable of growing. In addition, growth hormone is known to have the following basic effects on the metabolic process of the body:

10

1. Increased rate of protein synthesis in all cells of the body;
2. Decreased rate of carbohydrate utilization in cells of the body;
3. Increased mobilization of free fatty acids and use of fatty acids for energy.

15

A deficiency in growth hormone secretion can result in various medical disorders, such as dwarfism.

20

It is now widely recognized that most human growth deficiencies are due to hypothalamic defects that impair the release of pituitary growth hormone and are not the result of a primary deficit in the production of growth hormone by the pituitary. As a result, the development of synthetic growth hormone-releasing agents and the use of drugs acting through established neurotransmitter systems in the brain to stimulate growth hormone release are being considered as alternatives to highly expensive growth hormone replacement therapy for the restoration of normal serum growth hormone levels. Strobel and Thomas, Pharm. Rev. 46, No. 1, pg. 1-34 (1994).

25

Various ways are known to release growth hormone. For example, chemicals such as arginine, L-3,4-dihydroxyphenylalanine (L-DOPA), glucagon, vasopressin, and insulin induced hypoglycemia, as well as activities such as sleep and exercise, indirectly cause growth hormone to be released from the pituitary by acting in some fashion on the hypothalamus perhaps either to decrease somatostatin secretion or to increase the secretion of the known secretagogue growth hormone releasing factor (GRF) or an unknown endogenous growth hormone-releasing hormone or all of these.

30

In cases where increased levels of growth hormone were desired, the problem was generally solved by providing exogenous growth hormone or by administering an agent which stimulated growth hormone production and/or release. In either case the peptidyl nature of the compound necessitated that it be administered by injection. Initially the source of growth hormone was the extraction of the pituitary glands of cadavers. This resulted in a very expensive

35

-2-

product and carried with it the risk that a disease associated with the source of the pituitary gland could be transmitted to the recipient of the growth hormone. Recently, recombinant growth hormone has become available which, while no longer carrying any risk of disease transmission, is still a very expensive


5 product which must be given by injection or by a nasal spray.

Other compounds have been developed which stimulate the release of endogenous growth hormone such as analogous peptidyl compounds related to GRF or the peptides of U.S. Patent 4,411,890. These peptides, while considerably smaller than growth hormones are still susceptible to various

10 proteases. As with most peptides, their potential for oral bioavailability is low. WO 94/13696 refers to certain spiropiperidines and homologs which promote release of growth hormone. Preferred compounds are of the general structure shown below.

15

20




25

WO 94/11012 refers to certain dipeptides that promote release of growth hormone. These dipeptides have the general structure


30

35



where L is

-3-




5

The compounds of WO 94/11012 and WO 94/13696 are reported to be useful in the treatment of osteoporosis in combination with parathyroid hormone or a bisphosphonate.

### Summary of the Invention

10 This invention provides a compound of the formula:

15




wherein

20 Z is  $-\text{COC}^*\text{R}^1\text{R}^{2c}\text{LCOANR}^4\text{R}^5$ ;  
L is  $\text{NR}^6$ , O or  $\text{CH}_2$ ;


W is hydrogen or in combination with X is a benzo fusion in which W and X are linked to form a phenyl ring optionally substituted with one to three substituents independently selected from  $\text{R}^{3a}$ ,  $\text{T-R}^{3b}$  and  $\text{R}^{12}$ ;

25 Y is hydrogen,  $\text{C}_1\text{-C}_6$  alkyl,  $\text{C}_4\text{-C}_{10}$  cycloalkyl or aryl, each optionally substituted with one to three substituents selected from  $\text{R}^{3a}$ ,  $\text{R}^{3b}$  and  $\text{R}^{12}$ ;  
X is  $\text{OR}^2$ ,

30



35



or in combination with W is a benzofusion in which W and X are linked to form a phenyl ring optionally substituted with one to three substituents independently selected from  $R^{3a}$ ,  $-T-R^{3b}$  and  $R^{12}$ ;

- 5  $R^1$  is C1-C10 alkyl, aryl, aryl (C1-C6 alkyl) and C3-C7 cycloalkyl (C1-C6 alkyl) or C1-C5 alkyl-K-C1-C5 alkyl, aryl (C0-C5 alkyl)-K-(C1-C5 alkyl), (C3-C7) cycloalkyl(C-C5alkyl)-K-(C1-C5 alkyl) where K is O, S(O)<sub>m</sub>,  $N(R^2)C(O)$ ,  $C(O)N(R^2)$ ,  $OC(O)$ ,  $C(O)O$ ,  $-CR^2=CR^2-$  or  $-C\equiv C-$  where the aryl groups are defined below and  $R^2$  and the alkyl groups may be further  
 10 substituted by 1-5 halogens,  $S(O)_mR^{2a}$ , 1 to 3 of  $OR^{2a}$  or  $C(O)OR^{2a}$  and the aryl groups may be further substituted by phenyl, phenoxy, arylalkyloxy, halophenyl, 1 to 3 of C1-C6 alkyl, 1 to 3 of halogen, 1 to 2 of  $OR^2$ , methylenedioxy,  $S(O)_mR^2$ , 1 to 2 of  $CF_3$ ,  $OCF_3$ , nitro,  $N(R^2)(R^2)$ ,  $N(R^2)C(O)(R^2)$ ,  $C(O)OR^2$ ,  $C(O)N(R^2)(R^2)$ ,  $SO_2N(R^2)(R^2)$ ,  $N(R^2)SO_2$  aryl or  
 15  $N(R^2)SO_2R^2$ ;

$R^{2c}$  is hydrogen, C1-C6 alkyl, C3-C7 cycloalkyl, and may be joined with  $R^1$  to form a C3-C8 ring optionally including oxygen, sulfur or  $NR^{2a}$ ;

$R^2$  is hydrogen, C1-C6 alkyl, C3-C7 cycloalkyl;

$R^{2a}$  is hydrogen or C1-C6 alkyl;

- 20  $R^{2b}$  is hydrogen C1-C8 alkyl, C1-C8 halogenated alkyl, C3-C8 cycloalkyl, alkylaryl or aryl;

$R^{3a}$  is H, F, Cl, Br, I,  $CH_3$ ,  $OCH_3$  or  $CF_3$ ;

$R^{12}$  is H, F, Cl, Br, I,  $CH_3$ ,  $OCH_3$  or  $CF_3$ ;

- T is a bond or is phenyl or a 5 or 6-membered heterocycle containing 1  
 25 to 3 hetero atoms selected from nitrogen, sulfur or oxygen, each optionally substituted with one to three substituents selected from F, Cl, Br, I,  $CH_3$ ,  $OCH_3$ ,  $OCF_3$  and  $CF_3$ ;

- $R^{3b}$  is hydrogen,  $CONR^8R^9$ ,  $SO_2NR^8R^9$ ,  $COOH$ ,  $COO(C1-C6)alkyl$ ,  $NR^2SO_2R^9$ ,  $NR^2CONR^8R^9$ ,  $NR^2SO_2NR^8R^9$ ,  $NR^2C(O)R^9$ , imidazolyl, thiazolyl  
 30 or tetrazolyl;

- $R^4$  and  $R^5$  are independently hydrogen, C1-C6 alkyl, substituted C1-C6 alkyl where the substituents may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3 C1-C10 alkanoyloxy, 1 to 3 C1-C6 alkoxy, phenyl, phenoxy, 2-furyl, C1-C6 alkoxycarbonyl,  $S(O)_m(C1-C6 alkyl)$ ; or  $R^4$  and  $R^5$  can be taken together to  
 35 form  $-(CH_2)_rL_a(CH_2)_s-$  where  $L_a$  is  $C(R^2)_2$ , O,  $S(O)_m$  or  $N(R^2)$ , r and s are independently 1 to 3 and  $R^2$  is as defined above;

$R^6$  is hydrogen or C1-C6 alkyl, and may be joined with  $R^{2c}$  to form a C3-C8 ring;


-5-

R<sup>50</sup> is 4-morpholino, 4-(1-methylpiperazinyl), C<sub>3</sub>-C<sub>7</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl each optionally substituted with one to three substituents individually selected from F, OH, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub> and C<sub>3</sub>-C<sub>7</sub> cycloalkyl;

M is -C(O)- or -SO<sub>2</sub>-;

5 A<sup>1</sup> is a bond, C<sub>1</sub>-C<sub>6</sub> alkylene, C<sub>1</sub>-C<sub>6</sub> haloalkylenyl or C<sub>1</sub>-C<sub>6</sub> hydroxyalkylenyl;

A is a bond or is



where x and y are independently 0-3;

15 Z<sup>1</sup> is N-R<sup>2</sup> or O; or Z<sup>1</sup> is a bond;

R<sup>7</sup> and R<sup>7a</sup> are independently hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, trifluoromethyl, phenyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl where the substituents are imidazolyl, phenyl, indolyl, p-hydroxyphenyl, OR<sup>2</sup>, S(O)<sub>m</sub>R<sup>2</sup>, C(O)OR<sup>2</sup>, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, N(R<sup>2</sup>)(R<sup>2</sup>), C(O)N(R<sup>2</sup>)(R<sup>2</sup>); or R<sup>7</sup> and R<sup>7a</sup> can independently be joined to one  
 20 or both of R<sup>4</sup> and R<sup>5</sup> groups to form alkylene bridges between the terminal nitrogen and the alkyl portion of the R<sup>7</sup> and R<sup>7a</sup> groups, wherein the bridge contains 1 to 5 carbon atoms; or R<sup>7</sup> and R<sup>7a</sup> may be joined to form a 3- to 7-membered ring.

R<sup>9</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl  
 25 each optionally substituted with one to three substituents selected from Cl, F, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>;

R<sup>8</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl where the substituents may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3 C<sub>1</sub>-C<sub>10</sub> alkanoyloxy, 1 to 3 C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl, S(O)<sub>m</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl);  
 30 or R<sup>8</sup> and R<sup>9</sup> can be taken together to form -(CH<sub>2</sub>)<sub>r</sub>L<sub>a</sub>(CH<sub>2</sub>)<sub>s</sub>- where L<sub>a</sub> is C(R<sup>2</sup>)<sub>2</sub>, O, S(O)<sub>m</sub> or N(R<sup>2</sup>), r and s are independently 1 to 3.

F<sup>1</sup>, B, D and E are carbon, nitrogen, one of which is joined to A<sup>1</sup> and each of the remaining of F<sup>1</sup>, B, D and E may be optionally substituted with R<sup>aa</sup>, R<sup>bb</sup>; or F<sup>1</sup>, B, D and E, if not joined to A<sup>1</sup> may also be sulfur, oxygen or  
 35 carbonyl; F<sup>1</sup>, B, D and E may form a saturated or unsaturated ring; and one of F<sup>1</sup>, B, D or E may be optionally missing to afford a saturated or unsaturated five-membered ring;

-6-

R<sup>aa</sup> is H, C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one to three halogens; or phenyl optionally substituted with one to three substituents independently selected from halogen, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub> and C<sub>1</sub>-C<sub>8</sub> alkyl aryl;

5 R<sup>bb</sup> is C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>1</sub>-C<sub>8</sub> thioalkyl, each optionally substituted with one to three substituents independently selected from halogen, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub>; C<sub>3</sub>-C<sub>8</sub> cycloalkyl; phenyl optionally substituted with one to three substituents independently selected from halo, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>; -O-C<sub>1</sub>-C<sub>8</sub> alkyl; or -S-C<sub>1</sub>-C<sub>8</sub> alkyl.

10 G, H<sup>1</sup>, I and J are carbon, nitrogen, sulfur or oxygen atoms which form an aromatic ring, a partially saturated ring or a saturated ring; one of G, H<sup>1</sup>, I or J may be optionally missing to afford a five-membered ring;

aryl is phenyl, naphthyl or a 5- or 6- membered ring with 1 to 3 heteroatoms selected from oxygen, sulfur and nitrogen; or a bicyclic ring system consisting of a 5 or 6 membered heterocyclic ring with 1 to 3  
15 heteroatoms of nitrogen, sulfur or oxygen, fused to a phenyl ring, each aryl ring being optionally and independently substituted with up to three substituents selected from R<sup>3a</sup>, R<sup>3b</sup> and R<sup>12</sup>;

m is 0 to 2;

n is 0 to 2;

20 q is 0 to 3;

and pharmaceutically acceptable salts and individual diastereomers thereof.

In one aspect, this invention provides a compound of Formula I wherein Z is -COCR<sup>1</sup>R<sup>2c</sup>NHCOANR<sup>4</sup>R<sup>5</sup>

25 A is -CR<sup>7</sup>R<sup>7a</sup>(CH<sub>2</sub>)<sub>Y</sub>; Y is 0 to 3;

and R<sup>2c</sup> is H or CH<sub>3</sub>;

R<sup>7</sup> is C<sub>1</sub>-C<sub>3</sub> alkyl;

R<sup>7a</sup> is H or C<sub>1</sub>-C<sub>3</sub> alkyl;

30 R<sup>4</sup> is hydrogen or C<sub>1</sub>-C<sub>3</sub> alkyl; or R<sup>4</sup> and R<sup>7a</sup> are combined to form an alkylene bridge;

R<sup>5</sup> is hydrogen or C<sub>1</sub>-C<sub>3</sub> alkyl optionally substituted with one or two hydroxyl groups.

In another aspect, this invention provides a compound of Formula I wherein


35 R<sup>1</sup> is selected from the group consisting of 1-indolyl-CH<sub>2</sub>-; 2-indolyl-CH<sub>2</sub>-; 3-indolyl-CH<sub>2</sub>-; 1-naphthyl-CH<sub>2</sub>-; 2-naphthyl-CH<sub>2</sub>-; 1-benzimidazolyl-CH<sub>2</sub>-; 2-benzimidazolyl-CH<sub>2</sub>-; phenyl-(C<sub>1</sub>-C<sub>4</sub>) alkyl-; 2-, 3- or 4-pyridyl-(C<sub>1</sub>-C<sub>4</sub>) alkyl; thienyl-(C<sub>1</sub>-C<sub>4</sub>) alkyl; phenyl-(CH<sub>2</sub>)<sub>n</sub>-O-CH<sub>2</sub>- where n is zero to



-7-


three; phenyl-CH<sub>2</sub>O-phenyl-CH<sub>2</sub>-; and 3-benzothieryl-CH<sub>2</sub>-; or any of the above groups substituted in the aryl portion with one to three F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub> substituents.

In another aspect, this invention provides a compound of Formula I wherein R<sup>1</sup> is C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>OCH<sub>2</sub>-, 1-naphthyl-CH<sub>2</sub>-, 2-naphthyl-CH<sub>2</sub>-, phenylpropyl or



and the aryl portion of R<sup>1</sup> is optionally substituted with fluorine, CH<sub>3</sub> or CF<sub>3</sub>.

In another aspect, this invention provides a compound of Formula I which is:



wherein n is zero or one; m is one or two;

R<sup>3a</sup> is H, F, Cl, Br, I, CH<sub>3</sub>, OCH<sub>3</sub> or CF<sub>3</sub>;

R<sup>12</sup> is H, F, Cl, Br, I, CH<sub>3</sub>, OCH<sub>3</sub> or CF<sub>3</sub>;

T is a bond or is an aryl group selected from phenyl, pyridyl, pyrimidyl, thienyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl and tetrazolyl; each optionally substituted with one to three F, CH<sub>3</sub>, Cl, OR<sup>8</sup>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>;

R<sup>3b</sup> is hydrogen, CONR<sup>8</sup>R<sup>9</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, COOH, COO(C<sub>1</sub>-C<sub>6</sub>)alkyl, NH<sub>2</sub>SO<sub>2</sub>R<sup>9</sup>, NHC(O)NR<sup>8</sup>R<sup>9</sup>, NHC(O)R<sup>9</sup>, NR<sup>8</sup>R<sup>9</sup>, imidazolyl, thiazolyl or tetrazolyl;

-8-

$R^9$  is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl each optionally substituted with one to three substituents selected from Cl, F, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>;

$R^8$  is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl  
 5 where the substituents may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3 C<sub>1</sub>-C<sub>10</sub> alkanoyloxy, 1 to 3 C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl, S(O)<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl); or  $R^8$  and  $R^9$  can be taken together to form -(CH<sub>2</sub>)<sub>r</sub>L<sub>a</sub>(CH<sub>2</sub>)<sub>s</sub>- where L<sub>a</sub> is C(R<sup>2</sup>)<sub>2</sub>, O, S(O)<sub>2</sub> or N(R<sup>2</sup>), r and s are independently 1 to 3.

10 In another aspect, this invention provides a compound of Formula II wherein:

m is one;

$R^{12}$  is hydrogen;

$R^{3a}$  is hydrogen, F or Cl;

15 T is phenyl, thienyl, thiazolyl, oxazolyl, isoxazolyl or pyrazolyl, each optionally substituted with one to three substituents, selected from F, OH, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub> and CH<sub>3</sub>;

$R^{3b}$  is hydrogen, C(O)NR<sup>8</sup>R<sup>9</sup>, NHC(O)NR<sup>8</sup>R<sup>9</sup>, NHS(O)<sub>2</sub>R<sup>9</sup>, NHC(O)R<sup>9</sup>;

20 In another aspect, this invention provides a compound of Formula II wherein:

m is one;

T is a bond;

n is one;

$R^{3a}$  and  $R^{12}$  are hydrogen.

25  $R^{3b}$  is CONR<sup>8</sup>R<sup>9</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, COOH, COO(C<sub>1</sub>-C<sub>6</sub>)alkyl, NHSO<sub>2</sub>R<sup>9</sup>, NHSO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, NHC(O)NR<sup>8</sup>R<sup>9</sup>, NHC(O)R<sup>9</sup>, NR<sup>8</sup>R<sup>9</sup> or OR<sup>8</sup>;

$R^9$  is hydrogen, phenyl, or thienyl optionally substituted with one to three substituents selected from F, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; or  $R^9$  is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one to three substituents selected from F, OH,  
 30 OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; or R<sup>8</sup>;

$R^8$  is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl or C<sub>3</sub>-C<sub>7</sub> cycloalkyl where the substituents may be 1 to 5 halo or 1 to 3 hydroxy; or  $R^8$  and  $R^9$  can be taken together to form -(CH<sub>2</sub>)<sub>r</sub>L<sub>a</sub>(CH<sub>2</sub>)<sub>s</sub>- wherein L<sub>a</sub> is C(R<sup>2</sup>)<sub>2</sub>, O, S(O)<sub>m</sub> or N(R<sup>2</sup>) where r and s are independently 1 to 3.

35 In another aspect, this invention provides a compound of Formula I


n is one;

m is one;

W is hydrogen;

-9-

Y is hydrogen or methyl;



5 M is -C(O)- ;

R<sup>50</sup> is 4-morpholino, 4-(1-methylpiperazinyl), C<sub>3</sub>-C<sub>7</sub> cycloalkyl or C<sub>1</sub>-C<sub>6</sub> alkyl each optionally substituted with one to three substituents individually selected from F, OH, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>;

Ar is phenyl, pyridyl, thienyl, pyrimidyl or thiazolyl, each optionally substituted with one to three substituents individually selected from F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>.

In another aspect, this invention provides a compound of Formula I wherein

n and m are one;

15 Y is phenyl, pyridyl, pyrimidyl, thienyl, or thiazolyl, each optionally substituted with one to three substituents independently selected from F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>;

X is -C(O)-NR<sup>8</sup>R<sup>9</sup>;

20 R<sup>9</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl each optionally substituted with one to three substituents selected from Cl, F, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>;

R<sup>8</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, substituted C<sub>1</sub>-C<sub>6</sub> alkyl where the substituents may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3 C<sub>1</sub>-C<sub>10</sub> alkanoyloxy, 1 to 3 C<sub>1</sub>-C<sub>6</sub> alkoxy, phenyl, phenoxy, C<sub>1</sub>-C<sub>6</sub> alkoxycarbonyl, S(O)<sub>m</sub>(C<sub>1</sub>-C<sub>6</sub> alkyl);  
 25 or R<sup>8</sup> and R<sup>9</sup> can be taken together to form -(CH<sub>2</sub>)<sub>r</sub>L<sub>a</sub>(CH<sub>2</sub>)<sub>s</sub>- where L<sub>a</sub> is C(R<sup>2</sup>)<sub>2</sub>, O, S(O)<sub>m</sub> or N(R<sup>2</sup>), r and s are independently 1 to 3.

In another aspect, this invention provides a compound of Formula I wherein

Z is -COCR<sup>1</sup>R<sup>2c</sup>NHCOANR<sup>4</sup>R<sup>5</sup>

30 A is -CR<sup>7</sup>R<sup>7a</sup>(CH<sub>2</sub>)<sub>Y</sub>; Y is 0 to 3;

and R<sup>2c</sup> is H or CH<sub>3</sub>;

R<sup>7</sup> is C<sub>1</sub>-C<sub>3</sub> alkyl;

R<sup>7a</sup> is H or C<sub>1</sub>-C<sub>3</sub> alkyl;

R<sup>4</sup> is hydrogen or C<sub>1</sub>-C<sub>3</sub> alkyl; or R<sup>4</sup> and R<sup>7a</sup> are combined to form an  
 35 alkylene bridge;


R<sup>5</sup> is hydrogen or C<sub>1</sub>-C<sub>3</sub> alkyl optionally substituted with one or two hydroxyl groups;

-10-

R<sup>1</sup> is selected from the group consisting of 1-indolyl-CH<sub>2</sub>-; 2-indolyl-CH<sub>2</sub>-; 3-indolyl-CH<sub>2</sub>-; 1-naphthyl-CH<sub>2</sub>-; 2-naphthyl-CH<sub>2</sub>-; 1-benzimidazolyl-CH<sub>2</sub>-; 2-benzimidazolyl-CH<sub>2</sub>-; phenyl-C<sub>1</sub>-C<sub>4</sub> alkyl-; 2-, 3- or 4-pyridyl-C<sub>1</sub>-C<sub>4</sub> alkyl-; thienyl-C<sub>1</sub>-C<sub>4</sub> alkyl-; phenyl-(CH<sub>2</sub>)<sub>n</sub>-O-CH<sub>2</sub>- where n is zero to three;  
 5 phenyl-CH<sub>2</sub>O-phenyl-CH<sub>2</sub>-; and 3-benzothienyl-CH<sub>2</sub>-; or any of the above groups substituted in the aryl portion with one to three F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub> substituents;

Y is hydrogen and X is:


10



15

wherein n and m are one and A<sup>1</sup> is a bond, or X is:


20



25

or

30



wherein R<sup>3a</sup> is hydrogen, F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>;


R<sup>12</sup> is hydrogen, F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>;

35 R<sup>aa</sup> is hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one to three halogens; or phenyl optionally substituted with one to three substituents independently selected from halogen, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub> and C<sub>1</sub>-C<sub>8</sub> alkyl aryl;


-11-

$R^{bb}$  is hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl, optionally substituted with one to three substituents independently selected from halogen, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub>; C<sub>3</sub>-C<sub>8</sub> cycloalkyl; phenyl optionally substituted with one to three substituents independently selected from halo, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>; -O-C<sub>1</sub>-C<sub>8</sub> alkyl; or -S-C<sub>1</sub>-C<sub>8</sub> alkyl.

In a preferred aspect this invention provides a compound of Formula I wherein Z is



15 or




20

wherein each n is independently zero or one;

R<sub>1</sub> is 1-naphthyl CH<sub>2</sub>-, 2-naphthyl CH<sub>2</sub>, phenyl CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-

25



30 or -CH<sub>2</sub>OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>.

Preferred compounds of Formula I are:

(R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[6-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide hydrochloride;

(R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride;

-12-

- (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid phenylamide hydrochloride;
- (R)-2-Amino-N-(2-(1H-indol-3-yl)-1-[6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl)-isobutyramide hydrochloride;
- 5 (R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide;
- (R)-N-{2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl}-benzamide;
- 10 (R)-2-Amino-N-{1-benzyloxymethyl-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethyl}-isobutyramide;
- 2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(phenyl-propionyl-amino)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;
- (R)-N-[1-[4-(Acetyl-phenyl-amino)-piperidine-1-carbonyl]-2-(1H-indol-3-yl)-ethyl]-2-amino-isobutyramide;
- 15 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride;
- 2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-;
- 20 2-Amino-N-{1-(R)-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide;
- (R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide;
- 2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(1H-indol-3-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride;
- 25 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(2-methyl-benzoimidazol-1-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride;
- (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;
- 30 (R)-2-Amino-N-[2-[4-(6-fluoro-benzo[d]isoxazol-3-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide hydrochloride;
- (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-1,2-dihydro-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;
- (R)-2-Amino-N-[2-[4-(5-chloro-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide
- 35 (R)-2-Amino-N-{1-benzo[b]thiophen-3-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;

-13-

- 2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide;
- (R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[5-(toluene-4-sulfonylamino)-1,3-dihydro-isoindol-2-yl]-ethyl}-amide
- 5 hydrochloride;
- 2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide;
- (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-phenyl-4-(pyrrolidine-1-carbonyl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;
- 10 (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (4-hydroxy-butyl)-amide hydrochloride;
- 2-Amino-N-{1-(5-fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;
- 15 2-Amino-N-{1-(1-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;
- (R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride;
- (R)-3-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-3-methyl-butyramide hydrochloride;
- 20 and
- (R)-Piperidine-4-carboxylic acid {2-[4-(3-methyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethyl}-amide hydrochloride
- 25 This invention also provides:
- a method for increasing levels of endogenous growth hormone in a human or an animal which comprises administering to such human or animal an effective amount of a compound of Formula I;
- a composition useful for increasing the endogenous production or
- 30 release of growth hormone in a human or an animal which comprises an inert carrier and an effective amount of a compound of Formula I;
- a composition useful for increasing the endogenous production or release of growth hormone in a human or an animal which comprises an inert carrier and an effective amount of a compound of Formula I used in
- 35 combination with other growth hormone secretagogues such as, GHRP-6, Hexorelan, GHRP-1, growth hormone releasing factor (GRF) or one of its analogs or IGF-1 or IGF-2, or B-HT920; and

-14-

a method for the treatment of osteoporosis which comprises administering to a patient with osteoporosis a combination of a bisphosphonate compound such as alendronate, and a compound of Formula I.

5 This invention further provides a method for treating or preventing diseases or conditions which may be treated or prevented by growth hormone which comprises administering to a mammal in need of such treatment or prevention an amount of a compound of Formula I which is effective in promoting release of said growth hormone.

10 In another aspect this invention provides compositions and methods which are useful for treating obesity, frailty associated with old age, and cachexia associated with AIDS and cancer.

The instant compounds are highly substituted dipeptide analogs for promoting the release of growth hormone which are stable under various  
15 physiological conditions which may be administered parenterally, nasally or by the oral route.

#### Detailed Description of the Invention

One of ordinary skill will recognize that certain substituents listed in this invention will be chemically incompatible with one another or with the  
20 heteroatoms in the compounds, and will avoid these incompatibilities in selecting compounds of this invention.

In general the compounds of Formula I can be made by processes which include processes known in the chemical arts for the production of structurally analogous compounds. Certain processes for the manufacture of  
25 Formula I compounds are provided as further features of the invention and are illustrated by the following reaction schemes.

In the above structural formulae and throughout the instant specification, the following terms have the indicated meanings:

The alkyl groups specified above are intended to include those alkyl  
30 groups of the designated length in either a straight or branched configuration which may optionally contain double or triple bonds. Exemplary of such alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, tertiary butyl, pentyl, isopentyl, hexyl, isohexyl, allyl, ethinyl, propenyl, butadienyl, hexenyl and the like.

35 The alkoxy groups specified above are intended to include those alkoxy groups of the designated length in either a straight or branched configuration which may optionally contain double or triple bonds. Exemplary of such alkoxy groups are methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tertiary



-15-

butoxy, pentoxy, isopentoxy, hexoxy, isohexoxy, allyloxy, 2-propinyloxy, isobutenyloxy, hexenyloxy and the like.

The term "halogen" or "halo" is intended to include the halogen atoms fluorine, chlorine, bromine and iodine.

- 5       The term "aryl" is intended to include phenyl and naphthyl and aromatic 5- and 6-membered rings with 1 to 3 heteroatoms or fused 5- or 6-membered bicyclic rings with 1 to 3 heteroatoms of nitrogen, sulfur or oxygen. Examples of such heterocyclic aromatic rings are pyridine, thiophene, furan, benzothiophene, tetrazole, indole, N-methylindole, dihydroindole, indazole, N-  
10 formylindole, benzimidazole, thiazole, pyrimidine, and thiadiazole.

Certain of the above defined terms may occur more than once in the above formula and upon such occurrence each term shall be defined independently of the other.

- 15       Throughout the instant application the following abbreviations are used with the following meanings:

-16-

	BOC	t-butyloxycarbonyl
	BOP	Benzotriazol-1-yloxy
		tris(dimethylamino) phosphonium
		hexafluorophosphate
5	CBZ	Benzyloxycarbonyl
	CDI	N,N'-Carbonyldiimidazole
	CH <sub>2</sub> Cl <sub>2</sub>	Methylene chloride
	CHCl <sub>3</sub>	Chloroform
	DCC	Dicyclohexylcarbodiimide
10	DMF	Dimethylformamide
	EDC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
	EtOAc	Ethyl acetate
	Fmoc	9-Fluorenylmethoxycarbonyl
15	Hex	Hexane
	HOBT	Hydroxybenzotriazole hydrate
	HPLC	High pressure liquid chromatography
	MHz	Megahertz
	MS	Mass Spectrum
20	NMR	Nuclear Magnetic Resonance
	PTH	Parathyroid hormone
	TFA	Trifluoroacetic acid
	THF	Tetrahydrofuran
	TLC	Thin layer chromatography
25	TRH	Thyrotropin releasing hormone

The compounds of the instant invention all have at least one asymmetric center as noted by the asterisk in the structural Formula I, group Z above. Additional asymmetric centers may be present on the molecule depending upon the nature of the various substituents on the molecule. Each such asymmetric center will produce two optical isomers and it is intended that all such optical isomers, as separated, pure or partially purified optical isomers, racemic mixtures or diastereomeric mixtures thereof, be included within the ambit of the instant invention. In the case of the asymmetric center represented by the asterisk, it has been found that the absolute stereochemistry of the more active and thus more preferred isomer is shown in Formula IA. This preferred absolute configuration also applies to Formula I. With the R<sub>2</sub> substituent as hydrogen, the spatial configuration of the asymmetric center corresponds to

-17-

that in a D-amino acid. In most cases this is also designated an R-configuration although this will vary according to the values of R<sub>1</sub> and R<sub>2</sub> used in making R- or S-stereochemical assignments.

The instant compounds are generally isolated in the form of their  
5 pharmaceutically acceptable acid addition salts, such as the salts derived from using inorganic and organic acids. Examples of such acids are hydrochloric, nitric, sulfuric, phosphoric, formic, acetic, trifluoroacetic, propionic, maleic, succinic, malonic, methane sulfonic and the like. In addition, certain  
10 compounds containing an acidic function such as a carboxy can be isolated in the form of their inorganic salt in which the counterion can be selected from sodium, potassium, lithium, calcium, magnesium and the like, as well as from organic bases.

The growth hormone releasing compounds of Formula I are useful *in vitro* as unique tools for understanding how growth hormone secretion is  
15 regulated at the pituitary level. This includes use in the evaluation of many factors thought or known to influence growth hormone secretion such as age, sex, nutritional factors, glucose, amino acids, fatty acids, as well as fasting and non-fasting states. In addition, the compounds of this invention can be used in the evaluation of how other hormones modify growth hormone releasing  
20 activity. For example, it has already been established that somatostatin inhibits growth hormone release. Other hormones that are important and in need of study as to their effect on growth hormone release include the gonadal hormones, e.g., testosterone, estradiol, and progesterone; the adrenal hormones, e.g., cortisol and other corticoids, epinephrine and norepinephrine;  
25 the pancreatic and gastrointestinal hormones, e.g., insulin, glucagon, gastrin, secretin; the vasoactive peptides, e.g., bombesin, the neurokinins; and the thyroid hormones, e.g., thyroxine and triiodothyronine. The compounds of Formula I can also be employed to investigate the possible negative or positive feedback effects of some of the pituitary hormones, e.g., growth hormone and  
30 endorphin peptides, on the pituitary to modify growth hormone release. Of particular scientific importance is the use of these compounds to elucidate the subcellular mechanisms mediating the release of growth hormone.

The compounds of Formula I can be administered to animals, including man, to release growth hormone *in vivo*. For example, the compounds can be  
35 administered to commercially important animals such as swine, cattle, sheep and the like to accelerate and increase their rate and extent of growth, to improve feed efficiency and to increase milk production in such animals. In addition, these compounds can be administered to humans *in vivo* as a

-18-

diagnostic tool to directly determine whether the pituitary is capable of releasing growth hormone. For example, the compounds of Formula I can be administered *in vivo* to children. Serum samples taken before and after such administration can be assayed for growth hormone. Comparison of the  
5 amounts of growth hormone in each of these samples would be a means for directly determining the ability of the patient's pituitary to release growth hormone.

Accordingly, the present invention includes within its scope pharmaceutical compositions comprising, as an active ingredient, at least one  
10 of the compounds of Formula I in association with a pharmaceutical by acceptable carrier. Optionally, the active ingredient of the pharmaceutical compositions can comprise an anabolic agent in addition to at least one of the compounds of Formula I or another composition which exhibits a different activity, e.g., an antibiotic growth permittant or an agent to treat osteoporosis or  
15 in combination with a corticosteroid to minimize the catabolic side effects or with other pharmaceutically active materials wherein the combination enhances efficacy and minimizes side effects.

Growth promoting and anabolic agents include, but are not limited to, TRH, PTH, diethylstilbesterol, estrogens,  $\beta$ -agonists, theophylline, anabolic  
20 steroids, enkephalins, E series prostaglandins, compounds disclosed in U.S. Patent No. 3,239,345, e.g., zeranol, and compounds disclosed in U.S. Patent No. 4,036,979, e.g., sulbenox or peptides disclosed in U.S. Patent No. 4,411,890.

A still further use of the growth hormone secretagogues of this invention  
25 is in combination with other growth hormone secretagogues such as the growth hormone releasing peptides GHRP-6, GHRP-1 as described in U.S. Patent Nos. 4,411,890 and publications WO 89/07110, WO 89/07111 and B-HT920 as well as hexarelin and the newly discovered GHRP-2 as described in WO 93/04081 or growth hormone releasing hormone (GHRH, also designated  
30 GRF) and its analogs or growth hormone and its analogs or somatomedins including IGF-1 and IGF-2 or  $\mu$ -adrenergic agonists such as clonidine or serotonin 5HT<sub>1D</sub> agonists such as sumatriptan or agents which inhibit somatostatin or its release such as physostigmine and pyridostigmine.

As is well known to those skilled in the art, the known and potential uses  
35 of growth hormone are varied and multitudinous. See "Human Growth Hormone", Strobel and Thomas, Pharmacological Reviews 46, pg. 1-34 (1994) which describes possible use of growth hormone. Thus, the administration of the compounds of this invention for purposes of stimulating the release of

-19-

endogenous growth hormone can have the same effects or uses as growth hormone itself. These varied uses of growth hormone may be summarized as follows: stimulating growth hormone release in elderly humans; treating growth hormone deficient adults; prevention of catabolic side effects of glucocorticoids, treatment of osteoporosis, stimulation of the immune system, acceleration of wound healing, accelerating bone fracture repair, treatment of growth retardation, treating acute or chronic renal failure or insufficiency, treatment of physiological short stature, including growth hormone deficient children, treating short stature associated with chronic illness, treatment of obesity, treating growth retardation associated with Prader-Willi syndrome and Turner's syndrome; accelerating the recovery and reducing hospitalization of burn patients or following major surgery such as gastrointestinal surgery; treatment of intrauterine growth retardation, skeletal dysplasia, hypercortisonism and Cushings syndrome; replacement of growth hormone in stressed patients; treatment of osteochondrodysplasias, Noonans syndrome, sleep disorders, Alzheimer's disease, delayed wound healing, and psychosocial deprivation; treatment of pulmonary dysfunction and ventilator dependency; attenuation of protein catabolic response after a major operation; treating malabsorption syndromes, reducing cachexia and protein loss due to chronic illness such as cancer or AIDS; accelerating weight gain and protein accretion in patients on TPN (total parenteral nutrition); treatment of hyperinsulinemia including nesidioblastosis; adjuvant treatment for ovulation induction and to prevent and treat gastric and duodenal ulcers; to stimulate thymic development and prevent for age-related decline of thymic function; adjunctive therapy for patients on chronic hemodialysis; treatment of immunosuppressed patients and to enhance antibody response following vaccination; improvement in muscle strength, mobility, maintenance of skin thickness, metabolic homeostasis, renal hemeostasis in the frail elderly; stimulation of osteoblasts, bone remodeling, and cartilage growth; treatment of neurological diseases such as peripheral and drug induced neuropathy, Guillian-Barre Syndrome, amyotrophic lateral sclerosis, multiple sclerosis, cerebrovascular accidents and demyelinating diseases; stimulation of the immune system in companion animals and treatment of disorders of aging in companion animals; growth promotant in livestock; and stimulation of wool growth in sheep.

It will be known to those skilled in the art that there are numerous compounds now being used in an effort to treat the diseases or therapeutic indications enumerated above. Combinations of these therapeutic agents

-20-

some of which have also been mentioned above with the growth promotant, anabolic and desirable properties of these various therapeutic agents. In these combinations, the therapeutic agents and the growth hormone secretagogues of this invention may be independently present in dose ranges from one one-hundredth to one times the dose levels which are effective when these compounds and secretagogues are used singly.

Combined therapy to inhibit bone resorption, prevent osteoporosis and enhance the healing of bone fractures can be illustrated by combinations of bisphosphonates and the growth hormone secretagogues of this invention.

The use of bisphosphonates for these utilities has been reviewed, for example, by Hamdy, N.A.T., Role of Bisphosphonates in Metabolic Bone Diseases. *Trends in Endocrinol. Metab.*, 1993, 4, 19-25. Bisphosphonates with these utilities include alendronate, tiludronate, dimethyl - APD, risedronate, etidronate, YM-175, clodronate, pamidronate, and BM-210995. According to their potency, oral daily dosage levels of the bisphosphonate of between 0.1 mg and 5 g and daily dosage levels of the growth hormone secretagogues of this invention of between 0.01 mg/kg to 20 mg/kg of body weight are administered to patients to obtain effective treatment of osteoporosis.

Compounds that have the ability to stimulate GH secretion from cultured rat pituitary cells are identified using the following protocol. This test is also useful for comparison to standards to determine dosage levels. Cells are isolated from anterior pituitaries of 6-week old male Wistar rats following decapitation. Tissues are finely minced, then subjected to mechanical and enzymatic dispersion using 10 U/mL bacterial protease (EC 3.4.24.4, Sigma P-6141) in Hank's balanced salt solution without calcium or magnesium. The cell suspension is plated at  $5 \times 10^4$  cells per square cm in 24-well Costar dishes and cultured for 3 days in Dulbecco's Modified Eagles Medium (D-MEM) supplemented with 4.5 g/L glucose, 10% horse serum, 2.5% fetal bovine serum, non-essential amino acids, 100 U/mL nystatin and 50 mg/mL gentamycin sulfate.

Just prior to beginning the assay, culture wells are rinsed twice, then equilibrated for 30 minutes in release medium (D-MEM buffered with 25 mM Hepes, pH 7.4 and containing 0.5% bovine serum albumin at 37°C. Test compounds are dissolved in DMSO, then diluted into pre-warmed release medium. The assay is initiated by adding 1 mL of release medium (with test compounds) to each culture well. Incubation is carried out at 37°C for 15 minutes, then terminated by removal of the culture medium, which is centrifuged at 2000 x g for 15 minutes to remove cellular material before being

-21-

assayed for rat growth hormone by a radioimmunoassay procedure using reagents provided by A.F. Parlow (Harbor-UCLA Medical Center, Torrance, CA). Active compounds typically stimulate growth hormone release by 3-4 fold.

- Nonpeptidyl GH secretagogues are shown to mimic growth hormone releasing peptides based on one or more of the following criteria: synergistic stimulation of GH release when added together with GHRH, inability to further increase GH secretion by GHRP-6, sensitivity to protein kinase C inhibitors, and selective stimulation of biphasic calcium flux in GH-containing cells.

- The compounds of this invention can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous or subcutaneous injection, or implant), nasal, vaginal, rectal, sublingual, or topical routes of administration and can be formulated with pharmaceutically acceptable carriers to provide dosage forms appropriate for each route of administration.

- Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound is admixed with at least one inert pharmaceutically acceptable carrier such as sucrose, lactose, or starch. Such dosage forms can also comprise, as is normal practice, additional substances other than such inert diluents, e.g., lubricating agents such as magnesium stearate. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. Tablets and pills can additionally be prepared with enteric coatings.

- Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, solutions, suspensions, syrups, the elixirs containing inert diluents commonly used in the art, such as water. Besides such inert diluents, compositions can also include adjuvants, such as wetting agents, emulsifying and suspending agents, and sweetening, flavoring and perfuming agents.

- Preparations according to this invention for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, or emulsions. Examples of non-aqueous solvents or vehicles are propylene glycol, polyethylene glycol, vegetable oils, such as olive oil and corn oil, gelatin, and injectable organic esters such as ethyl oleate. Such dosage forms may also contain adjuvants such as preserving, wetting, emulsifying, and dispersing agents. They may be sterilized by, for example, filtration through a bacteria-retaining filter, by incorporating sterilizing agents into the compositions, by irradiating the compositions, or by heating the compositions. They can also be manufactured in the form of sterile solid compositions which can be dissolved

-22-

in sterile water, or some other sterile injectable medium immediately before use.

Compositions for rectal or vaginal administration are preferably suppositories which may contain, in addition to the active substance, excipients  
5 such as coca butter or a suppository wax.

Compositions for nasal or sublingual administration are also prepared with standard excipients well known in the art.


The dosage of active ingredient in the compositions of this invention may be varied; however, it is necessary that the amount of the active ingredient  
10 be such that a suitable dosage form is obtained. The selected dosage depends upon the desired therapeutic effect, on the route of administration, and on the duration of the treatment. Generally, dosage levels of between 0.0001 to 100 mg/kg of body weight daily are administered to patients and animals, e.g., mammals, to obtain effective release of growth hormone.

15 A preferred dosage range is 0.01 to 5.0 mg/kg of body weight daily.

The following structures prepared in the Examples indicated exemplify the nomenclature used in this document:

(R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-  
20 1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride  
**Example 26**

25






-23-

(R)-Piperidine-4-carboxylic acid [1-naphthalen-2-ylmethyl-2-oxo-2-[5-(toluene-4-sulfonylamino)-1,3-dihydro-indol-2-yl]-ethyl]-amide

**Example 83**

5


10



(R)-N-[1-[4-(Acetyl-phenyl-amino)-piperidine-1-carbonyl]-2-(1H-indol-3-yl)-ethyl]-2-amino-isobutyramide

**Example 34**

20




(R)-2-Amino-N-[1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-phenyl-4-(pyrrolidine-1-carbonyl)-piperidin-1-yl]-ethyl]-isobutyramide

**Example 46**

30

35



-24-


2-Amino-N-(1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-  
benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-isobutyramide

**Example 1**

5

10

15




(R)-2-Amino-N-(2-(1H-indol-3-yl)-1-[6-[2-(3-methyl-ureido)-phenyl]-3,4-  
dihydro-1H-isoquinoline-2-carbonyl]-ethyl)-isobutyramide

**Example 29**

20

25




-25-

(R)-3-Amino-N-[1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl]-3-methyl-butylamide

**Example 91**

5

10



The preparation of the compounds of Formula I of the present invention can be carried out in sequential or convergent synthetic routes. Syntheses detailing the preparation of the compounds of Formula I in a sequential manner are presented in the following reaction schemes.

Many protected amino acid derivatives are commercially available, where the protecting groups P are, for example, BOC, CBZ, Fmoc, benzyl or ethoxycarbonyl groups. Other protected amino acid derivatives can be prepared by literature methods. Some substituted pyrrolidines, piperidines and tetrahydroisoquinolines are commercially available, and many other pyrrolidines, 4-substituted piperidines, and 1,2,3,4-tetrahydroisoquinolines are known in the literature. Various phenyl or heteroaryl substituted piperidines and 1,2,3,4-tetrahydroisoquinolines can be prepared following literature methods using derivatized phenyl and heteroaryl intermediates. Alternatively the phenyl or heteroaryl rings of such compounds can be derivatized by standard means, such as halogenation, nitration, sulfonylation etc.

Many of the schemes illustrated below describe compounds which contain protecting groups P. Benzyloxycarbonyl groups can be removed by a number of methods including, catalytic hydrogenation with hydrogen in the presence of a palladium or platinum catalyst in a protic solvent such as methanol. Preferred catalysts are palladium hydroxide on carbon or palladium on carbon. Hydrogen pressures from 1-1000 psi may be employed; pressures from 10 to 70 psi are preferred. Alternatively, the benzyloxycarbonyl group can be removed by transfer hydrogenation.

Removal of BOC protecting groups can be carried out using a strong acid such as trifluoroacetic acid or hydrochloric acid with or without the


-26-

presence of a cosolvent such as dichloromethane or methanol at a temperature of about -30 to 70°C, preferably about -5 to about 35°C.

- Benzyl esters of amines can be removed by a number of methods including, catalytic hydrogenation with hydrogen in the presence of a
- 5 palladium catalyst in a protic solvent such as methanol. Hydrogen pressures from 1-1000 psi may be employed; pressures from 10 to 70 psi are preferred. The addition and removal of these and other protecting groups is discussed by T. Greene in Protective Groups in Organic Synthesis, John Wiley & Sons, New York, 1981.

-27-

## SCHEME 1



The protected amino acid derivatives 1 are in many cases commercially available, where the protecting group P is, for example, BOC or CBZ groups. Other amino acids can be prepared by literature methods.


-28-

As illustrated in Scheme 1, coupling of amines of formula 2 with protected amino acids of formula 1, where P is a suitable protecting group, is conveniently carried out in an inert solvent such as dichloromethane or DMF by a coupling reagent such as EDC or DCC in the presence of HOBT. In the case  
5 where the amine is present as the hydrochloride salt, it is preferable to add one equivalent of a suitable base such as triethylamine to the reaction mixture. Alternatively, the coupling can be effected with a coupling reagent such as BOP in an inert solvent such as methanol. Such coupling reactions are generally conducted at temperatures of about -30 to about 80°C, preferably 0  
10 to about 25°C. For a discussion of other conditions used for coupling peptides see Houben-Weyl, Vol. XV, part II, E. Wunsch, Ed., George Thieme Verlag, 1974, Stuttgart. Separation of unwanted side products and purification of intermediates is achieved by chromatography on silica gel, employing flash chromatography (W. C. Still, M. Kahn and A. Mitra, J. Org. Chem. 43 2923  
15 1978), by crystallization, or by trituration.

Transformation of 3 into intermediates of formula 4 can be carried out by removal of the protecting group P as described above. Coupling of intermediates of formula 4 to amino acids of formula 5 can be effected as described above to give intermediates of formula 6. Deprotection of the amine  
20 6 gives compounds of formula 7.

-29-

## SCHEME 2



25


Alternatively, compounds of formula 7 can be prepared by a convergent route as shown in Scheme 2. Intermediate esters of formula 8 can be prepared by treating amino acids 1, where P is a suitable protecting group, with a base such as potassium carbonate followed by an alkyl halide such as iodomethane in a suitable solvent such as DMF. Deprotection of the amine transforms 8 into 9. Alternatively, many amino acids of formula 9 are commercially available. Intermediate 10 is generated by coupling 9 to amino acid 5. The ester of intermediate 10 can be converted to intermediate acid 11 by a number of methods known in the art; for example, methyl and ethyl esters can be hydrolyzed with lithium hydroxide in a protic solvent such as aqueous methanol or aqueous THF at a temperature of about -20 to 120°C, preferably about 20 to 70°C. In addition, removal of a benzyl group can be accomplished by a number of reductive methods including hydrogenation in the presence of

-30-

platinum or palladium catalyst in a protic solvent such as methanol. Acid 11 can then be coupled to amine 2 to give intermediates of formula 6.

Transformation of 6 to 7 can be achieved by removal of the protecting group P.

## SCHEME 3




Compounds of formula 6 where  $R = CH_2COOR$ , can be converted to intermediate acids of formula 13 by any of the methods outlined in Scheme 2. Coupling the acid 13 to amine 147 generates the intermediates of formula 14 as illustrated in Scheme 3. Transformation of 14 to 15 can be achieved by removal of the protecting group P.



-31-


## SCHEME 4



As shown in Scheme 4, compounds of formula 18 can be prepared by treating 2-nitrobenzaldehydes of formula 16 with amines of formula 17 in the presence of a suitable reducing agent which include alkali metal borohydrides and cyanoborohydrides. The preferred reducing agent is sodium cyanoborohydride. Sodium borohydride and sodium triacetoxyborohydride may also be used. For a general review of reductive aminations see R. F. Borch, *Aldrichimica Acta*, 8, 3-10 (1975). The nitro group of compound 18 can be reduced by a number of methods including hydrogenation with a catalyst such as palladium in a protic solvent such as methanol to give compounds of formula 19. Cyclization of the diamine with *N,N'*-carbonyldiimidazole (CDI) or other phosgene equivalents generates compounds of formula 20. Removal of the protecting group transforms 20 into 12.

-32-

## SCHEME 5




As shown in Scheme 5, compounds of formula 23 can be prepared by treating 2-nitroanilines of formula 21, where  $X=N$  or  $C$ , with N-protected piperidones of formula 22 in the presence of a hydride reducing agent such as

-33-

sodium borohydride. Alternatively intermediates of formula 23 can be prepared by heating a protected 4-aminopiperidine 17 and a compound of formula 30, where X=N or C, in a solvent such as ethanol or DMF in the presence of an acid acceptor such as triethylamine or potassium carbonate.

- 5 The nitro group of formula 23 can be reduced by a number of methods including hydrogenation with a catalyst such as palladium in a solvent such as methanolic HCl to give compounds of formula 24. Cyclization with CDI or other phosgene equivalents produces the benzimidazolinones of formula 25. Transformation of 25 into 26 can be achieved by removal of the protecting
- 10 group P.

SCHEME 6




As shown in Scheme 6, the benzimidazolinones of formula 26 can also be prepared by treating 2-anthranilic acids of formula 27 with N-protected piperidones of formula 22 in the presence of a hydride reducing agent such as

-34-

sodium cyanoborohydride to give compounds of formula 28. The acid of formula 28 can then be treated with diphenylphosphoryl azide to give the aroyl azide of formula 29. Heating 29 in xylene produces the benzimidazolinones of formula 30. Transformation of 30 into 26 can be achieved by removal of the protecting group P.

SCHEME 7




-35-

Compounds of formula 33 can be prepared by treating a 4-piperidone of formula 22, where P is a protecting group such as benzyl, with a primary ( $R_2=H$ ) or secondary amine of formula 147 in the presence of a hydride reducing agent such as sodium cyanoborohydride as shown in Scheme 7.

- 5 Compounds of formula 33 can either be deprotected by conventional means to give compounds of formula 34, or in the case where  $R_{2a}=H$ , the amine can be coupled to a carboxylic acid of formula 35 using a coupling agent such as EDC in the presence of HOBT in an inert solvent such as dichloromethane to give compounds of formula 36. Subsequent removal of the protecting group  
10 generates compounds of formula 37. Alternatively, 33 can be treated with an aryl or alkyl sulfonyl chloride 38 in the presence of an acid scavenger to generate the sulfonamide 39. Transformation of 39 into 40 proceeds by removal of the protecting group.

SCHEME 8


15



- As illustrated in Scheme 8, piperidinols of formula 41 can be prepared  
25 by treating a suitably protected 4-piperidone of formula 22 with organometallics of formula  $YLi$  or  $YMgX$  or  $YZnX$  where  $X=Cl, Br, \text{ or } I$  in aprotic solvents such as tetrahydrofuran or diethyl ether at temperatures at or below  $0^\circ C$ . Intermediates of formula 41 can then be deprotected to give compounds of formula 42.


-36-

## SCHEME 9



- 20 Treating a methyl arylacetate 43 and a protected bis(chloroethyl)amine 44 (Bercz and Ice; J. Pharm. Sci. 1972 61 pp 1316-1317) with two equivalents of a base such as lithium diisopropylamide in a solvent such as THF generates the piperidine ester 45 as shown in Scheme 9. The ester can then be hydrolyzed under mild basic conditions to give intermediates of formula 46.
- 25 The resulting acid can then be coupled to amines of formula 147 to give 4,4'-disubstituted piperidines of formula 47. Deprotection of the amine generates compounds of formula 48.


## SCHEME 10



-37-

As illustrated in Scheme 10, compounds of formula 50 can be prepared by treating a 4-piperidone of formula 22, with equimolar amounts of potassium cyanide and a primary or secondary amine hydrochloride in aqueous ethanol to give intermediates of formula 49. Intermediates of formula 49 can then be  
 5 deprotected to give compounds of formula 50.


## SCHEME 11



15 Treating an aryl nitrile of formula 51 with intermediate 44 with two equivalents of a base such as sodium amide in a suitable solvent such as DMSO generates the 4,4'-disubstituted piperidines of formula 52. Subsequent deprotection of the piperidine amine gives rise to compounds of formula 53 as shown in Scheme 11.

20

## SCHEME 12




The nitrile group of 52 can be reduced by a number of methods (March, J., Advanced Org. Chem.: Reactions, Mechanisms, and Structure, John Wiley &

-38-

Sons, New York, 1985), including lithium aluminum hydride to give the primary amine of compound 54. Intermediate 54 can then be coupled to acids of formula 35 as outlined in Scheme 12 to give compounds of formula 55.

Subsequent deprotection of the piperidine amine gives compounds of formula 56.

## SCHEME 13






-39-

The nitrile 52 can also be converted to an ester as described in Scheme 13 by heating intermediate 52 with concentrated sulfuric acid and a small amount of water to temperatures around 150°C, followed by heating the mixture in an alcohol solvent to temperatures around 125°C. Deprotection of the piperidine amine 57 generates compounds of formula 58. Alternatively, the nitrile can be hydrolyzed to the amide 161 using a number of methods known in the literature (March, J., *Advanced Org. Chem.; Reactions, Mechanisms, and Structure*, p. 788, John Wiley & Sons, New York, 1985); for example, heating the amide in concentrated sulfuric acid to temperatures around 100-150°C will convert the nitrile to the amide as will treating the nitrile with hydrogen peroxide and sodium hydroxide in aqueous acetone. Transformation of 161 to 162 proceeds by removal of the protecting group.


SCHEME 14



Many 1,2,3,4-tetrahydroisoquinolines and substituted 1,2,3,4-tetrahydroisoquinolines can be prepared by methods described in the literature (R. Adams, *Org. Reactions* vol. VI; pp 74-206, John Wiley & Sons, New York, NY 1964). As illustrated in Scheme 14, treating an amine of formula 59, such as 1,2,3,4-tetrahydroisoquinoline, with potassium nitrate in concentrated sulfuric acid at temperatures below 5°C generates the nitro amine of formula 60. Reduction of the nitro functionality to the corresponding amine using iron powder and ammonium chloride in refluxing aqueous ethanol is one of many suitable literature procedures (see March, J., p. 1103-4, *Advanced Org. Chem.; Reactions, Mechanisms, and Structure*, John Wiley & Sons, New York, 1985) for carrying out this transformation to generate compounds of formula 61.

-40-


## SCHEME 15



- As described in Scheme 15, the  $\beta$ -phenethylamide of compound 63 can be prepared by treating 4-bromophenethylamine, 62, with ethyl formate.
- 20 Treating intermediate 63 with polyphosphoric acid and phosphorous pentoxide in a Bischler-Napieralski cyclodehydration produces the 7-bromo-3,4-dihydroisoquinoline 64. Sodium borohydride reduction of the hydrochloride salt of compound 64 in water produces the 7-bromo-1,2,3,4-tetrahydroisoquinoline 65. Following the method outlined in Scheme 14,
- 25 nitration of 7-bromo-1,2,3,4-tetrahydroisoquinoline generates intermediate 66. Protection of the tetrahydroisoquinoline amine 66 with a suitable protecting group P, such as tert-butoxycarbonyl, can be achieved by methods known to those skilled in the art and produces intermediates of formula 67. Reduction of the nitro group and dehalogenation can be accomplished by hydrogenation
- 30 using a catalyst such as palladium in a solvent such as acetic acid with ammonium acetate as a buffer and gives compounds of formula 68.


-41-

SCHEME 16



Treating the amines of formula 68 with an isocyanate of formula 69 gives the urea of formula 70 as illustrated in Scheme 16. Alternatively the amine 68 can be functionalized with an acid chloride or acid to give the amide of formula 72, a sulfonyl chloride of formula 38 to give the sulfonamide 74, or a ketone of formula 154 and a reducing agent to give the amine of formula 155. The resulting amines can be deprotected to give intermediates 71, 73, 75 and 156 respectively.


SCHEME 17



The synthesis of 6-, 7-, and 8-methoxy substituted 1,2,3,4-tetrahydroisoquinolines is described by Sall and Grunewald in J. Med. Chem. 30, 1987 pp 2208-2216. The methoxy group can be removed by a number of methods; for example, refluxing compounds of formula 76 with 48% HBr generates compounds of formula 77 as shown in Scheme 17.

-42-


## SCHEME 18



-43-

- Treating compound of formula 78, where P is a suitable protecting group, with pyridine and triflic anhydride in a suitable solvent such as dichloromethane generates the triflate of formula 79 as shown in Scheme 18.
- 5 The triflate can then be carbonylated with carbon monoxide and a catalyst such as palladium acetate in the presence of a base such as triethylamine and a ligand such as 1,3-bis(diphenylphosphine)propane in an alcohol solvent such as methanol to generate the ester of formula 80. The ester can then be hydrolyzed by a number of different methods to give the acid of formula 81.
  - 10 The acid can then be coupled to amines of formula 147 to give compounds of formula 82. Subsequent deprotection of the amine generates compounds of formula 83. Alternatively, compound of formula 80 can be deprotected to give intermediate 165. The amine can then be coupled to acids of formula 11 to give compounds of formula 166. Subsequent deprotection of the amine
  - 15 generates intermediate 167. Hydrolysis of the ester transforms 167 into 168.


SCHEME 19



- Alternatively, as shown in Scheme 19, the triflate of formula 79 can undergo a biaryl cross-coupling reaction with a suitable nucleophile such as an arylboronic acid or aryl zincate 86 (where M = metal) in the presence of a
- 35 catalyst such as tetrakis (triphenylphosphine) palladium(0) and potassium carbonate in toluene and aqueous ethanol to give the intermediate of formula 84. Deprotection of the isoquinoline amine transforms 84 to 85.

-44-


SCHEME 20



As shown in Scheme 20, the boronic acid of formula 88 can be prepared by treating a suitably protected aniline 87 with t-butyllithium and trimethylborate at temperatures near  $-78^\circ\text{C}$ . Intermediate 88 can then be coupled to triflate 79 using conditions defined in Scheme 19 to give 89. The aniline protecting group can then be selectively removed, and the resulting amine 90 treated with an isocyanate 69 and a suitable base to give the urea of formula 91. Deprotection of the amine transforms 91 to 92. Alternatively, the amine of

formula 90 can also be sulfonylated with a sulfonyl chloride in the presence of an acid scavenger to give compounds of formula 93. Deprotection of the amine transforms 93 to 94. Acylation of 90 with a carboxylic acid 35 using a coupling agent such as EDC in the presence of HOBT in an inert solvent such as DMF gives the compound of formula 95. Deprotection of the amine transforms 95 to 96.

**SCHEME 21**



-46-

As illustrated in Scheme 21, phthalimides of formula 99 can be prepared from 4-aminopyridines 97 and phthalic anhydride 98 as described by Ciganek et al. U.S. Pat. Appl. 92/876542. The pyridine ring of 99 can be reduced by a number of methods including hydrogenation in the presence of a platinum catalyst in a solvent such as methanolic HCl to give the piperidine ring of formula 157. Alternatively, quaternization of the pyridine ring with benzyl bromide in ethanol leads to the quaternary salt 100. The pyridine ring can then be reduced by a number of methods including hydrogenation in the presence of a platinum catalyst in glacial acetic acid or methanol to give 101. Many other anhydrides are commercially available and one skilled in the art can apply such chemistry to these compounds.

Addition of organometallics such as RLi or RMgX where X=Cl, Br, or I to compounds of formula 101 followed by hydrolysis produces compounds of formula 102. Deprotection of the piperidine amine transforms 102 to 103.

Compounds of formula 101 can be converted to compounds of formula 104 with hydride reducing agents such as sodium borohydride in methanol, or lithium borohydride in an aprotic solvent such as THF. Phthalimides may also be reduced to compounds of type 104 with zinc in acetic acid. Deprotection of the piperidine ring transforms 104 to 105.

Compounds of formula 104 can be treated with a base such as sodium hydride in appropriate solvents such as tetrahydrofuran, or metal alkoxides such as sodium methoxide in alcohol solvents such as methanol, followed by addition of an alkylating agent to give compounds of formula 106.


Alternatively, such compounds can be prepared by treating compounds of formula 104 with an alcohol in the presence of an acid such as hydrochloric or methanesulfonic at temperatures of about 0-100°C. Deprotection of the piperidine ring transforms 106 into 107.

Treating compounds of formula 101 or 104 with zinc in acetic acid or tin in acetic acid in the presence of hydrochloric acid generates intermediates of formula 108. Deprotection of the amine transforms 108 to 109.



-47-

## SCHEME 22




Treating 110 with one equivalent of a base such as potassium tert-butoxide in a solvent such as THF, followed by addition of 111 and then by a second equivalent of base generates intermediate of formula 112 as illustrated in Scheme 22. Conversion of 112 to 113 can be accomplished by a number of methods, including hydrogenation with a catalyst such as palladium in a mixture of ethanol and aqueous HCl. Alternatively, the ester of 112 can be deprotected to give the intermediate acid 158. The acid 158 can be coupled to

-48-

amine 147 to give the amide of formula 159. Conversion of 159 to 160 can be accomplished by a number of methods including hydrogenation with a catalyst such as platinum in methanolic HCl.

5

SCHEME 23



35


As illustrated in Scheme 23, a differentially protected 3-aminopyrrolidine of formula 114 is selectively deprotected at the 3-amino position to give intermediate 115. The amine can then be sulfonylated with a sulfonyl chloride

-49-

38 in the presence of an acid scavenger in a solvent such as dichloromethane to give 116. Alternatively, the amine can be coupled to acid 35 as above to give 117. Deprotection of the pyrrolidine nitrogen of 116 and 117 generates compounds of formula 118 and 119 respectively.

5


SCHEME 25



-50-

As shown in Scheme 25, isoindolines of formula 129 can be prepared from 126 by radical bromination using a suitable source of bromine radical such as N-bromosuccinimide and a suitable radical initiator such as AIBN (2,2'-azobis(isobutyronitrile) in an inert solvent such as carbon tetrachloride to produce intermediates 127. The dibromide can then be treated with a suitably protected primary amine such as benzylamine in aqueous acetone using a base such as sodium carbonate to give compounds of formula 128. Deprotection of the amine and nitro reduction to give 129 can be accomplished by hydrogenation with hydrogen in a protic solvent such as ethanol using a catalyst such as palladium. The coupling of intermediate 129 to acid 11 gives 130. The free amine can then be acylated or sulfonylated by methods mentioned above to give 131 and 132. Transformation of 131 into 133 and 132 into 134 can be accomplished by removal of the protecting group P.

SCHEME 26




As shown in Scheme 26, an N-protected 4-hydroxypiperidine 142 can be treated with methanesulfonyl chloride in the presence of a base such as triethylamine in an inert solvent such as methylene chloride to give the intermediate mesylate of formula 143. The mesylate can then be displaced with an imidazo[4,5-b]pyridine of formula 135 (for example, see Carpino et al., *Biorg. & Med. Chem. Lett.* 1994 4, pp 93-98 and references cited within) using

-51-

a base such as sodium hydride in a solvent such as DMF or dioxane to give 144. Deprotection of the amine transforms 144 into 145.

SCHEME 27


5



Benzisoxazoles of formula 136 can be prepared as described by Villabolos et al., J. Med. Chem. 1994 37 pp 2721-2724. The amine 136 can be coupled to acids of formula 35 as illustrated in Scheme 27 to give 30 intermediates 137. Deprotection of the piperidine amine gives 138. Alternatively the amine 136 can be sulfonlated to give 139. Deprotection of the amine generates compounds of formula 140.

-52-


## SCHEME 28



Esters of formula 146 can be prepared by treating an acid of formula 5  
 15 with hydroxysuccinimide in the presence of a coupling agent such as EDC in  
 an inert solvent such as methylene chloride as illustrated in Scheme 28.  
 Treating the ester with an amino acid of formula 1 in a solvent such as DMF in  
 the presence of a base such as diisopropylethylamine produces 11.

20


## SCHEME 29



-53-


The preparation of compounds of formula 148 has been described by Blank et al., J. Med. Chem. 1980 23 pp 837-840. Treating the sulfonyl chloride 148 with a phenol of formula 149 and a base such as potassium carbonate in a polar solvent such as acetonitrile or acetone generates intermediates of formula 150. Deprotection of the amine transforms 150 into 151. Alternatively the sulfonyl chloride can be treated with an amine 147 in the presence of an acid scavenger such as triethylamine in an inert solvent such as dichloromethane to give the sulfonamide of formula 152. Deprotection of the tetrahydroisoquinoline amine transforms 152 into 153.

SCHEME 30



As illustrated in Scheme 30, treating an indole of formula 169 with a base such as methoxide ion in a solvent such as methanol, followed by addition of a ketone of formula 170 produces intermediates of formula 171. Conversion of 171 to 172 can be accomplished by a number of reductive methods including hydrogenation in the presence of a protic solvent such as ethanol using a catalyst such as palladium.


SCHEME 31



-54-

As shown in Scheme 31, treating an amine of formula 24 with three equivalents of anhydride 173, neat, at temperatures near or around 160°C overnight, followed by decomposition of excess anhydride with a base such as sodium hydroxide generates compounds of formula 174. Deprotection of the amine transforms 174 into 190.

SCHEME 32




As illustrated in Scheme 32, alkylation of the diphenyloxazinone of formula 175 with cinnamyl bromide in the presence of sodium bis(trimethylsilyl)amide generates 176 which is then converted to the desired (D)-2-amino-5-phenylpentanoic acid 177 by removing the protecting group and hydrogenation over a  $\text{PdCl}_2$  catalyst.



-55-

SCHEME 33




Compounds of formula 179 can be prepared by treating an indole of formula 178 with benzoyl chloride in pyridine as illustrated in Scheme 33. Subsequent heating of 179 with dibenzylamine and a catalyst such as palladium on carbon at temperatures at or near 210°C in a solvent such as diphenyl ether generates the intermediate of formula 180. The indole nitrogen can be alkylated by treating 180 with cuprous bromide and bromobenzene in the presence of a base such as potassium carbonate in a suitable solvent such as N-methylpyrrolidine at reflux overnight to give intermediates of formula 181.

-56-

Quaternization of the pyridine nitrogen can be accomplished by treating 181 with benzyl bromide in a solvent such as benzene at refluxing temperatures to give intermediates of formula 192. Treating 192 with excess sodium borohydride in methanol generates the tetrahydropyridine 182. Hydrogenation of 182 with hydrogen in a solvent such as ethanol using a catalyst such as palladium produces the piperidine of formula 183.

## SCHEME 34




Compound of formula 185 is prepared by refluxing a mixture of 184 in acetic anhydride for several hours. The acid chloride can be prepared by treating 185 with either thionyl chloride or oxalyl chloride to give intermediates

-57-

- of formula 191. Friedel-Crafts acylation of m-difluorobenzene with 191 in the presence of aluminum chloride as a catalyst produces 186. Removal of the acetyl protecting group can be accomplished by heating 186 in a mixture of concentrated HCl and glacial acetic acid to give intermediate of formula 187.
- 5 The oxime 188 can then be prepared by refluxing 187 with triethylamine and hydroxylamine hydrochloride in ethanol for several hours. Refluxing the oxime in 50% NaOH produces the benzoisoxazole 189.

SCHEME 35



- As shown in Scheme 35, compounds of formula 193 can be prepared by treating anilines of formula 194 with N-protected piperidones of formula 22 in the presence of a hydride reducing agent such as sodium triacetoxyborohydride in a solvent such as acetic acid which contains an excess of sodium sulfate. Treating compounds of formula 193 with chloroacetonitrile at temperatures at or near 0°C in the presence of a Lewis acid such as boron trichloride followed by refluxing the mixture overnight and then treating the solution with 10% HCl and refluxing again for 0.5h produces intermediate of formula 195. Treating 195 with sodium borohydride in a solvent such as ethanol generates the indole of formula 196. Deprotection of the amine transforms 196 into 197.
- 30
- 35

-58-

The following examples are provided for the purpose of further illustration only and are not intended to be limitations on the disclosed invention.

#### General Experimental Procedures

- 5 Amicon silica 30  $\mu$ M, 60 Å pore size, was used for column chromatography. Melting points were taken on a Buchi 510 apparatus and are uncorrected. Proton and carbon NMR spectra were recorded on a Varian XL-300, Bruker AC-300, or Bruker AC-250 at 25 °C. Chemical shifts are expressed in parts per million downfield from trimethylsilane. Particle beam
- 10 mass spectra (MS) were obtained on a Hewlett-Packard 5989A spectrometer using ammonia as the source of chemical ionization. For initial sample dissolution chloroform or methanol were employed. Liquid secondary ion mass spectra (LSIMS) were obtained on a Kratos Concept-1S high resolution spectrometer using cesium ion bombardment on sample dissolved in a 1:5
- 15 mixture of dithioerythritol and dithiothreitol or in a thioglycerol matrix. For initial sample dissolution chloroform or methanol were employed. Reported data are sums of 3-20 scans calibrated against cesium iodide. TLC analyses were performed using E. Merck Kieselgel 60 F254 silica plates visualized (after elution with the indicated solvent(s)) by UV, iodine or by staining with 15%
- 20 ethanolic phosphomolybdic acid or ceric sulfate/ammonium molybdate and heating on a hot plate. The terms "concentrated" and "coevaporated" refer to removal of solvent at water aspirator pressure on a rotary evaporator with a bath temperature of less than 40°C.

- General Procedure A (Peptide coupling using EDC). A 0.2-0.5 M
- 25 solution of the primary amine (1.0 equivalent) in dichloromethane (or a primary amine hydrochloride and 1.0-1.3 equivalents of triethylamine) was treated sequentially with 1.0-1.2 equivalents of the carboxylic acid coupling partner, 1.5-1.8 equivalents hydroxybenzotriazole hydrate (HOBT), and 1.0-1.2 equivalents (stoichiometrically equivalent to the quantity of carboxylic acid) 1-
- 30 (3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) and the mixture was stirred overnight in an ice bath (the ice bath was allowed to warm, thus the reaction mixture was typically held at 0-20 °C for 4-6 h and 20-25 °C for the remaining period). The mixture was diluted with ethyl acetate or other solvent as specified, and the resulting mixture washed twice with 1N NaOH,
- 35 twice with 1N HCl (if the product is not basic), once with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated giving the crude product which was purified as specified. The carboxylic acid component could be used as the

-59-

dicyclohexylamine salt in coupling to the primary amine or hydrochloride of the latter; in this case no triethylamine was employed.

General Procedure B. (Cleavage of a t-Boc-protected Amine using TFA). Cold trifluoroacetic acid (usually 0-10°C) was added to the t-Boc amine (typically 10 mL per mmol amine) neat or dissolved in a minimum volume of CH<sub>2</sub>Cl<sub>2</sub> and the resulting solution was stirred at 0°C for 0.25-2h (the time required for complete disappearance of the starting material to a more polar product as judged by TLC). The resulting solution or suspension was concentrated, and the residue coevaporated several times with added methylene chloride. The residue was then dissolved in ethyl acetate and washed twice with 1N NaOH and once with brine. The organic phase was then dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated to give the free amine which was used without further purification or purified as specified.

General Procedure C. (Cleavage of a t-Boc-protected Amine and HCl salt exchange). Cold trifluoroacetic acid (usually 0-10°C) was added to the t-Boc amine (typically 10 mL per mmol amine) and the resulting solution was stirred at 0°C for 0.25-2 h (the time required for complete disappearance of the starting material to a more polar product as judged by TLC). The resulting solution or suspension was concentrated, the residue coevaporated several times with added methylene chloride and then dried in vacuo. The trifluoroacetate salt was dissolved in ethanol (typically 5 mL per mmol of salt) and cooled to 0°C. Two equivalents of either aqueous 1N HCl or 1M HCl in ether was added to the cold solution, and the mixture stirred for 10 min. at 0°C. The mixture was then concentrated and the residue coevaporated several times with ethanol and then dried in vacuo. The resulting oil in most cases could be triturated to a solid with diethyl ether.

General Procedure D. (Lithium Hydroxide Hydrolysis of Esters). To a 0.20-0.50M solution of ester dissolved in THF, 3.5 equivalents of lithium hydroxide hydrate dissolved in a volume of water equal to 25% the volume of THF was added. The mixture was stirred overnight at room temperature. Excess THF was removed by evaporation, and the basic aqueous mixture was extracted three times with ethyl acetate, and then acidified to pH4 with dilute acetic or hydrochloric acid. The product was extracted with ethyl acetate and the combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and evaporated to give the desired acid which was used without further purification, or triturated to a solid as specified.

-60-

**Example 1****2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride****5     A. {1-(R)-Benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-carbamic acid tert-butyl ester**

According to General Procedure A, 420 mg (1.94 mmol) of N-t-BOC-O-benzyl-D-serine and 571 mg (1.94 mmol) of 4-(2-keto-1-benzimidazoliny) piperidine were coupled and the product purified by silica gel chromatography (1:1 v/v EtO Ac/hexanes) to afford 630 mg of **1A**.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ 7.18-7.44 (m, 5H), 6.85-7.10 (m, 4H), 4.16-4.84 (m, 6H), 3.60-3.74 (m, 2H), 2.73-2.90 (m, 1H), 2.25-2.67 (m, 2H), 1.71-1.90 (m, 3H), 1.45-1.55 (m, 9H).

**15     B. (R)-1-[1-[(2-Amino-3-benzyloxy-propionyl)]-piperidin-4-yl]-1,3-dihydro-benzoimidazol-2-one**

According to General Procedure B, 630 mg (1.27 mmol) of the product from **1A** was deprotected to afford 550 mg of **1B**.

**20     C. {1-[1-(R)-Benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl]carbonyl}-1-methyl-ethyl}-carbamic acid tert-butyl ester**

According to General Procedure A, 44 mg (0.22 mmol) of N-t-BOC-α-methylalanine and 78 mg (0.20 mmol) of **1B** were coupled, and the product was purified by silica gel chromatography (10:1.5 v/v EtO Ac:hexanes) to give 24 mg of **1C**.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) δ 9.25-9.40 (d, 1H), 6.78-7.41 (m, 9H), 4.79-5.30 (m, 3H), 4.45-4.65 (m, 3H), 4.17-4.42 (m, 1H), 3.58-3.79 (m, 2H), 2.62-3.45 (m, 2H), 2.19-2.47 (m, 2H), 1.75-1.98 (m, 3H), 1.38-1.55 (m, 15H).

**30     D. 2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride**

The product from **1C** was deprotected according to General Procedure C to give 15 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ 7.39-7.41 (m, 7H), 6.93-7.15 (m, 4H), 5.12-5.22 (m, 1H), 4.42-4.76 (m, 5H), 4.20-4.30 (m, 1H), 3.70-3.86 (m, 2H), 3.28-3.25 (m, 1H), 2.22-2.49 (m, 1H), 1.75-1.92 (m, 2H), 1.53-1.61 (s, 6H).

35 MS (Cl, NH<sub>3</sub>) 480 (MH<sup>+</sup>)

**Example 2****3-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-3-methyl-butyramide hydrochloride**

-61-

A. (2-{1-(R)-Benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-1,1-dimethyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 78 mg (0.20 mmol) of **1B** and 48 mg (0.22 mmol) of 3-tert-Butoxycarbonylamino-3-methyl-butyric acid were coupled and the product purified by silica gel chromatography (10:1.5 v/v EtOAc/hexanes) to give 108 mg of **2A** as a clear oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) δ 7.19-7.38 (m, 5H), 6.79-7.17 (m, 5H), 5.22-5.52 (m, 2H), 4.82-4.97 (m, 1H), 4.49-4.63 (m, 3H), 4.20-4.38 (m, 1H), 3.60-3.72 (m, 2H), 3.03-3.32 (m, 1H), 2.50-2.82 (m, 3H), 2.11-2.49 (m, 3H), 1.77-1.98 (m, 2H), 1.35-1.53 (m, 15H).

B. 3-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-3-methyl-butyramide hydrochloride

According to General Procedure C, 94 mg (0.16 mmol) of **2A** was deprotected to give 58 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ 7.20-7.44 (m, 6H), 6.88-7.24 (m, 4H), 5.13-5.23 (m, 1H), 4.20-4.83 (m, 6H), 3.65-3.82 (m, 2H), 2.21-2.90 (m, 6H), 1.78-1.91 (m, 2H), 1.32-1.40 (bs, 6H).

MS (CI, NH<sub>3</sub>) 494 (MH<sup>+</sup>).

**Example 3**

(R)-3-Benzyloxy-2-(2-tert-butoxycarbonylamino-2-methyl-propionylamino)-propionic acid

To 1.83 g (6.2 mmol) of N-t-BOC-O-benzyl-D-serine in 35 mL of DMF was added 1.02 g (7.4 mmol) of potassium carbonate followed by 0.92 g (6.5 mmol) of iodomethane. The mixture was stirred overnight at 24°C under an atmosphere of nitrogen. The reaction mixture was diluted with 200 mL of water, and extracted three times with ethyl acetate. The combined organics were washed five times with water and once with brine, dried over MgSO<sub>4</sub> and concentrated. The crude (R)-3-Benzyloxy-2-tert-butoxycarbonylamino-propionic acid methyl ester was then deprotected according to General Procedure B and 0.84 g (4.02 mmol) of the resulting (R)-2-Amino-3-benzyloxy-propionic acid methyl ester was coupled to 0.81 g (4.02 mmol) of N-t-BOC-α-methylalanine to give 1.80 g of (R)-3-Benzyloxy-2-(2-tert-butoxycarbonylamino-2-methyl-propionylamino)-propionic acid methyl ester. The crude product was hydrolyzed according to the method outlined in General Procedure D, and 1.60 g of the title compound was recovered as an oil which solidified on standing.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 300 MHz) δ 7.30 (m, 5H), 7.10 (d, 1H), 5.07 (bs, 1H), 4.68 (m, 1H), 4.53 (q, 2H), 4.09 (m, 1H), 3.68 (m, 1H), 1.3-1.5 (m, 15H)

**Example 4****2-Amino-N-(1-(R)-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-isobutyramide****A. (R)-2-Amino-3-[(1H-indol-3-yl)-propionic acid methyl ester]**

5 To 4.92 g (16.2 mmol) of N- $\alpha$ -t-BOC-D-tryptophan in 100 mL of DMF was added 2.46 g (17.8 mmol) of potassium carbonate followed by 2.41 g (17.0 mmol) of iodomethane, and the mixture was stirred overnight at 24°C under an atmosphere of nitrogen. The reaction mixture was diluted with water, and extracted three times with ethyl acetate. The combined organics were washed  
10 five times with 500 mL of water and once with brine, dried over MgSO<sub>4</sub> and concentrated to give 4.67 g of a white solid. The crude (R)-2-tert-Butoxycarbonylamino-3-(1H-indol-3-yl)-propionic acid methyl ester was deprotected according to General Procedure B to give **4A** as an orange oil in quantitative yield.

**B. (R)-2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionic acid methyl ester.**

According to General Procedure A 1.55 g (7.1 mmol) of **4A** was coupled to 1.44 g (7.1 mmol) of N-t-BOC- $\alpha$ -methylalanine to give an oil which was purified by silica gel chromatography using a gradient of 10%, 20%, 30%, 40%  
20 and 50% ethyl acetate in hexane to afford 1.32 g of (R)-2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionic acid methyl ester.

**C. (R)-2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionic acid**

25 The product from **4B** was hydrolyzed according to the method described in General Procedure D to give 1.03 g of **4C** as an orange foam.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 300 MHz)  $\delta$  7.61 (d, 1H), 7.48 (d, 1H), 7.27 (t, 1H), 7.10 (t, 1H), 4.81 (bs, 1H), 3.35 (m, 1H), 1.49 (s, 6H), 1.32 (s, 9H).  
MS (CI, NH<sub>3</sub>) 390 (MH<sup>+</sup>)

**D. (1-(1-(R)-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester**

According to General Procedure A, 75 mg (0.19 mmol) of **4C** was coupled to 39 mg (0.18 mmol) of 4-(2-keto-1-benzimidazoliny)piperidine and  
35 the product was purified by silica gel chromatography (99:0.5 v/v CHCl<sub>3</sub>:MeOH) to give 90 mg of **4D** as a white foam.

**E. 2-Amino-N-(1-(R)-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-isobutyramide**



-63-

According to General Procedure B, 90 mg (0.15 mmol) of **4D** was deprotected to give 70 mg of the title compound as a yellow solid.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) (1:1 mixture of rotamers) δ 8.80 (bs, 0.5H), 8.58 (bs, 0.5H), 8.48-8.52 (dd, 1H), 7.83 (d, 1H), 7.52 (d, 1H), 7.34-7.40 (m, 1.5H), 6.93-7.12 (m, 7H), 6.65-6.71 (m, 1H), 4.70-4.88 (m, 1H), 4.39-4.45 (m, 0.5H), 4.08-4.39 (m, 0.5H), 3.72-3.92 (m, 1H), 2.80-2.99 (0.5H), 2.50-2.66 (m, 0.5H), 1.92-2.41 (m, 3H), 1.30-1.82 (m, 11H).  
MS (CI, NH<sub>3</sub>) 489 (MH<sup>+</sup>)

### Example 5

10 (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid methyl ester hydrochloride  
A. 6-Hydroxy-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester

To 4.75 g (20.7 mmol) of 1,2,3,4-Tetrahydro-isoquinolin-6-ol  
15 hydrobromide (see **7D**) in 150 mL of 1:1 water:dioxane at 0°C was added 4.95 g (22.7 mmol) of di-tert-butyl dicarbonate and sufficient 1N NaOH to maintain the pH of the solution between 10 and 10.5. The ice bath was removed after 30 min and the reaction was stirred at room temperature for an additional 2 h. The solution was concentrated, diluted with water and acidified with 1N HCl.  
20 The mixture was extracted three times with ethyl acetate. and the organic phase was washed with water and brine, dried over MgSO<sub>4</sub> and concentrated to give 5.24 g of crude product. The product was crystallized from hexane/ether to give 3.99 g of **5A** as a solid.

B. 6-(4-Trifluoro-methanesulfonyloxy)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester

To 3.97 g (15.94 mmol) of **5A** in 13 mL of methylene chloride was added 8.2 mL (101.4 mmol) of pyridine and the mixture was cooled to 0°C. A solution of 3.16 mL (18.81 mmol) of triflic anhydride in 26 mL of methylene chloride was then added dropwise to the stirring mixture over 20 min. The mixture was  
30 stirred for 30 min. at 0°C and then poured into saturated brine and extracted with ether. The ether phase was washed three times with 1N HCl, and once each with water and brine. The solution was dried over MgSO<sub>4</sub> and concentrated to give 6.08 g of **5B** as an oil which crystallized on standing.

C. 3,4-Dihydro-1H-isoquinoline-2,6-dicarboxylic acid 2-tert-butyl ester-6-methyl ester

A Parr bottle containing 381 mg (1.0 mmol) of **5B**, 6 mg (0.03 mmol) of palladium acetate, 25 mg (0.06 mmol) of 1,3-bis(diphenylphosphino)propane, 0.28 mL (2.0 mmol) of triethylamine in 4 mL of dimethylformamide and 1.8 mL

-64-

of methanol was charged to 5 psi with carbon monoxide, and the mixture agitated for 0.5 h. The mixture was then heated to 60°C and the pressure increased to 30 psi of carbon monoxide, and the mixture agitated overnight. After cooling to room temperature, the mixture was dissolved in ether and  
5 washed three times with water and once with brine. The solution was dried over MgSO<sub>4</sub> and concentrated to give 365 mg of 5C as an oil.

D. 1,2,3,4-Tetrahydro-isoquinoline-6-carboxylic acid methyl ester

According to General Procedure B, 563 mg (1.93 mmol) of 5C was deprotected to give 370 mg of 5D as an oil.

10 E. (R)-2-[2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid methyl ester

According to General Procedure A, 303 mg (1.59 mmol) of 5D was coupled to 617 mg (1.59 mmol) of 4C and the product was purified by silica  
15 gel chromatography (1:1 v/v ethyl acetate:hexanes) to give 866 mg of 5E.

F. (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid methyl ester hydrochloride

According to General Procedure C, 60 mg (0.107 mmol) of 5E was  
20 deprotected to give 41 mg of the title compound as a solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) δ (1:1 mixture of rotamers) 7.79 (d, 0.5H), 7.54-7.65 (m, 2.5H), 7.29 (d, 0.5H), 7.17 (d, 0.5H), 6.95-7.10 (m, 3.5H), 6.61 (d, 0.5H), 5.28-5.30 (m, 1H), 4.61-4.70 (d, 0.5H), 4.42-4.51 (m, 1H), 4.10-4.19 (d, 0.5H), 3.87 (d, 3H), 3.54-3.12 (m, 1H), 3.17-3.34 (m, 3H), 2.53-2.74 (m, 1H),  
25 2.36-2.51 (m, 0.5H), 2.08-2.21 (m, 0.5H), 1.56 (s, 6H).  
MS (Cl, NH<sub>3</sub>) 463 (MH<sup>+</sup>)

### Example 6

2-Amino-N-[1-(R)-benzyloxymethyl-2-[7-(3-methyl-ureido)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethyl]-isobutyramide

30 A. 7-Nitro-1,2,3,4-tetrahydro-isoquinoline

To 43.1 g (324 mmol) of 1,2,3,4-tetrahydro-isoquinoline was added 160 mL of concentrated sulfuric acid with ice bath cooling. Potassium nitrate 35.2 g (348 mmol) was added in portions to the stirring mixture, while maintaining an internal temperature below 5°C. The mixture was allowed to stand at ambient  
35 temperature for 72 h, and then basified with aqueous ammonia and extracted four times with chloroform. The combined organics were dried over MgSO<sub>4</sub> and concentrated to give a dark brown oil. The oil was dissolved in 240 mL of ethanol and 40 mL of concentrated hydrochloric acid was added to the mixture

-65-

with cooling. The precipitated material was collected by filtration and washed with cold ethanol to give 25.0 g of a tan solid. The crude material was partitioned between 1N NaOH and ethyl acetate. The ethyl acetate layer was washed once with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The dinitrated tetrahydroisoquinoline was removed by crystallization from ether/hexanes. The mother liquor was concentrated to give 2.19 g of 7-Nitro-1,2,3,4-tetrahydroisoquinoline.

B. {1-[1-(R)-Benzyloxymethyl-2-(7-nitro-3,4-dihydro-1H-isoquinolin-2-yl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 100 mg (0.57 mmol) of **6A** and 228 mg (0.60 mmol) of **3** were coupled, and the product was purified by silica gel chromatography (99:1 v/v CH<sub>2</sub>Cl<sub>2</sub>:MeOH) to give 220 mg of **6B** as a white foam.

C. {1-[2-(7-Amino-3,4-dihydro-1H-isoquinolin-2-yl)-1-(R)-benzyloxymethyl-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

To a solution of 222 mg (0.41 mmol) of **6B** in 2.5 mL of ethanol and 1.5 mL of water was added 110 mg (1.96 mmol) of iron powder, and 12 mg (0.22 mmol) of ammonium chloride. The mixture was refluxed for 45 min, and another 110 mg of iron powder, and 24 mg of ammonium chloride was added to the reaction mixture and refluxing continued for another 1 h. The hot solution was filtered through celite and concentrated. The residue was dissolved in ethyl acetate and washed once each with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (98:2 v/v CHCl<sub>3</sub>:MeOH) to give 147 mg of **6C** as a white foam. MS (Cl, NH<sub>3</sub>) 511 (MH<sup>+</sup>)

D. {1-[1-(R)-Benzyloxymethyl-2-[7-(3-methyl-ureido)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

To a solution of 68 mg (0.13 mmol) of **6C** in 3 mL of methylene chloride was added 0.40 mL (6.67 mmol) of methyl isocyanate, and the mixture was stirred for 17 h under nitrogen at 24°C. The mixture was concentrated and the product was purified by silica gel chromatography (98:2 v/v CHCl<sub>3</sub>:MeOH) to give 55 mg of the title compound as a white foam.

E. 2-Amino-N-{1-(R)-benzyloxymethyl-2-[7-(3-methyl-ureido)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethyl}-isobutyramide

According to General Procedure B, 55 mg (0.097 mmol) of **6D** was deprotected to give 41 mg of the title compound as a white foam.

-66-

<sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) (2:1 mixture of rotamers) δ 8.37-8.52 (m, 1H), 7.54 (bs, 0.67H), 7.41 (bs, 0.33H), 7.11-7.31 (m, 7H), 6.89-7.03 (m, 2H), 5.52-5.60 (m, 0.67H), 5.42-5.50 (m, 0.33H), 4.64 (bs, 1H), 4.59 (bs, 1H), 4.50 (d, 2H), 4.42 (bs, 1H), 3.61-3.90 (m, 4H), 2.62-2.77 (m, 4H), 1.55-1.72 (bs, 2H), 1.32 (s, 6H).  
 MS (CI, NH<sub>3</sub>) 468 (MH<sup>+</sup>)

### Example 7

#### 2-Amino-N-(1-(R)-benzyloxymethyl-2-(6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl)-2-oxo-ethyl)-isobutyramide

##### 10 A. Phenyl-carbamic acid tert-butyl ester

To a solution of 21.8 g (100 mmol) of di-tert-butyl-dicarbonate in 1 L of methylene chloride was added 10.0 g (100 mmol) of aniline. The mixture was stirred at 24°C overnight and then concentrated. The residue was dissolved in ether and washed ten times with dilute aqueous acetic acid, once each with 1N NaOH and brine, dried over MgSO<sub>4</sub> and concentrated to give 8.6 g of **7A** as a white solid.

##### B. 2-tert-Butoxycarbonylamino-phenylboronic acid

To 8.00 g (41.4 mmol) of Phenyl-carbamic acid tert-butyl ester in 103 mL of tetrahydrofuran at -78°C, 58.5 mL (99.5 mmol) of 1.7M tert-butyllithium in pentane was added dropwise. The mixture was warmed to -20°C and stirred at that temperature for 1 h after which time the reaction was cooled to -78°C, and 12.2 mL (107.7 mmol) of trimethylborate was added dropwise. The mixture was concentrated and 500 mL of ether and 125 mL of 1N HCl was added. The mixture was stirred at 24°C for 10 min. The layers were separated and the ethereal portion was washed once each with water and brine, dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (50:50 v/v EtO Ac:hexanes) to give 4.49 g of **7B** as a foam.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 300MHz) δ 9.16 (s, 1H), 8.85 (d, 1H), 7.37 (t, 1H), 7.06 (t, 1H), 1.48 (s, 9H).

##### 30 C. 6-Methoxy-1,2,3,4-tetrahydro-isoquinoline

To 20.0 g (132 mmol) of β-(m-methoxy) phenylethylamine was added 20.4 g (140 mmol) of 20% formaldehyde solution, and the mixture heated at 85°C for 1 h. After cooling to room temperature, the mixture was extracted with benzene. The organic phase was washed three times with water, once with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 26.8 g of an oil. To the resulting oil was added 26.0 g of a 20% aqueous hydrochloric acid solution, and the mixture was heated at 85°C for 45 min. The mixture was basified with 1N NaOH to pH 11 and extracted with ether. The ether layer was washed with

-67-

water and brine, dried over  $\text{MgSO}_4$ , and evaporated. The crude product was purified by silica gel chromatography using a gradient of 5%, 10%, and 20% methanol in chloroform to give 8.00 g of **7C** as an oil.

5 D. 6-Hydroxy-3,4-dihydro-1H-isoquinoline-2-carboxylic acid benzyl ester

A solution of 8.00 g (49 mmol) of 6-Methoxy-1,2,3,4-tetrahydro-isoquinoline in 196 mL of 48% hydrobromic acid was refluxed for 3 h. The mixture was then concentrated and coevaporated several times with ethanol. The resulting slurry was filtered and dried under vacuum to give 7.43 g of 1,2,3,4-Tetrahydro-isoquinolin-6-ol hydrobromide as a solid. To 7.38 g of the crude hydrobromide dissolved in 117 mL each of dioxane and water was added 6.34 g (35.3 mmol) of benzyl chloroformate with ice bath cooling. A 1M solution of potassium carbonate was slowly added until a pH of 10 was maintained. The mixture was stirred overnight at room temperature. Excess dioxane was removed by evaporation, and the resulting aqueous mixture was acidified with 1N HCl and the product extracted with ethyl acetate. The organic portion was washed with water and brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to give 10.4 g of an oil which was crystallized from ether/hexane to give 7.05 g of **7D** as a solid.

20 E. 6-(2-tert-Butoxycarbonylamino-phenyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid benzyl ester

To a solution of 4.00 g (14.1 mmol) of **7D** in 11 mL of methylene chloride was added 7.09 g (89.7 mmol) of pyridine. The mixture was cooled in an ice bath, and a of 4.70 g (16.7 mmol) of triflic anhydride was added to the stirring solution over 20 min. Once the addition was complete, the mixture was stirred at 0°C for 30 min. The reaction mixture was then poured into brine and the product extracted with ether. The ether phase was then washed three times with 1N HCl, once each with water and brine, dried over  $\text{MgSO}_4$  and evaporated to give the 6-(Trifluoro-methanesulfonyloxy)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid benzyl ester as an oil which crystallized on standing.

A mixture of 4.10 g (10.3 mmol) of crude 6-(Trifluoro-methanesulfonyloxy)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid benzyl ester, 3.25 g (13.7 mmol) of **7B**, 1.0 g of tetrakis(triphenylphosphine)palladium(0), 3.77 g (27.3 mmol) potassium carbonate, 45 mL toluene, 28 mL ethanol, and 18 mL of water were heated at 90°C overnight. The mixture was diluted with ethyl acetate and washed twice with saturated aqueous sodium bicarbonate, once with brine, dried over

-68-

MgSO<sub>4</sub> and concentrated to give an oil which was purified by silica gel chromatography (85:15 v/v hexanes:EtO Ac) to give 3.80 g of **7E**.

F. 6-(2-Amino-phenyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid benzyl ester

5 The product from **7E** 3.80 g (8.3 mmol) was deprotected according to General Procedure B to give 2.63 g of **7F** as a glass.

G. 6-[2-(3-Methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinoline-2-carboxylic acid benzyl ester

To a solution of 1.36 g (3.8 mmol) of **7F** in 45 mL of methylene chloride  
10 and 0.403 g (3.99 mmol) triethylamine was added a solution of 0.376 g (1.27 mmol) triphosgene in 8 mL of methylene chloride dropwise. The mixture was stirred at room temperature for 5 h. The mixture was purged with nitrogen for 25 min and then concentrated. The residue was dissolved in 21 mL of methanol and 15.8 mL of 40% aqueous methylamine was added to the stirring  
15 solution. The mixture was stirred at room temperature for 72 h and then concentrated. The mixture was diluted with water and extracted with ethyl acetate. The organic phase was washed with brine, dried over MgSO<sub>4</sub> and concentrated to give a foam which was crystallized from ethyl acetate/hexanes to give 0.711 g of **7G** as a white solid.

20 H. 1-Methyl-3-[2-(1,2,3,4-tetrahydro-isoquinolin-6-yl)-phenyl]-urea

To 0.70 g (1.69 mmol) of **7G** in 10 mL ethanol was added 0.350 g of 10% palladium on carbon and the mixture hydrogenated at 55 psi for 1.5 h. The catalyst was removed by filtration through celite, and the mixture concentrated. **7H** was obtained in quantitative yield and used in the following  
25 step without purification.

I. (1-(R)-Benzyloxymethyl-2-[6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 150 mg (0.53 mmol) of **7H** and 157 mg (0.53 mmol) of N-t-BOC-O-benzyl-D-serine were coupled. The product  
30 was purified by silica gel chromatography (60:40 v/v EtOAc:hexanes) to give 211 mg of **7I** as a foam.

J. 1-[2-[2-(2-Amino-3-benzyloxy-propionyl)-1,2,3,4-tetrahydro-isoquinolin-6-yl]-phenyl]-3-methyl-urea

Following the method outlined in General Procedure B, 205 mg (0.367  
35 mmol) of **7I** was deprotected to give 132 mg of **7J**.

K. [1-(1-(R)-Benzyloxymethyl-2-[6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethylcarbamoyl)-1-methyl-ethyl]-carbamic acid tert-butyl ester

-69-

According to General Procedure A, 130 mg (0.28 mmol) of 7J and 58 mg (0.28 mmol) of N-t-BOC- $\alpha$ -methylalanine were coupled and the resulting product was purified by silica gel chromatography (75:25 v/v EtOAc:hexanes) to give 142 mg of 7K as a foam.

5        L. 2-Amino-N-(1-(R)-benzyloxymethyl-2-(6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinolin-2-yl)-2-oxo-ethyl)-isobutyramide

According to General Procedure B, 142 mg (0.22 mmol) of 7K was deprotected to give 96 mg of the title compound as a white foam.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 300MHz)  $\delta$  8.32 (m, 1H), 7.86 (m, 1H), 7.01-7.45 (m, 12H), 6.10-6.33 (m, 1H), 5.09-5.27 (m, 1H), 4.43-4.84 (m, 5H), 3.60-3.85 (m, 3H), 2.62-2.96 (m, 5H), 1.32 (bs, 6H).  
10        MS (CI, NH<sub>3</sub>) 544 (MH<sup>+</sup>)

### Example 8

15        (R)-N-[3-(2-{1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-piperidin-4-yl}-ethyl)-benzo[d]isoxazol-6-yl]-benzamide

A. N-[3-(2-Piperidin-4-yl-ethyl)-benzo[d]isoxazol-6-yl]-benzamide  
4-[2-(6-Benzoylamino-benzo[d]isoxazol-3-yl)-ethyl]-piperidine-1-carboxylic acid tert-butyl ester was prepared as described by Villabolos et al., in J. Med. Chem. 1994 37 p2721-24 (example 2h). According to General  
20        Procedure B, 200 mg (0.44 mmol) of this material was then deprotected to give 8A in quantitative yield.

B. [1-(2-[4-[2-(6-Benzoylamino-benzo[d]isoxazol-3-yl)-ethyl]-piperidin-1-yl]-1-(R)-benzyloxymethyl-2-oxo-ethylcarbonyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

25        Following the method outlined in General Procedure A, 55 mg (0.16 mmol) of 8A and 60 mg (0.16 mmol) of 3 were coupled and the product was purified by silica gel chromatography (60:40 v/v ethyl acetate:hexanes) to give 8B as a foam.

30        C. (R)-N-[3-(2-{1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-piperidin-4-yl}-ethyl)-benzo[d]isoxazol-6-yl]-benzamide

According to General Procedure B, 29 mg (0.04 mmol) of 8B was deprotected to give 20 mg of the title compound as a foam.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) (1:1 mixture of rotamers)  $\delta$  8.19-8.31 (m, 2H),  
35        8.09 (m, 1H), 7.93 (d, 2H) 7.38-7.52 (m, 5H), 7.20-7.38 (m, 4H), 5.01-5.16 (m, 1H), 4.54-4.67 (m, 1H), 4.42-4.53 (q, 2H), 3.90-4.03 (m, 1H), 3.54-3.68 (m, 2H), 2.84-3.01 (m, 3H), 2.37-2.70 (m, 6H), 1.49-1.88 (m, 5H), 1.40 (s, 6H).

-70-

**Example 9**

N-[2-(7-Acetylamino-3,4-dihydro-1H-isoquinolin-2-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl]-2-amino-isobutyramide

5     A. {1-[2-(7-Acetylamino-3,4-dihydro-1H-isoquinolin-2-yl)-1-(R)-benzyloxymethyl-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

To a mixture of 75 mg (0.15 mmol) of 6C, and 36 mg (0.44 mmol) of sodium acetate in 2 mL of methylene chloride was added 12 mL (0.16 mmol) of methanesulfonyl chloride and the resulting mixture was stirred overnight at  
10   24°C. The mixture was diluted with 20 mL of ethyl acetate, and washed three times with 1N NaOH, and once each with 1N HCl and brine. The solution was dried over MgSO<sub>4</sub> and concentrated to give 52 mg of 9A as a clear oil.

B. N-[2-(7-Acetylamino-3,4-dihydro-1H-isoquinolin-2-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl]-2-amino-isobutyramide

15   According to General Procedure B, 52 mg (0.091 mmol) of 9A was deprotected and 26 mg of the title compound was obtained as a clear oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) (2:1 mixture of rotamers) δ 8.33 (d, 1H), 7.65 (s, 0.67H), 7.56 (s, 0.33H), 7.15-7.38 (m, 7H), 6.99-7.19 (m, 1H), 5.10-5.23 (m, 1H), 4.65-4.72 (m, 2H), 4.42-4.50 (m, 2H), 3.55-3.87 (m, 5H), 2.65-2.82 (m,  
20   2H), 2.16 (s, 3H), 1.34 (s, 6H).  
MS (CI, NH<sub>3</sub>) 453 (MH<sup>+</sup>)

**Example 10**

2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-1,4-dihydro-2H-quinazolin-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

25   A. {1-[1-(R)-Benzyloxymethyl-2-oxo-2-[4-(2-oxo-1,4-dihydro-2H-quinazolin-3-yl)-piperidin-1-yl]-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

Following the method outlined in General Procedure A, 14 mg (0.052 mmol) of 3,4-Dihydro-1H-quinazolin-2-one and 21 mg (0.055 mmol) of 3 were  
30   coupled and the product was purified by silica gel chromatography (99:0.75 v/v CHCl<sub>3</sub>:MeOH) to give 27 mg of 10A as a clear oil.

B. 2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-1,4-dihydro-2H-quinazolin-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

According to General Procedure C, 27 mg (0.045 mmol) of 10A was  
35   deprotected to give 21 mg of the title compound as a yellow solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ 7.20-7.41 (m, 6H), 7.03-7.20 (m, 2H), 6.73-7.01 (m, 1H), 5.11-5.22 (m, 1H), 3.98-4.79 (m, 8H), 3.70-3.84 (m, 2H), 3.10-3.26 (m, 1H), 2.61-2.90 (m, 1H), 1.85-2.32 (m, 2H), 1.55-1.78 (m, 9H).



MS (CI, NH<sub>3</sub>) 494 (MH<sup>+</sup>)

### Example 11

2-Amino-N-{1-(R)-benzyloxymethyl-2-[6-(2-methanesulfonylamino-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

A. 6-(2-Methanesulfonylamino-phenyl)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid benzyl ester

To 249 mg (0.695 mmol) of **7F** and 84 mg (0.083 mmol) of triethylamine in 3.5 mL of methylene chloride was added 87 mg (0.765 mmol) of methanesulfonyl chloride and the mixture stirred overnight at 24°C. An additional 16 mg of triethylamine and 18 mg of methanesulfonyl chloride were added, and the mixture stirred for an additional 6 h. The reaction mixture was poured into an aqueous sodium bicarbonate solution, and the product extracted twice with methylene chloride. The combined organics were dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (2:1 v/v hexanes:EtOAc) to give 291 mg of **11A** as an oil.

**B. N-[2-(1,2,3,4-Tetrahydro-isoquinolin-6-yl)-phenyl]-methanesulfonamide**

A mixture of 280 mg (0.642 mmol) of **11A** and 160 mg of 10% palladium on carbon in 20 mL of ethanol was hydrogenated at 50 psi for 2 h. An additional 100 mg of catalyst was added and the hydrogenation continued for an additional 1.5 h. The catalyst was removed by filtration through celite, and the solution concentrated to give 120 mg of a white solid which was recrystallized from methanol/hexane to give 61 mg of **11B** as a white solid.

25 C. (1-{1-(R)-Benzyloxymethyl-2-[6-(2-methanesulfonylamino-phenyl)-  
3,4-dihydro-1H-isquinolin-2-yl]-2-oxo-ethylcarbamoyl}-1-methyl-ethyl)-  
 carbamic acid tert-butyl ester

According to the method outlined in General Procedure A, 55 mg (0.18 mmol) of **11B**, and 68 mg (0.18 mmol) of **3** were coupled and the product was  
30 purified by silica gel chromatography (1:1 v/v hexanes:ethyl acetate) to give 15 mg of **11C** as a foam.

D. 2-Amino-N-[1-(R)-benzyloxymethyl-2-[6-(2-methanesulfonylamino-phenyl)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethyl]-isobutyrarnide hydrochloride

35 According to General Procedure C, 15 mg (0.02 mmol) of **11C** was deprotected to give 10 mg of the title compound as a white solid.

-72-

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (1:1 mixture of rotamers) δ 8.28 (d, 1H), 7.14-7.52 (m, 12H), 5.14-5.37 (m, 1H), 4.69-4.82 (m, 3H), 4.46-4.56 (m, 2H), 3.70-3.93 (m, 4H), 2.75-3.06 (m, 2H), 2.72 (s, 3H), 1.52 (bs, 6H).  
MS (CI, NH<sub>3</sub>) 565 (MH<sup>+</sup>)

5

**Example 12**

2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(phenyl-propionyl-amino)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

A. (1-{1-(R)-Benzyloxymethyl-2-oxo-2-[4-(phenyl-propionyl-amino)-piperidin-1-yl]-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

10

According to the method outlined in General Procedure A, 30 mg (0.13 mmol) of 4-(N-propionylanilino)-piperidine and 48 mg (0.13 mmol) of **3** were coupled and the product was purified by silica gel chromatography (1:1 v/v hexanes:ethyl acetate) to give 67 mg of **12A** as an oil.

B. 2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(phenyl-propionyl-amino)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

15

According to General Procedure C, 67 mg (0.11 mmol) of **12A** was deprotected to give 50 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ 7.11-7.47 (m, 12H), 5.00-5.10 (m, 1H), 4.63-4.81 (m, 2H), 4.49-4.62 (m, 2H), 4.27-4.38 (m, 1H), 3.94-4.21 (m, 1H), 3.45-3.68 (m, 2H), 2.99-3.23 (m, 1H), 2.62-2.80 (m, 1H), 1.81-2.00 (m, 4H), 1.52 (s, 6H), 1.07-1.38 (m, 3H), 0.97 (t, 3H).  
MS (CI, NH<sub>3</sub>) 495 (MH<sup>+</sup>)

20

**Example 13**

(R)-N-{2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-2,3-dihydro-1H-isoindol-5-yl}-benzamide hydrochloride

25

A. (R)-{1-[1-(5-Amino-1,3-dihydro-isoindole-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 330 mg (2.46 mmol) of **43C** and 960 mg (2.46 mmol) of **4C** were coupled and 1.10 g of **13A** was recovered as a yellow solid.

30

B. (R)-{1-[1-(5-Benzoylamino-1,3-dihydro-isoindole-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

A mixture of 300 mg (0.59 mmol) of **13A**, 98 mg (0.69 mmol) of benzoyl chloride and 85 mg (0.69 mmol) of 4-dimethylaminopyridine in 5 mL of methylene chloride was stirred overnight at room temperature. The mixture was diluted with chloroform and washed twice each with 10% hydrochloric acid, saturated aqueous sodium bicarbonate and brine. The solution was

35

-73-

dried over MgSO<sub>4</sub> and concentrated and the product was purified by silica gel chromatography using a gradient of 75% ethyl acetate in hexane to 100% ethyl acetate to give 60 mg of **13B** as a white solid.

5     C. (R)-N-{2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-2,3-dihydro-1H-isoindol-5-yl}-benzamide hydrochloride

To 60 mg (0.10 mmol) of **13B** in 2 mL of ethanol was added 2 mL of concentrated hydrochloric acid, and the mixture was stirred at room temperature for 1 h. The mixture was concentrated and dissolved in a trace amount of methanol. Methylene chloride was added until the product  
10   precipitated from solution giving 50 mg of **13C** as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) δ 10.12 (d, 1H), 8.45 (d, 1H), 7.93 (d, 2H), 7.49-7.72 (m, 7H), 7.04-7.35 (m, 5H), 4.98-5.08 (m, 1H), 4.65-4.81 (m, 1H), 4.46-4.60 (m, 1H), 3.34-3.52 (m, 2H), 1.62 (s, 6H).

LSIMS-MS 533 (M<sup>+</sup>Na)

15

#### Example 14

N-{2-[4-(Acetylamino-methyl)-4-phenyl-piperidin-1-yl]-1-(R)-benzyloxymethyl-2-oxo-ethyl}-2-amino-isobutyramide

20   A. (1-{2-[4-(Acetylamino-methyl)-4-phenyl-piperidin-1-yl]-1-(R)-benzyloxymethyl-2-oxo-ethyl}carbamoyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to the method outlined in General Procedure A, 46 mg (0.21 mmol) of N-(4-Phenyl-piperidin-4-ylmethyl)-acetamide and 85 mg (0.22 mmol) of **3** were coupled, and the product was purified by silica gel chromatography (95:5 v/v chloroform:methanol) to give 111 mg of **14A** as a clear oil.

25   B. N-{2-[4-(Acetylamino-methyl)-4-phenyl-piperidin-1-yl]-1-(R)-benzyloxymethyl-2-oxo-ethyl}-2-amino-isobutyramide

According to General Procedure B, 110 mg (0.18 mmol) of **14A** was deprotected to give 72 mg of the title compound as a white foam.

30   <sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) (1:1 mixture of rotamers) δ 7.16-7.40 (m, 10H), 5.92-6.03 (t, 0.5H), 5.61-5.68 (t, 0.5H), 4.93-5.04 (m, 1H), 4.43-4.58 (q, 1H), 4.30-4.43 (q, 1H), 3.84-4.03 (m, 1H), 3.46-3.78 (m, 3H), 3.00-3.42 (m, 6H), 1.87-2.17 (m, 2H), 1.83 (d, 3H), 1.61-1.82 (m, 1H), 1.28 (d, 3H), 1.24 (d, 3H).  
MS (CI, NH<sub>3</sub>) 495 (MH<sup>+</sup>)

-74-

**Example 15**

(R)-N-[2-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl]-benzamide hydrochloride

- 5 A. [1-[2-(7-Benzoylamino-3,4-dihydro-1H-isoquinolin-2-yl)-1-(R)-benzyloxymethyl-2-oxo-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

To a mixture of 50 mg (0.10 mmol) of **6C** in 1 mL of methylene chloride was added 14 mg (0.14 mmol) of triethylamine. The mixture was cooled to 0°C, and 17 mg (0.12 mmol) of benzoyl chloride was added. The stirring  
10 mixture was allowed to warm to room temperature over 1.5 h, and the mixture was then poured into a saturated solution of sodium bicarbonate and extracted with ethyl acetate. The organic phase was washed once with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified by silica gel  
15 chromatography using 100% chloroform followed by 5% methanol in chloroform to elute. Recovered 48 mg of **15A** as a foam.

B. (R)-N-[2-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl]-benzamide hydrochloride

According to General Procedure C, 48 mg (0.08 mmol) of **15A** was deprotected, to give 44 mg of the title compound as a white solid.

- 20 <sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ 7.78-7.92 (m, 2H), 7.41-7.22 (m, 5H), 7.04-7.38 (m, 8H), 5.22-5.40 (m, 1H), 4.41-4.72 (m, 4H), 3.62-3.90 (m, 5H), 2.71-2.90 (m, 2H), 1.60 (bs, 6H).  
MS (CI, NH<sub>3</sub>) 515 (MH<sup>+</sup>)

**Example 16**

- 25 2-Amino-N-[1-(R)-benzyloxymethyl-2-[4-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-piperidin-1-yl]-2-oxo-ethyl]-isobutyramide

A. 2-Piperidin-4-ylmethyl-isoindole-1,3-dione hydrochloride

- To a mixture of 148.0 g (1.0 mol) of phthalic anhydride and 15 mL of triethylamine in 1.5L of xylene was slowly added 108.0 g (1.0 mol) of 4-  
30 (Aminomethyl)pyridine with mechanical stirring. The resulting solid was collected by filtration. The N-Pyridin-4-ylmethyl-phthalamide was then cyclized, by heating 235 g (0.88 mol) to the melt stage for 15 min. The mixture was allowed to cool slightly and 2.5 L of ethanol was added to the warm reaction mixture, and the solution filtered. Upon cooling, the mixture  
35 crystallized to give 188.0 g of 2-Pyridin-4-ylmethyl-isoindole-1,3-dione (m.p. 166-167.5°C). A mixture of 23.8 g (0.1 mol) of 2-Pyridin-4-ylmethyl-isoindole-1,3-dione and 1.3 g of platinum oxide in 400 mL of methanolic HCl was reduced at 50 psi until no starting material could be detected by TLC. The

-75-

mixture was filtered through celite and concentrated to give 22.0 g of 2-Piperidin-4-ylmethyl-isoindole-1,3-dione hydrochloride as a white solid. A small sample was recrystallized from ethanol to give an analytically pure sample of **16A** (m.p. 241-242.5°C).

- 5        B. (1-{1-(R)-Benzyloxymethyl-2-[4-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-piperidin-1-yl]-2-oxo-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to the method outlined in General Procedure A, 83 mg (0.21 mmol) of **3** and 53 mg (0.21 mmol) of 2-Piperidin-4-ylmethyl-isoindole-1,3-dione hydrochloride were coupled, and the product was purified by silica gel chromatography (95:5 v/v chloroform:methanol) to give 14 mg of **16B** as a clear oil.

- 10       C. 2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(1,3-dioxo-1,3-dihydro-isoindol-2-ylmethyl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide

15       According to General Procedure B, 10 mg (0.016 mmol) of **16B** was deprotected to give 7 mg of the title compound as a clear oil.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (1:1 mixture of rotamers) δ 7.75-7.89 (m, 4H), 7.20-7.39 (m, 6H), 4.98-5.09 (m, 1H), 4.41-4.59 (m, 4H), 3.94-4.07 (m, 1H), 3.60-3.78 (m, 2H), 3.57-3.60 (d, 1H), 2.96-3.12 (m, 1H), 2.57-2.74 (m, 1H), 2.00-2.12 (bm, 1H), 1.60-1.78 (m, 2H), 1.26-1.37 (m, 8H).  
MS (CI, NH<sub>3</sub>) 507 (MH<sup>+</sup>)

### Example 17

2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide

- 25       A. (1-{1-(R)-Benzyloxymethyl-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 26 mg (0.068 mmol) of Morpholin-4-yl-(4-phenyl-piperidin-4-yl)-methanone hydrochloride and 20 mg (0.065 mmol) of **3** were coupled and the product was purified by silica gel chromatography (99:1 v/v chloroform:methanol) to give 42 mg of **17A** as a clear oil.

- B. 2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide

35       According to General Procedure B, 42 mg (0.66 mmol) of **17A** was deprotected to give 27 mg of the title compound as a clear oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) δ 7.13-7.39 (m, 10H), 6.97-7.02 (d, 1H), 5.00-5.12 (m, 1H), 4.38-4.52 (m, 3H), 3.82-4.00 (m, 1H), 3.51-3.67 (m, 3H), 2.96-3.50 (m, 9H), 2.20-2.34 (m, 1H), 1.57-2.11 (bm, 3H), 1.29 (s, 6H).

-76-

MS (Cl, NH<sub>3</sub>) 537 (MH<sup>+</sup>)**Example 18**

(R)-1'-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-  
[1,4']bipiperidiny-4'-carboxylic acid amide hydrochloride

5     A. {1-[1-(R)-Benzyloxymethyl-2-(4'-carbamoyl-[1,4']bipiperidiny-1'-yl)-2-  
oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to the method outlined in General Procedure A, 21 mg (0.055 mmol) of **3** and 11 mg (0.052 mmol) of 4-carbamyl-4-piperidinopiperidine were coupled and the product was purified by silica gel chromatography (99:0.75 v/v  
10 chloroform:methanol) to give 25 mg of **18A** as a glass.

B. (R)-1'-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-  
[1,4']bipiperidiny-4'-carboxylic acid amide hydrochloride

According to General Procedure C, 25 mg (0.044 mmol) of **18A** was deprotected to give 19 mg of the title compound as a white solid.

15     <sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (1:1 mixture of rotamers) δ 8.32 (d, 0.5H), 8.27 (d, 0.5H), 7.26-7.42 (m, 5H), 5.06-5.19 (m, 1H), 4.48-4.70 (m, 3H), 4.21-4.34 (m, 1H), 3.64-3.79 (m, 3H), 3.38-3.50 (m, 1H), 2.91 3.18 (m, 2H), 2.48-2.91 (m, 4H), 1.76-2.12 (m, 8H), 1.57 (s, 6H).

MS (Cl, NH<sub>3</sub>) 474 (MH<sup>+</sup>)

20

**Example 19**

(R)-2-Amino-N-[2-(5-diphenylacetyl-amino-1,3-dihydro-isoindol-2-yl)-1-(1H-  
indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide hydrochloride

A. (R)-{1-[2-(5-Diphenylacetyl-amino-1,3-dihydro-isoindol-2-yl)-1-(1H-  
indol-3-ylmethyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid

25     A mixture of 300 mg (0.59 mmol) of **13A**, 160 mg (0.68 mmol) of diphenylacetylchloride and 83 mg (0.68 mmol) of 4-dimethylaminopyridine in 5 mL of methylene chloride was stirred overnight at room temperature. The mixture was diluted with chloroform and washed twice each with 10% hydrochloric acid, saturated aqueous sodium bicarbonate and brine. The  
30 solution was dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography using a gradient of 75% ethyl acetate in hexane to 100% ethyl acetate to give 150 mg of **19A** as a white solid.

B. (R)-2-Amino-N-[2-(5-diphenylacetyl-amino-1,3-dihydro-isoindol-2-yl)-  
1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide hydrochloride

35     To 150 mg (0.20 mmol) of **19B** in 2 mL of ethanol was added 2 mL of concentrated hydrochloric acid, and the mixture was stirred at room temperature for 1 h. The mixture was concentrated to give 140 mg of the title compound as a white solid.

-77-

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture rotamers) δ 7.60 (d, 1H), 7.52 (s, 0.5H), 7.41 (s, 0.5H), 7.21-7.34 (m, 12H), 6.93-7.13 (m, 4H), 5.13 (s, 1H), 4.92-5.00 (m, 1H), 4.06-4.18 (m, 1H), 4.40-4.51 (m, 1H), 4.01-4.19 (m, 2H), 3.20-3.42 (m, 2H), 1.59 (s, 3H), 1.51 (s, 3H).

5 MS (CI, NH<sub>3</sub>) 600 (MH<sup>+</sup>)

### Example 20

(R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[6-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide hydrochloride

10 A. 6-(Toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert butyl ester

To a mixture of 275 mg (1.10 mmol) of 6-amino-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester, 134 mg (1.33 mmol) of triethylamine in 6 mL of methylene chloride at 0°C was added 233 mg (1.2 mmol) of toluenesulfonyl chloride in one portion. The mixture was allowed to  
15 warm to room temperature and was stirred overnight. An additional 62 mg (0.61 mmol) of triethylamine and 105 mg (0.55 mmol) of toluenesulfonylchloride was added and stirring was continued for an additional 4 h. The mixture was concentrated and the residue dissolved in ethyl acetate. The organic portion was washed twice each with 1N NaOH and brine, dried  
20 over Na<sub>2</sub>SO<sub>4</sub> and the product was purified by silica gel chromatography (1:3 v/v ethyl acetate:hexane) to give 274 mg of **20A** as a foam.

B. 4-Methyl-N-(1,2,3,4-tetrahydro-isoquinolin-6-yl)-benzenesulfonamide

According to General Procedure B, 253 mg (0.63 mmol) of **20A** was deprotected to give 175 mg of **20B** as a foam.

25 C. (R)-(1-(2-(1H-Indol-3-yl)-1-[6-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethylcarbamoyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 84 mg (0.278 mmol) of **20B** was coupled to 108 mg (0.278 mmol) of **4C** and the product was purified by silica  
30 gel chromatography (70:30 v/v ethyl acetate:hexane) to give 114 mg of **20C**.

D. (R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[6-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide hydrochloride

According to General Procedure C, 114 mg (0.17 mmol) of **20C** was deprotected to give 68 mg of the title compound as a solid.

35 <sup>1</sup>H NMR (CDCl<sub>3</sub> 250MHz) (mixture of rotamers) δ 8.41 (d, 0.5H), 8.33 (d, 0.5H), 8.15 (bs 0.5H), 7.75 (d, 2H), 7.67 (d, 2H), 7.53 (bs, 0.5H), 6.51-7.31 (m, 8.5H), 6.07 (d, 0.5H), 5.13-5.32 (m, 1H), 4.38-4.61 (q, 1H), 4.04-4.19 (m, 1H), 3.77-4.04 (m, 1H), 2.98-3.44 (m, 3H), 2.46-2.62 (m, 5H), 1.30 (bs, 6H).

-78-

**Example 21**

(R)-Piperidine-4-carboxylic acid {1-benzyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

5 A. (R)-4-(1-Carboxy-2-phenyl-ethylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester

A mixture of 1.00 g (6.05 mmol) of D-phenylalanine, 1.98 g (6.05 mmol) of **77A** and 1.84 g (18.2 mmol) of triethylamine in 10 mL of water and 40 mL of dioxane was stirred at room temperature for 15 h. The mixture was diluted with chloroform and acidified to pH 4 with acetic acid. The layers were separated  
10 and the organic portion was washed three times with brine, dried over MgSO<sub>4</sub> and concentrated to give 2.16 g of **21A** as a white solid.

B. (R)-4-{1-Benzyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 80 mg (0.29 mmol) of **50D** and 109  
15 mg of **96A** were coupled and the product was purified by silica gel chromatography (75:25 v/v ethyl acetate:hexanes) to give 140 mg of **21B**.

C. (R)-Piperidine-4-carboxylic acid {1-benzyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

To a solution of 140 mg (0.22 mmol) of **21B** in 5 mL of ethanol was  
20 added 5 mL of concentrated HCl and the mixture was stirred at room temperature for 1 h. The mixture was concentrated and the residue crystallized from methanol/ethyl acetate to give 107 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 8.32 (m, 0.5H), 7.66-7.89 (m, 8.5H), 7.19-7.40 (m, 5H), 5.13-5.21 (m, 0.33H), 5.02-5.10 (m, 0.67H), 4.15-4.27 (m, 0.33H), 4.03-4.14 (m, 0.67H).

MS (CI, NH<sub>3</sub>) 536 (MH<sup>+</sup>)

**Example 22**

30 N-[2-(4-Acetyl-4-phenyl-piperidin-1-yl)-1-(R)-benzyloxymethyl-2-oxo-ethyl]-2-amino-isobutyramide hydrochloride

According to General Procedure A, 3.15 mg (13.1 mmol) of 4-acetyl-4-phenylpiperidine hydrochloride and 5.0 mg (13.1 mmol) of **3** were coupled to give {1-[2-(4-Acetyl-4-phenyl-piperidin-1-yl)-1-(R)-benzyloxymethyl-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester. The crude  
35 product was deprotected according to General Procedure C to give 5.09 mg of the title compound.

MS (CI, NH<sub>3</sub>) 508 (MH<sup>+</sup>)



-79-

**Example 23**

(R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-4-phenyl-piperidine-4-carboxylic acid ethyl ester hydrochloride

A. 4-Phenyl-piperidine-4-carboxylic acid ethyl ester hydrochloride

- 5 A mixture of 10.0 g (45 mmol) 4-cyano-4-phenylpiperidine hydrochloride, 10 g of sulfuric acid and 2.5 mL of water was heated at 150°C for 1 h. The mixture was allowed to cool to 110°C and 10 mL of ethanol was added. The ethanol was then distilled off. This ethanol addition/distillation was repeated four more times. Following the last addition, the mixture was heated  
10 to 125°C cooled to room temperature and poured onto ice. The mixture was basified with 10% NaOH and extracted twice with ether. The organic extracts were dried over MgSO<sub>4</sub> and concentrated to give 2.3 g of a yellow oil. A solution of 2.0 g of the free base in 100 mL of ether was treated with an ether solution saturated with HCl. The white precipitate was filtered and  
15 recrystallized from EtOAc/hexanes to give 1.73 g of **23A**

B. (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-4-phenyl-piperidine-4-carboxylic acid ethyl ester hydrochloride

- According to the method outlined in General Procedure A, 3.55 mg (13.1 mmol) of **23A** and 5.0 mg (13.1 mmol) of **3** were coupled to give 1-[3-benzyloxy-2-(2-tert-butoxycarbonylamino-2-methyl-propionylamino)-propionyl]-4-phenyl-piperidine-4-carboxylic acid ethyl ester. The crude product  
20 was then deprotected according to General Procedure C to give 6.97 mg of the title compound as a solid.

MS (Cl, NH<sub>3</sub>) 496 (MH<sup>+</sup>)

25

**Example 24**

2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-hydroxy-4-(4-trifluoromethyl-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

A. 4-Oxo-piperidine-1-carboxylic acid benzyl ester

- To a mixture of 50.0 g (0.325mol) of 4-piperidone hydrate hydrochloride  
30 and 181.9 g (1.82mol) of potassium bicarbonate in 750 mL of EtOAc and 75 mL of water, was added 49 mL (0.343mol) of benzyl chloroformate over 10 min. with stirring. The mixture was stirred at 24°C for 2.5 h and then diluted with water. The layers were separated, and the aqueous portion extracted with 500 mL of ethyl acetate. The combined organic extracts were washed once each  
35 with water and brine, dried over MgSO<sub>4</sub> and concentrated to give 81.41 g of **24A** as a yellow oil. The crude product was used without further purification.

B. 4-Hydroxy-4-trifluoromethyl-phenyl-piperidine-1-carboxylic acid benzyl ester

-80-

To 1.25 g (51.4 mmol) of magnesium in 10 mL of ether was added a solution of 4-bromobenzotrifluoride in 10 mL of ether. The reaction initiated soon after the addition was complete. The mixture was stirred for 1.5 h at 24°C and then cooled to 0°C. A solution of 10.0 g (42.87 mmol) of **24A** in 50 mL of ether was added and the mixture allowed to warm to room temperature and was stirred for 17 h. The reaction mixture was quenched by adding 150 mL of saturated aqueous ammonium chloride. The organic layer was separated and the aqueous phase was extracted with ether. The combined organic extracts were washed once each with water and brine, dried over MgSO<sub>4</sub> and concentrated. The crude material was purified by silica gel chromatography (3:7 v/v EtO Ac:hexanes) to give 12.48 g of **24B** as an orange oil which solidified on standing.

C. 4-(4-Trifluoromethyl-phenyl)-piperidin-4-ol

To 12.3 g (32.4 mmol) of **24B** in 150 mL of ethanol was added 1.4 g of 10% palladium on carbon. The mixture was hydrogenated on a Parr shaker at 48 psi for 2.5 h. The solution was filtered through celite and concentrated. The resulting solid was triturated with ether/hexane to give 4.98 g of **24C** as a white solid.

D. 2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-hydroxy-4-(4-trifluoromethyl-phenyl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

According to General Procedure A, 3.22 mg (13.1 mmol) of **24C**, and 5 mg (13.1 mmol) of **3** were coupled to give (1-{1-(R)-Benzyloxymethyl-2-[4-hydroxy-4-(4-trifluoromethyl-phenyl)-piperidin-1-yl]-2-oxo-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester. The crude product was deprotected according to General Procedure C to give 7.35 mg of the title compound.

MS (Cl, NH<sub>3</sub>) 466 (MH<sup>+</sup>)

**Example 25**

(R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-piperidine-4-carboxylic acid cyclohexylamide hydrochloride

A. Piperidine-1,4-dicarboxylic acid monobenzyl ester

To a slurry of 66.4 g (0.50 mol) of isonipecotic acid, and 145 mL (0.50 mol) of triethylamine in 600 mL of methylene chloride was added a solution of 72 mL (0.50 mol) of benzyl chloroformate in 200 mL of toluene at 10°C. The slurry was allowed to warm to room temperature and after stirring for 17 h was treated with 1 L of a 10% aqueous sodium carbonate solution. The layers were separated, and the aqueous portion was extracted once with ether and then acidified to pH 3. The aqueous portion was then extracted with methylene

-81-

chloride. The organic extract was washed with brine, dried over MgSO<sub>4</sub> and concentrated to give 35.2 g of the title compound as an oil which was used without further purification.

B. 4-Cyclohexylcarbamoyl-piperidine-1-carboxylic acid benzyl ester

5 To 3.82 g (14.3 mmol) of 25A was added 1.2 mL (15.7 mmol) of thionyl chloride in 50 mL of methylene chloride which contained 1 drop of DMF. The slurry was refluxed on a steam bath for 10 min. to give a clear solution. The mixture was concentrated, and the resulting oil was dissolved in 30 mL of THF and was added to a solution of 3.54 g (35.8 mmol) of cyclohexylamine in 20 mL  
10 of THF at 0°C. The mixture was stirred at 24°C for 0.5 h and then filtered. The filtrate was concentrated and then dissolved in methylene chloride. The organic solution was washed once each with water, 1N HCl, saturated aqueous sodium bicarbonate, and brine, dried over MgSO<sub>4</sub> and concentrated to give 3.46 g of 25B as a solid.

15 C. Piperidine-4-carboxylic acid cyclohexylamide

To 3.46 g (10 mmol) of 25B was added 30 mL of methanol, and the mixture refluxed in 20% NaOH for 8 h. The methanol was removed, and the aqueous phase extracted with methylene chloride. The aqueous solution was acidified to pH 5 with acetic acid and extracted with methylene chloride. The  
20 combined organic extracts were dried over MgSO<sub>4</sub> and concentrated to give 1.81 g of the title compound as an off-white solid.

D. (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-piperidine-4-carboxylic acid cyclohexylamide hydrochloride

According to General Procedure A, 2.76 mg (13.1 mmol) of 25C, and 5  
25 mg (13.1 mmol) of 3 were coupled to give {1-[1-(R)-Benzyloxymethyl-2-(4-cyclohexylcarbamoyl-piperidin-1-yl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester. The crude product was deprotected according to General Procedure C to give 6.60 mg of the title compound.

MS (Cl, NH<sub>3</sub>) 473 (MH<sup>+</sup>)

30 **Example 26**

(R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride

A. (R)-2-[2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid

35 To 756 mg (1.35 mmol) of 5E in 10.8 mL of THF was added a solution of 97 mg (4.04 mmol) of lithium hydroxide in 2.7 mL of water. The mixture was stirred for 17 h at room temperature. The mixture was concentrated and dissolved in 100 mL of water. The aqueous solution was extracted twice with

-82-

methylene chloride, acidified with acetic acid, and extracted twice with ethyl acetate. The ethyl acetate extracts were washed twice with brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to give 632 mg of **26A** as a white solid.

5     B. (R)-[1-[1-(6-Ethylcarbamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid

According to General Procedure A, 60 mg (0.11 mmol) of **26A** and 13.4 mg (0.16 mmol) of ethylamine hydrochloride were coupled and the product was purified by silica gel chromatography (100% ethyl acetate) to give 47 mg of **26B** as a foam.

10    C. (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride

According to General Procedure C, 47 mg (0.081 mmol) of **26B** was deprotected to give 39 mg of the title compound as a solid.

15    <sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (mixture of rotamers) (partial) δ 8.32 (d, 1H), 7.53-7.64 (m, 1.5H), 7.40-7.50 (m, 1.5H), 7.30 (d, 0.5H), 6.91-7.18 (m, 4H), 6.61 (d, 0.5H), 5.14-5.32 (m, 1H), 4.63-4.74 (d, 0.5H), 4.39-4.54 (m, 1H), 4.07-4.18 (d, 0.5H), 3.80-3.91 (m, 0.5H), 3.54-3.16 (m, 0.5H), 1.57 (s, 6H), 1.06-1.27 (m, 3H).

20    MS (CI, NH<sub>3</sub>) 476 (MH<sup>+</sup>)

### Example 27

(R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid phenylamide hydrochloride

25    A. (R)-[1-[2-(1H-Indol-3-yl)-1-(6-phenylcarbamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

To 60 mg (0.11 mmol) of **26A** in 0.6 mL of methylene chloride was added 11.4 mg (0.112 mmol) of triethylamine. The mixture was cooled to 0°C and 13.5 mg (0.112 mmol) of pivaloyl chloride was added in one portion. The mixture was stirred for 1.5 h at 0°C and then 10 mg (0.107 mmol) of aniline was added and the mixture was allowed to warm to room temperature and stirred for 17 h. The mixture was diluted with ethyl acetate and washed twice with 1N NaOH and brine. The solution was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (2:1 v/v ethyl acetate:hexanes) to give 39 mg of **27A** as a foam.

35    B. (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid phenylamide hydrochloride

-83-

According to the method outlined in General Procedure C, 37 mg (0.06 mmol) of **27A** was deprotected to give 32 mg of the title compound as a solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (mixture of rotamers) (partial) δ 7.53-7.79 (m, 6H), 7.37-7.42 (m, 3H), 6.95-7.22 (m, 5.5H), 6.67 (d, 0.5H), 5.29-5.32 (m, 1H), 4.63-4.79 (m, 0.5H), 4.43-4.59 (m, 1H), 4.09-4.19 (m, 0.5H), 3.78-3.92 (m, 0.5H), 3.39-3.63 (m, 2.5H), 1.59 (s, 6H).  
MS (CI, NH<sub>3</sub>) 524 (MH<sup>+</sup>)

### Example 28

10 (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-tetrahydro-isoquinoline-6-carboxylic acid hydrochloride

A. (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid hydrochloride

According to General Procedure C, 40 mg (0.073 mmol) of **26A** was deprotected to give 25 mg of the title compound as a solid.

15 <sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (mixture of rotamers) (partial) δ 7.78 (d, 0.5H), 7.52-7.68 (m, 2.5H), 7.39 (d, 0.5H), 7.23 (d, 0.5H), 6.95-7.19 (m, 3.5H), 6.58 (d, 0.5H), 5.17-5.31 (m, 1H), 4.63-4.74 (d, 0.5H), 4.42-4.53 (q, 1H), 4.09-4.20 (d, 1H), 1.58 (s, 6H).  
MS (CI, NH<sub>3</sub>) 449 (MH<sup>+</sup>)

20

### Example 29

(R)-2-Amino-N-(2-(1H-indol-3-yl)-1-[6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl)-isobutyramide hydrochloride

A. (R)-[1-(2-(1H-Indol-3-yl)-1-[6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-H-isoquinoline-2-carbonyl]-ethylcarbamoyl)-1-methyl-ethyl]-carbamic acid tert-butyl ester

25 According to General Procedure A, 72 mg (0.26 mmol) of **7H** and 100 mg (0.26 mmol) of **4C** were coupled and the product was purified by silica gel chromatography using 100% chloroform, followed by 2% methanol in chloroform to give 130 mg of **29A** as a foam.

30 B. (R)-2-Amino-N-(2-(1H-indol-3-yl)-1-[6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl)-isobutyramide hydrochloride

According to General Procedure C, 130 mg (0.20 mmol) of **29A** was deprotected to give 120 mg of the title compound as a tan solid.

35 <sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (1:1 mixture of rotamers) δ 7.57-7.72 (m, 2H), 6.99-7.37 (m, 10.5H), 6.68 (d, 0.5H), 5.18-5.32 (m, 1H), 4.37-4.75 (m, 2H), 3.58-3.81 (m, 0.5H), 3.53-3.68 (m, 0.5H), 3.17-3.32 (m, 2H), 2.62-2.80 (m, 4H), 2.41-2.60 (m, 0.5H), 2.14-2.39 (m, 0.5H), 1.58 (s, 6H).  
MS (CI, NH<sub>3</sub>) 552 (MH<sup>+</sup>)

-84-

**Example 30**

2-Amino-N-(1-(R)-benzyloxymethyl-2-oxo-2-[3-(toluene-4-sulfonylamino)-pyrrolidin-1-yl]-ethyl)-isobutyramide trifluoroacetate

A. 3-tert-Butoxycarbonylamino-pyrrolidine-1-carboxylic acid benzyl

5 ester

To a mixture of 19.0 g (102 mmol) of 3-(t-butoxycarbonylamino) pyrrolidine and 14.28 g (117 mmol) of 4-dimethylaminopyridine in 115 mL of methylene chloride at 0°C was added 20.0 g (117 mmol) of benzyl chloroformate dropwise. The mixture was allowed to warm to 24°C and was  
10 stirred overnight. The mixture was diluted with chloroform and washed twice with 10% HCl, twice with saturated aqueous sodium bicarbonate and once with brine. The mixture was dried over MgSO<sub>4</sub> and concentrated to give **30A** in quantitative yield. The crude material was used without further purification.

B. 3-Amino-pyrrolidine-1-carboxylic acid benzyl ester trifluoroacetate

15 To 9.4 g (29.3 mmol) of **30A** at 0°C was added 100 mL of trifluoroacetic acid and the mixture stirred at 24°C for 1 h. The mixture was concentrated, and coevaporated three times from ethyl acetate/heptane to give 11.0 g of **30B** as an orange oil.

C. 3-(Toluene-4-sulfonylamino)-pyrrolidine-1-carboxylic acid benzyl

20 ester

To a mixture of 2.0 g (6.0 mmol) of **30B**, 0.84 g (6.9 mmol) of 4-dimethylaminopyridine, and 0.61 g (6.0 mmol) of triethylamine in 10 mL of methylene chloride was added 1.32 g (6.9 mmol) of toluenesulfonyl chloride and the mixture was stirred overnight. The mixture was diluted with chloroform  
25 and washed twice each with 10% HCl, saturated aqueous bicarbonate and brine, dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography using a gradient elution of 35% ethyl acetate in hexane to 100% ethyl acetate to give **30C** as a white solid (700 mg).

D. 4-Methyl-N-pyrrolidin-3-yl-benzenesulfonamide

30 To 0.70 g (1.9 mmol) of **30C** in 20 mL of ethanol was added 0.40 g of 10% palladium on carbon and the mixture was hydrogenated on a Parr shaker for 17 h at 45 psi. The mixture was diluted with 110 mL of a (9:1:1) mixture of ethanol:water:ammonium hydroxide and was stirred for 20 min. The solution was filtered through celite and concentrated to give 300 mg of **30D** as a yellow  
35 solid.

E. (1-(1-(R)-Benzyloxymethyl-2-oxo-2-[3-(toluene-4-sulfonylamino)-pyrrolidin-1-yl]-ethylcarbamoyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

-85-

According to General Procedure A, 94 mg (0.39 mmol) of **30D** and 100 mg (0.26 mmol) of **3** were coupled and the residue was purified by silica gel chromatography using a gradient elution of 50% ethyl acetate in hexane, to 100% ethyl acetate to give 100 mg of **30E**.

5        F. 2-Amino-N-[1-(R)-benzyloxymethyl-2-oxo-2-[3-(toluene-4-sulfonylamino)-pyrrolidin-1-yl]-ethyl]-isobutyramide trifluoroacetate

To 100 mg (0.16 mmol) of **30E** was added 2 mL of trifluoroacetic acid at 0°C and the mixture was stirred for 1.5 h at 24°C. The mixture was concentrated, and coevaporated once each with ethyl acetate and hexane.

10      The product was precipitated from methylene chloride/hexane and was collected by filtration and dried under vacuum to give 95 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ (rotamers) 7.75-7.84 (m, 2H), 7.29-7.50 (m, 7H), 4.76-4.93 (m, 1H), 4.57 (d, 2H), 3.69-3.80 (m, 5H), 3.39-3.65 (m, 2.5H),  
15      3.22-3.31 (m, 0.5H), 2.45 (m, 3H), 1.62 (m, 6H).  
MS (CI, NH<sub>3</sub>) 503 (MH<sup>+</sup>)

**Example 31**

(R)-2-Amino-N-[1-(3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-isobutyramide hydrochloride

20      A. (R)-[1-[1-(3,4-Dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]carbonyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

According to General Procedure A, 60 mg (0.154 mmol) of **4C** and 26 mg (0.154 mmol) of 1,2,3,4-tetrahydroisoquinoline were coupled and the product was purified by silica gel chromatography (40:60 v/v ethyl  
25      acetate:hexanes) to give 71 mg of **31A**.

B. (R)-2-Amino-N-[1-(3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-isobutyramide hydrochloride

According to the method outlined in General Procedure C, 57 mg (0.106 mmol) of **31B** was deprotected to give 49 mg of the title compound.

30      <sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (mixture of rotamers) δ 8.32 (d, 1H), 7.56 (d, 1H), 7.32 (d, 0.5H), 6.96-7.22 (m, 7H), 6.61 (d, 0.5H), 5.17-5.32 (m, 1H), 4.61-4.73 (d, 0.5H), 4.45-4.58 (d, 0.5H), 4.29-4.41 (d, 0.5H), 3.96-4.05 (d, 0.5H), 3.37-3.72 (m, 2H), 3.13-3.26 (m, 2H), 2.54-2.74 (m, 1H), 2.39-2.55 (m, 0.5H), 2.11-2.25 (m, 0.5H), 1.49-1.62 (m, 6H).  
35      MS (CI, NH<sub>3</sub>) 405 (MH<sup>+</sup>)

-86-

**Example 32**

(R)-N-{1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-pyrrolidin-3-yl}-benzamide trifluoroacetate

A. 3-Benzoylamino-pyrrolidine-1-carboxylic acid benzyl ester

- 5 To a mixture of 2.00 g (6.0 mmol) of **30B**, 0.84 g (6.9 mmol) of 4-dimethylaminopyridine and 0.610 g (6.9 mmol) of triethylamine in 10 mL of methylene chloride was added 0.97 g (6.9 mmol) of benzoyl chloride and the mixture stirred for 72 h. The reaction mixture was diluted with chloroform and washed twice with 10% HCl, twice with saturated sodium bicarbonate solution  
10 and once with brine. The mixture was dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (75:25 v/v ethyl acetate/hexanes) to give 690 mg of **32A**.

B. N-Pyrrolidin-3-yl-benzamide

- To 0.69 g (2.1 mmol) of **32A** in 20 mL of ethanol was added 0.30 g of  
15 10% palladium on carbon and the mixture was hydrogenated on the Parr shaker for 17 h at 45 psi. The mixture was diluted with 110 mL of a (9:1:1) mixture of ethanol:water:ammonium hydroxide and stirred for 15 min. The solution was filtered through celite and concentrated to give 265 mg of **32B**.

- C. {1-[2-(3-Benzoylamino-pyrrolidin-1-yl)-1-(R)-benzyloxymethyl-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester  
20

According to General Procedure A, 100 mg (0.52 mmol) of **32B** and 75 mg (0.20 mmol) of **3** were coupled and the residue was purified by silica gel chromatography using a gradient elution of 75% ethyl acetate in hexane, to 100% ethyl acetate to give 50 mg of **32C**.

- 25 D. (R)-N-{1-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-pyrrolidin-3-yl}-benzamide trifluoroacetate

- To 50 mg (0.10 mmol) of **32C** at 0°C was added 2 mL of trifluoroacetic acid. The reaction mixture was stirred for 4 h at 24°C, and the residue was diluted with ethyl acetate and coevaporated with heptane and ethyl  
30 acetate/hexane. The residue was diluted with chloroform by addition of hexane to give 22 mg of the title compound as a solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) δ 7.86-7.99 (m, 2H), 7.46-7.65 (m, 3H), 7.23-7.45 (m, 5H), 4.52-4.74 (m, 3H), 3.39-4.26 (m, 7H), 1.59 (bs, 6H).  
LSIMS-MS 453 (MH<sup>+</sup>)



-87-

**Example 33**

2-Amino-N-{1-(5-fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

5     A. 2-tert-Butoxycarbonylamino-2-methyl-propionic acid 2,5-dioxo-pyrrolidin-1-yl ester

To a slurry of 5.0 g (24.6 mmol) of N-t-BOC- $\alpha$ -methylalanine in 13.5 mL of methylene chloride was added 3.40 g (29.6 mmol) of N-hydroxysuccinimide and 5.65 g (29.6 mmol) of EDC. The slurry was stirred for 17 h at room temperature. The mixture was diluted with ethyl acetate and washed twice  
10 each with water, saturated aqueous sodium bicarbonate and brine. Dried over MgSO<sub>4</sub> and concentrated. The residue was purified by silica gel chromatography (1:1 v/v ethyl acetate:hexanes) to give 5.2 g of **33A** as a white solid.

15     B. -2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(5-fluoro-1H-indol-3-yl)-propionic acid

A mixture of 100 mg (0.33 mmol) of **33A**, 62 mg (0.278 mmol) of 5-fluoro-DL-tryptophan and 79 mg (0.611 mmol) of diisopropylethylamine in 1.0 mL of DMF was stirred for 17 h at room temperature. The reaction mixture was poured into water and extracted twice with ethyl acetate. An equal volume of  
20 hexane was added to the ethyl acetate and the mixture was washed three times with water and once with brine. The organic phase was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (9:1 v/v CHCl<sub>3</sub>:MeOH) to give 52 mg of **33B**.

25     C. -(1-[1-(5-Fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl]carbonyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 44 mg (0.11 mmol) of **33B** and 23.5 mg (0.11 mmol) of 4-(2-keto-1-benzimidazol-1-yl)piperidine were coupled and the product was purified by silica gel chromatography (100% ethyl acetate) to  
30 give 39 mg of **33C** as an amorphous solid.

D. -2-Amino-N-{1-(5-fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

According to General Procedure C, 39 mg (0.064 mmol) of **33C** was deprotected to give 32 mg of the title compound as a white solid.

35     <sup>1</sup>H NMR (CD<sub>3</sub>OD 250MHz) (partial) (1:1 mixture of diastereomers)  $\delta$  8.36 (d, 1H), 7.03-7.39 (m, 7.5H), 6.89 (t, 1H), 6.76 (d, 0.5H), 5.14-5.35 (m, 1H), 4.59-4.71 (m, 1H), 4.23-4.46 (m, 1H), 3.94-4.11 (m, 1H), 2.45-2.72 (m, 2H), 1.50-1.64 (m, 6H).

-88-

MS (Cl, NH<sub>3</sub>) 507 (MH<sup>+</sup>)**Example 34**

(R)-N-[1-[4-(Acetyl-phenyl-amino)-piperidine-1-carbonyl]-2-(1H-indol-3-yl)-ethyl]-2-amino-isobutyramide

5 A. (R)-[1-[1-[4-(Acetyl-phenyl-amino)-piperidine-1-carbonyl]-2-(1H-indol-3-yl)-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

According to General Procedure A, 45 mg (0.116 mmol) of **4C** and 27 mg (0.124 mmol) of N-phenyl-N-piperidin-4-yl-acetamide were coupled and the product was purified by silica gel chromatography using a gradient of 1% to 10 5% methanol in methylene chloride to give 63 mg of **34A**.

B. (R)-N-[1-[4-(Acetyl-phenyl-amino)-piperidine-1-carbonyl]-2-(1H-indol-3-yl)-ethyl]-2-amino-isobutyramide

According to General Procedure B, 63 mg (0.107 mmol) of **34A** was deprotected to give 33 mg of the title compound as a white solid.

15 <sup>1</sup>H NMR (CD<sub>3</sub>OD 300MHz) (1:1 mixture of rotamers) δ 7.31-7.62 (m, 4H), 6.89-7.23 (m, 5H), 5.11-5.20 (t, 1H), 4.99-5.19 (t, 1H), 4.40-4.63 (m, 2H), 3.68-3.83 (m, 2H), 3.02-3.17 (t, 2H), 2.90-3.01 (t, 0.5H), 2.50-2.60 (t, 0.5H), 2.32-2.41 (m, 1H), 1.64-1.82 (m, 1H), 1.69 (s, 3H), 1.32 (s, 6H).  
MS (Cl, NH<sub>3</sub>) 490 (MH<sup>+</sup>)

20

**Example 35**

(R)-2-Amino-N-[1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethyl]-isobutyramide hydrochloride

A. [1-Naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethyl]-carbamic acid tert-butyl ester

25 According to General Procedure A, 605 mg (1.92 mmol) of BOC-D-3-2-(Naphthyl)alanine and 417 mg (1.92 mmol) of 4-(2-keto-1-benzimidazoliny) piperidine were coupled to give 890 mg of **35A** as a solid which was used without further purification.

30 B. 1-[1-(2-Amino-3-naphthalen-2-yl-propionyl)-piperidin-4-yl]-1,3-dihydro-benzimidazol-2-one

According to General Procedure B, 830 mg (1.61 mmol) of **35A** was deprotected to give 582 mg of **35B** as a tan solid.

C. (R)-[1-Methyl-1-[1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl]-ethyl]-carbamic acid tert-butyl ester

35

According to General Procedure A, 570 mg (1.38 mmol) of **35B** and 266 mg (1.31 mmol) of N-t-BOC-α-methylalanine were coupled to give 63 mg of

-89-

**35C** after silica gel chromatography using a gradient of 1% to 3% methanol in methylene chloride.

D. (R)-2-Amino-N-[1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl]-isobutyramide hydrochloride

5 According to General Procedure C, 620 mg (1.03 mmol) of **35C** was deprotected to give 404 mg of the title compound as a light yellow solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD 300MHz) (mixture of rotamers) (partial) δ 7.73-7.87 (m, 4H), 7.40-7.50 (m, 3H), 7.19-7.28 (m, 0.5H), 6.91-7.05 (m, 3H), 6.74-6.79 (d, 0.5H), 5.26-5.48 (m, 1H), 4.67-4.71 (m, 1H), 4.33-4.50 (m, 1H), 4.10-4.32 (m, 1H), 3.12-3.39 (m, 2H), 1.55-1.61 (d, 3H), 1.38 (s, 3H).

10 MS (CI, NH<sub>3</sub>) 500 (MH<sup>+</sup>)

### Example 36

(R)-2-Amino-N-[2-(1H-indol-3-yl)-1-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl]-isobutyramide

15 A. (R)-[1-[2-(1H-Indol-3-yl)-1-(7-nitro-3,4-dihydro-1H-isoquinoline-2-carbonyl)-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

According to General Procedure A, 750 mg (4.21 mmol) of **6A** and 1.64 g (4.21 mmol) of **4C** were coupled and the product was purified by silica gel chromatography (30:70 v/v EtOAc:hexanes) to give 1.50 g of **36A** as a foam.

20 B. (R)-[1-[1-(7-Amino-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

To 800 mg (1.46 mmol) of **36A** in 10 mL of methanol was added 300 mg of 10% palladium hydroxide on carbon and the mixture hydrogenated at 50 psi for 17 h. The mixture was filtered through celite and concentrated to give 700 mg of **36B** as a brown powder.

C. (R)-[1-[2-(1H-Indol-3-yl)-1-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

To 71 mg (0.14 mmol) of **36B** in 10 mL of methylene chloride at 0°C was added 17 mg (0.165 mmol) of triethylamine, followed by 29 mg (0.15 mmol) of toluenesulfonyl chloride. The mixture was allowed to warm to room temperature over 3 h. The reaction mixture was diluted with ethyl acetate and washed twice with 1N NaOH and once with brine. The solution was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated, and the residue was purified by silica gel chromatography (1:2 v/v EtOAc:hexanes) to give 90 mg of **36C** as a foam.

D. (R)-2-Amino-N-[2-(1H-indol-3-yl)-1-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl]-isobutyramide

-90-

According to General Procedure B, 50 mg (0.74 mmol) of **36C** was deprotected to give 20 mg of the title compound as a white powder.

MS (Cl, NH<sub>3</sub>) 574 (MH<sup>+</sup>)

### Example 37

5     (R)-N-{2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl}-benzamide

A. (R)-{1-[1-(7-Benzoylamino-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbonyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

To 86 mg (0.17 mmol) of **36B** in 10 mL of methylene chloride at 0°C was  
10   added 20 mg (0.20 mmol) of triethylamine and 26 mg (0.18 mmol) of benzoyl chloride. The mixture was warmed to room temperature over 2 h and diluted with ethyl acetate and washed twice with 1N NaOH and once with brine. The solution was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated, and the residue was purified by silica gel chromatography (1:1 v/v EtOAc/hexanes) to give 95 mg of  
15   **37A** as a foam.

B. (R)-N-{2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl}-benzamide

According to General Procedure B, 95 mg (0.15 mmol) of **37A** was deprotected to give 65 mg of the title compound.

20     MS (Cl, NH<sub>3</sub>) 524 (MH<sup>+</sup>)

### Example 38

2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(2-cyclopropyl-5,7-dimethyl-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide

A. 4-(2-Cyclopropyl-5,7-dimethyl-imidazo[4,5-b]pyridin-3-yl)-piperidine-1-carboxylic acid tert-butyl ester  
25

2-Cyclopropyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine was prepared according to literature methods (for preparation see Carpino et al., Bioorg. & Med. Chem. Lett. 4, pp. 93-98 1994 (ref.13)). To 625 mg (2.4 mmol) of triphenylphosphine in 10 mL of toluene was added 460 mg (2.0 mmol) of di-  
30   tert-butylazodicarboxylate, and the mixture was stirred for 15 min. A mixture of 430 mg (2.3 mmol) of 2-cyclopropyl-5,7-dimethyl-3H-imidazo[4,5-b]pyridine and 325 mg (1.62 mmol) of 4-hydroxy-piperidine-1-carboxylic acid tert-butyl ester was added to the stirring solution and the mixture was stirred for 17 h at room temperature. The mixture was concentrated and the product was purified  
35   by silica gel chromatography (50:50 v/v hexanes: ethyl acetate) to give 150 mg of **38A**.

B. 2-Cyclopropyl-5,7-dimethyl-3-piperidin-4-yl-3H-imidazo[4,5-b]pyridine

-91-

According to General Procedure B, 150 mg (0.39 mmol) of **38A** was deprotected to give 60 mg of **38B**.

- C. (1-{1-(R) Benzyloxymethyl-2-[4-(2-cyclopropyl-5,7-dimethyl-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-2-oxo-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 60 mg (0.22 mmol) of **38B** was coupled to 84 mg (0.22 mmol) of **3** and the product was purified by silica gel chromatography (9:1 v/v ethyl acetate:hexanes) to give 37 mg of **38C**.

- D. 2-Amino-N-[1-(R)-benzyloxymethyl-2-[4-(2-cyclopropyl-5,7-dimethyl-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-2-oxo-ethyl]-isobutyramide

According to General Procedure B, 37 mg of **38C** was deprotected to give 25 mg of the title compound.

- <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300MHz) δ 7.29-7.37 (m, 3H); 7.16-7.24 (m, 2H), 6.87-6.89 (s, 1H), 5.13-5.21 (m, 1H), 4.92-5.03 (m, 1H), 4.68-4.82 (m, 1H) 4.50-4.67 (m, 3H), 4.71-4.89 (m, 1H), 3.66-3.77 (m, 2H), 2.73-2.92 (m, 3H), 2.51 (s, 5H), 2.43 (s, 1H), 2.10-2.30 (m, 1H), 1.86-2.94 (m, 2H), 1.32 (d, 6H), 1.02-1.26 (m, 4H).  
MS (CI, NH<sub>3</sub>) 533 (MH<sup>+</sup>)

### Example 39

- (R)-2-Amino-N-[2-(4-cyano-4-phenyl-piperidin-1-yl)-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide

A. (R)-{1-[2-(4-Cyano-4-phenyl-piperidin-1-yl)-1-(1H-indol-3-ylmethyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

- According to General Procedure A, 53 mg (0.24 mmol) of 4-phenyl-piperidine-4-cyano hydrochloride and 92 mg (0.24 mmol) of **4C** were coupled and the product was purified by silica gel chromatography (1:1 v/v ethyl acetate:hexanes) to give 124 mg of **39A**.

B. (R)-2-Amino-N-[2-(4-cyano-4-phenyl-piperidin-1-yl)-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide

- According to General Procedure B, 105 mg (0.19 mmol) of **39C** was deprotected to give 82 mg of the title compound as a foam.

- <sup>1</sup>H NMR (CDCl<sub>3</sub>, 250MHz) (60:40 mixture of rotamers) δ 8.13-8.34 (m, 2H), 7.75 (d, 0.6H), 7.68 (d, 0.4H), 7.28-7.41 (m, 4H), 7.08-7.27 (m, 3H), 6.94-7.01 (m, 1H), 5.13-5.33 (m, 1H), 4.62-4.78 (m, 1H), 3.73-3.90 (m, 1H), 3.20-3.31 (m, 2H), 2.64-2.93 (m, 2H), 1.66-2.43 (bm, 6H), 1.43 (s, 6H).  
MS (CI, NH<sub>3</sub>) 458 (MH<sup>+</sup>)

-92-

**Example 40**

(R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

- 5     A. (R)-{1-[1-(1H-indol-3-ylmethyl)-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 27 mg (0.087 mmol) of morpholin-4-yl-(4-phenyl-piperidin-4-yl)-methanone hydrochloride and 32 mg (0.081 mmol) of **4C** were coupled and the product was purified by silica gel chromatography using a gradient of 50% ethyl acetate in hexane to 100% ethyl acetate to give 24 mg of **40A** as a foam.

B. (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

According to the method outlined in General Procedure C, 36 mg of **40A** was deprotected to give 17 mg of the title compound as a yellow solid.

- <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (rotamers) δ 7.52-7.67 (m, 1H), 6.99-7.40 (m, 8H), 6.87 (d, 1H), 5.07-5.21 (m, 1H), 4.03-4.30 (m, 1H), 3.52-3.68 (m, 1H), 2.74-3.39 (m, 13H), 1.94-2.18 (m, 1H), 1.74-1.90 (m, 1H), 1.58-1.66 (m, 1H), 1.30-1.47 (d, 6H), 1.25-1.30 (m, 2H).
- 20   MS (CI, NH<sub>3</sub>) 546 (MH<sup>+</sup>)

**Example 41**

1-Amino-cyclopentanecarboxylic acid {1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide trifluoroacetate

- 25     A. 1-tert-Butoxycarbonylamino-cyclopentanecarboxylic acid

To 1.66 g (12.88 mmol) of 1-amino-1-cyclopentane-carboxylic acid in 12.5 mL of 2N NaOH was added 283 mg (13.0 mmol) of di-tert-butyl-dicarbonate and the mixture was stirred for 17 h at room temperature. The mixture was acidified with 10% HCl and extracted three times with ethyl acetate. The combined organic extracts were washed with brine, dried over MgSO<sub>4</sub> and concentrated to give 1.71 g of **41A**.

30

-93-

B. 1-[1-(R)-Benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl]-cyclopentyl]-carbamic acid tert-butyl ester

According to General Procedure A, 20 mg (0.05 mmol) of **1B** and 11.6 mg (0.05 mmol) of **41A** were coupled, and 23 mg of **41B** was recovered. The crude material was used without further purification in the following step.

C. 1-Amino-cyclopentanecarboxylic acid {1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide trifluoroacetate

To 23 mg (0.04 mmol) of **41B** was added 2 mL of trifluoroacetic acid at 0°C. The ice bath was removed and the mixture was stirred for 2.5 h, and then was diluted with ethyl acetate and concentrated. The residue was coevaporated once from heptane, dissolved in methylene chloride and the product precipitated with hexane to give 10 mg of the title compound as a solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) δ 7.29-7.45 (m, 7H), 7.06-7.28 (m, 4H), 5.17-5.36 (m, 1H), 4.04-4.31 (m, 4H), 4.19-4.38 (m, 1H), 3.72-3.87 (m, 2H), 2.25-2.44 (m, 3H), 1.80-2.12 (m, 9H).

MS (CI, NH<sub>3</sub>) 506 (MH<sup>+</sup>)

### Example 42

20 2-Amino-N-[1-(R)-benzyloxymethyl-2-(3-diphenylacetyl-amino-pyrrolidin-1-yl)-2-oxo-ethyl]-isobutyramide trifluoroacetate

A. 3-Diphenylacetyl-amino-pyrrolidine-1-carboxylic acid benzyl ester

To a mixture of 2.40 g (7.19 mmol) of **30B**, 1.10 g (9.02 mmol) of 4-dimethylaminopyridine, and 0.73 g (7.19 mmol) of triethylamine in 10 mL of methylene chloride was added 1.91 g (8.26 mmol) of diphenylacetylchloride and the mixture stirred overnight at room temperature. The mixture was diluted with chloroform and washed twice each with 10% HCl, saturated aqueous sodium bicarbonate, and brine. The mixture was dried over MgSO<sub>4</sub> and concentrated to give 2.43 g of **42A** as a white solid.

30 B. 2,2-Diphenyl-N-pyrrolidin-3-yl-acetamide

To 2.40 g (5.8 mmol) of **42A** in 40 mL of ethanol was added 1.20 g of 10% palladium on carbon and the mixture was hydrogenated at 45 psi for 17 h. The reaction mixture was filtered through celite and was concentrated to give 1.67 g of **42B** as a yellow oil.

35 C. {1-[1-(R)-Benzyloxymethyl-2-(3-diphenylacetyl-amino-pyrrolidin-1-yl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 150 mg (0.53 mmol) of **42B** and 200 mg (0.53 mmol) of **3** were coupled and the product was purified by silica gel

-94-

chromatography using a gradient of 50% ethyl acetate in hexane to 100% ethyl acetate to give 100 mg of **42C** as a white solid.

D. 2-Amino-N-[1-(R)-benzyloxymethyl-2-(3-diphenylacetyl-amino-pyrrolidin-1-yl)-2-oxo-ethyl]-isobutyramide trifluoroacetate

- 5 To 100 mg (0.16 mmol) of **42C** was added 3 mL of trifluoroacetic acid at 0°C. The ice bath was removed and the mixture stirred for 3 h, diluted with ethyl acetate and concentrated. The residue was coevaporated once with heptane. Hexane and chloroform were added to the residue and the precipitated product was collected by filtration. Recovered 55 mg of the title  
10 compound as a solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (partial) δ 7.22-7.40 (m, 15H), 4.95-5.00 (m, 1H), 4.38-4.57 (m, 3H), 1.60 (s, 6H).

LSIMS-MS 543 (MH<sup>+</sup>)

**Example 43**

- 15 (R)-N-[2-[2-(2-Amino-2-methyl-propionyl-amino)-3-benzyloxy-propionyl]-2,3-dihydro-1H-isoindol-5-yl]-benzamide trifluoroacetate

A. 1,2-Bis-bromomethyl-4-nitro-benzene

- To 8.00 g (53.0 mmol) of 4-Nitro-o-xylene in 80 mL of carbon tetrachloride was added 18.8 g (106 mmol) of N-Bromosuccinimide followed by  
20 100 mg (0.60 mmol) of 2,2'-azobis (isobutyronitrile) and the mixture was refluxed overnight. The precipitated solid was removed by filtration and washed with carbon tetrachloride and the filtrate concentrated to give 15.4 g of **43A** as a yellow oil. The crude product was used without further purification in the following step.

- 25 B. 2-Benzyl-5-nitro-2,3-dihydro-1H-isoindole

- To a mixture of 15.4 g (50.5 mmol) of **43A** and 35.4 g (430 mmol) of sodium carbonate in 160 mL of acetone and 35 mL of water was added 5.68 g (53 mmol) of benzylamine in 10 mL of acetone over 3 h. The mixture was then stirred for 17 h at room temperature. The precipitated solids were removed by  
30 filtration and the filtrate concentrated. The residue was dissolved in ethyl acetate and washed three times with 1N HCl. The combined HCl washes were neutralized with sodium carbonate solution and extracted three times with ethyl acetate. The combined organics were washed with brine, dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography  
35 (15:85 v/v ethyl acetate:hexanes) to give 2.55 g of **43B** as a yellow solid.

C. 5-Amino-2,3-dihydro-1H-isoindole

To 1.50 g (5.9 mmol) of **43B** in 50 mL of ethanol was added 1.25 g of 10% palladium on carbon and the mixture hydrogenated at 40°C and 5 psi for



-95-

17 h. The mixture was filtered through celite and concentrated to give 600 mg of **43C** as a pale purple solid.

D. {1-[2-(5-Amino-1,3-dihydro-isoindol-2-yl)-1-(R)-benzyloxymethyl-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

5 According to General Procedure A, 300 mg (2.2 mmol) of **43C** and 850 mg (2.2 mmol) of **3** were coupled to give 950 mg of **43D** as a yellow solid. The crude product was used without further purification.

E. {1-[2-(5-Benzoylamino-1,3-dihydro-isoindol-2-yl)-1-(R)-benzyloxymethyl-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-  
10 butyl ester

To a mixture of 300 mg (0.60 mmol) of **43D** and 85 mg (0.69 mmol) of 4-dimethylaminopyridine in 5 mL of methylene chloride was added 98 mg (0.69 mmol) of benzoyl chloride, and the mixture stirred for 17 h at room temperature. The reaction mixture was diluted with chloroform and washed twice each with  
15 10% HCl, saturated aqueous sodium bicarbonate and brine, dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (87:13 v/v ethyl acetate:hexanes) to give 120 mg of the title compound as a white solid.

F. (R)-N-{2-[2-(2-Amino-2-methyl-propionylamino)-3-benzyloxy-propionyl]-2,3-dihydro-1H-isoindol-5-yl}-benzamide trifluoroacetate  
20

To 120 mg (0.20 mmol) of **43D** was added 3 mL of trifluoroacetic acid at 0°C. The ice bath was removed and the mixture stirred for 3 h. The reaction mixture was diluted with ethyl acetate and concentrated. The product crystallized upon addition of hexane and chloroform. Filtration of the  
25 precipitate provided 50 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) δ 7.82-8.00 (d, 2H), 7.78-7.85 (m, 1H), 7.00-7.17 (m, 4H), 7.37-7.42 (m, 6H), 5.08-5.32 (m, 1H), 4.99-5.10 (m, 2H), 4.75-4.80 (d, 2H), 4.61 (s, 2H), 3.82-3.90 (m, 2H), 1.62 (s, 6H)  
MS (CI, NH<sub>3</sub>) 501 (MH<sup>+</sup>)

30 **Example 44**

(R)-2-Amino-N-{1-(4-benzyloxy-benzyl)-2-oxo-2-[4-(2-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide

A. (R)-3-(4-Benzyloxy-phenyl)-2-tert-butoxycarbonylamino-propionic acid methyl ester

35 To a mixture of 2.00 g (5.38 mmol) of N-t-BOC-O-benzyl-D-Tyrosine and 0.819 g (5.92 mmol) of potassium carbonate in 30 mL of DMF was added 0.802 g (5.65 mmol) of iodomethane, and the mixture stirred for 17 h at room temperature. The mixture was diluted with 150 mL water and extracted three

-96-

times with ethyl acetate. The organics were washed four times with water and once with brine. Drying over  $\text{MgSO}_4$  and concentration provided 2.27 g of **44A** as a yellow oil.

B. (R)-2-Amino-3-(4-benzyloxy-phenyl)-propionic acid methyl ester

5 Following the method outlined in General Procedure B, 2.27 g (5.4 mmol) of **44A** was deprotected to give 1.00 g of **44B** as a yellow oil.

C. (R)-3-(4-Benzyloxy-phenyl)-2-(2-tert-butoxycarbonylamino-2-methyl-propionylamino)-propionic acid methyl ester

10 According to General Procedure A, 1.00 g (3.5 mmol) of **44B** and 0.72 g (3.9 mmol) of N-t-BOC- $\alpha$ -methylalanine were coupled and the product was purified by silica gel chromatography (25:75 v/v ethyl acetate:hexanes) to give 1.00 g of **44C** as a white solid.

D. (R)-3-(4-Benzyloxy-phenyl)-2-(2-tert-butoxycarbonylamino-2-methyl-propionylamino)-propionic acid

15 Following the method outlined in General Procedure D, 1.00 g (2.2 mmol) of **44C** was hydrolyzed to give 1.00 g of **44D**.

E. (R)-(1-(1-(4-Benzyloxy-benzyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-ethyl)-carbamic acid tert-butyl ester

20 According to General Procedure A, 150 mg (0.34 mmol) of **44D** and 75 mg (0.34 mmol) of 4-(2-keto-1-benzimidazolyl)piperidine were coupled and the product was purified by silica gel chromatography using a gradient of (75:25 v/v ethyl acetate:hexanes) to (5:95 v/v methanol:ethyl acetate) to give 100 mg of **44E** as a white solid.

25 F. (R)-2-Amino-N-{1-(4-benzyloxy-benzyl)-2-oxo-2-[4-(2-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide

According to General Procedure B, 100 mg (0.15 mmol) of **44E** was deprotected to give 60 mg of the title compound as a white solid.

30  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 250MHz) (rotamers)  $\delta$  7.26-7.52 (m, 6H), 6.91-7.22 (m, 7H), 5.04-5.18 (m, 2H), 4.96-5.01 (m, 1H), 4.60-4.81 (m, 1H), 4.36-4.60 (m, 1H), 3.95-4.14 (m, 1H), 2.94-3.10 (m, 2H), 2.45-2.80 (m, 2H), 2.08-2.40 (m, 1H), 1.69-1.92 (m, 2H), 1.55-1.68 (m, 1H), 1.28-1.40 (m, 8H).

MS (CI,  $\text{NH}_3$ ) 556 ( $\text{MH}^+$ )

### Example 45

35 (R)-2-Amino-N-{3-(3,4-dihydro-1H-isoquinolin-2-yl)-3-oxo-1-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carbonyl]-propyl}-isobutyramide

A. (R)-3-tert-Butoxycarbonylamino-4-oxo-4-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-butyric acid benzyl ester

-97-

According to General Procedure A, 2.33 g (7.21 mmol) of N-a-t-BOC-D-aspartic-b-benzyl ester and 1.49 g (6.86 mmol) of 4-(2-keto-1-benzimidazoliny) piperidine were coupled and the product was purified by silica gel chromatography using an elution gradient of 2% to 4% methanol in methylene chloride to provide 2.56 g of **45A**.

B. (R)-3-Amino-4-oxo-4-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-butyric acid benzyl ester

Following the method outlined in General Procedure B, 2.49 g (4.78 mmol) of **45A** was deprotected to give 1.77 g of **45B**.

C. (R)-3-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-4-oxo-4-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-butyric acid benzyl ester

According to General Procedure A, 1.60 g (3.8 mmol) of **45B** and 0.77 g (3.8 mmol) of N-t-BOC- $\alpha$ -methylalanine were coupled and the product was purified by silica gel chromatography using an elution gradient of 2% to 4% methanol in methylene chloride to provide 2.46 g of **45C**.

D. (R)-3-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-4-oxo-4-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-butyric acid

To 2.44 g (4.0 mmol) of **45C** in 40 mL of ethanol was added 250 mg of 10% palladium on carbon and the mixture hydrogenated at 50 psi for 2 h. The mixture was filtered through celite and concentrated to give 2.00 g of **45D**.

E. (R)-(1-{3-(3,4-Dihydro-1H-isoquinolin-2-yl)-3-oxo-1-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carbonyl]-propylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 80 mg (0.19 mmol) of **45D** and 22 mg (0.20 mmol) of 1,2,3,4-tetrahydroisoquinoline were coupled and the product was purified by silica gel chromatography using an elution gradient of 2% to 4% methanol in methylene chloride to provide 52 mg of **45E**.

F. (R)-2-Amino-N-{3-(3,4-dihydro-1H-isoquinolin-2-yl)-3-oxo-1-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carbonyl]-propyl}-isobutyramide

Following the method outlined in General Procedure B, 45 mg (0.07 mmol) of **45E** was deprotected to give 22 mg of the title compound.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 300MHz) (1:1 mixture of rotamers) (partial)  $\delta$  7.54-7.62 (m, 0.5H), 7.20-7.32 (m, 0.5H), 7.10-7.21 (m, 4H), 7.01-7.19 (m, 3H), 5.33-5.49 (m, 1H), 3.67-3.92 (m, 2H), 2.58-3.01 (m, 5H), 1.74-1.89 (m, 2H), 1.39-1.42 (m, 8H).

MS (CI,  $\text{NH}_3$ ) 533 ( $\text{MH}^+$ )

-98-

**Example 46**

(R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-phenyl-4-(pyrrolidine-1-carbonyl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

5 A. (1-{1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-phenyl-4-(pyrrolidine-1-carbonyl)-piperidin-1-yl]-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 71 mg (0.24 mmol) of (4-Phenyl-piperidin-4-yl)-pyrrolidin-1-yl-methanone hydrochloride and 94 mg (0.24 mmol) of **4C** were coupled and the product was purified by silica gel chromatography  
10 using 100% ethyl acetate to elute to give 142 mg of **46A**.

B. (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-phenyl-4-(pyrrolidine-1-carbonyl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

According to General Procedure C, 133 mg (0.21 mmol) of **46A** was deprotected to give 119 mg of the title compound as a solid.

15 <sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) δ 7.52-7.63 (m, 1H), 7.00-7.37 (m, 8H), 6.80-6.91 (m, 1H), 5.02-5.26 (m, 1H), 3.99-4.27 (m, 1H), 3.04-3.72 (m, 6H), 2.61-3.02 (m, 3H), 1.44-2.28 (m, 13H).  
MS (CI, NH<sub>3</sub>) 530 (MH<sup>+</sup>)

**Example 47**

20 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(1-phenyl-1H-indol-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

A. [4-(1H-Indol-3-yl)-4H-pyridin-1-yl]-phenyl-methanone

To a solution of 23.4 g (0.20 mol) of indole in 250 mL of pyridine was added 23.2 mL (0.20 mol) of benzoyl chloride with mechanical stirring and the  
25 mixture was stirred for 48 h. After dilution with 100 mL of water and acidification with 6N HCl, the mixture was extracted twice with ether and the ether was washed once with 6N HCl. Drying over MgSO<sub>4</sub> and filtration through celite provided an oil which was triturated with methanol to give 18.4 g of **47A** as a pale yellow solid.

30 B. 3-Pyridin-4-yl-1H-indole

A mixture of 12.0 g (0.04mol) of **47A**, 0.6 g of 10% palladium on carbon, 8 mL of dibenzylamine and 70 mL of diphenyl ether were heated at 210°C for 7.5 h. Off-white needles precipitated from solution on cooling and these needles were filtered and washed with ether. The solids were suspended in  
35 50 mL of 1N HCl and washed once with ether. The aqueous solution was filtered through celite and concentrated. To the residue was added 10 mL of 5N NaOH, and the resulting solid was filtered and washed with water to provide 4.02 g of **47B**.

-99-

C. 1-Phenyl-3-pyridin-4-yl-1H-indole

To a mixture of 4.0 g (20.6 mmol) of **47B**, 5.91 g (20.6 mmol) of cuprous bromide, 2.84 g (20.6 mmol) of potassium carbonate in 40 mL of N-methylpyrrolidine was added 9.1 mL (86.5 mmol) of bromobenzene and the mixture heated at 190°C for 20 h. After cooling to room temperature the mixture was filtered through celite and the filter pad washed with ethyl acetate. The solution was refiltered and washed with 5N ammonium hydroxide solution until a blue color no longer appeared in the aqueous layer. The organic portion was washed twice with water, once with brine, dried over MgSO<sub>4</sub> and concentrated to give 3.3 g of **47C** as a light brown solid.

D. 3-(1-Benzyl-pyridin-4-yl)-1-phenyl-1H-indole bromide

A mixture of 2.7 g (10.0 mmol) of **47C** and 1.18 mL (10.0 mmol) of benzyl bromide in 50 mL of benzene was heated on a steam bath for 3 h. After cooling to room temperature, the precipitated yellow solid was removed by filtration and washed with benzene to give 3.03 g of **47D**.

E. 3-(1-Benzyl-1,2,3,6-tetrahydro-pyridin-4-yl)-1-phenyl-1H-indole

To 3.0 g (6.8 mmol) of **47D** in 100 mL of methanol was added 18.5 g (490 mmol) of sodium borohydride over 40 min. The mixture was stirred at room temperature for 1 h and then heated on a steam bath for 1.5 h. After cooling to room temperature, the mixture was diluted with water and ether, and potassium carbonate was added until the lower layer clouded. The mixture was filtered and the ether phase dried over MgSO<sub>4</sub> and potassium carbonate. Filtration and concentration provided 2.13 g of **47E** as a yellow oil.

F. 1-Phenyl-3-piperidin-4-yl-1H-indole hydrochloride

A mixture of 1.7 g (6.21 mmol) **47E** and 1.0 g of 10% palladium on carbon in 200 mL of ethanol was hydrogenated at 50°C for 5 h. 0.78 mL of concentrated HCl was added and hydrogenation continued at 50°C for another 17 h. The mixture was filtered through celite and concentrated to give 1.26 g of **47F** as a tan solid.

G. (R)-(1-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(1-phenyl-1H-indol-3-yl)-piperidin-1-yl]-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 70 mg (0.21 mmol) of **47F** and 83 mg of **4C** were coupled and the product was purified by silica gel chromatography (65:35 v/v ethyl acetate:hexane) to give 90 mg of **47G** as a white solid.

H. (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(1-phenyl-1H-indol-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

-100-

To a solution of 90 mg (0.14 mmol) of **47G** in 4 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 1 h. The mixture was concentrated and the residue crystallized from chloroform/hexane to give 72 mg of the title compound as a solid.

- 5        <sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) δ (partial) 7.34-7.74 (m, 8H), 7.01-7.29 (m, 6H), 6.78 (s, 1H), 5.22-5.39 (m, 1H), 4.45-4.69 (m, 1H), 3.93-4.06 (m, 1H), 1.68 (s, 6H).  
MS (CI, NH<sub>3</sub>) 548 (MH<sup>+</sup>)

#### Example 48

- 10    (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(1H-indol-3-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

A. 3-(1,2,3,6-Tetrahydro-pyridin-4-yl)-1H-indole

- 50 mL of methanol was treated with 3.45 g (150 mmol) of sodium metal with stirring until the sodium had completely dissolved. To the methoxide  
15    solution was added 3.00 g (25.6 mmol) of indole and 10.23 g (66.6 mmol) of 4-piperidone monohydrate hydrochloride. The mixture was refluxed for 16 h and then concentrated. The residue was dissolved in ethyl acetate and washed once with water. The aqueous phase was extracted three times with ethyl acetate and the combined organics were washed once with brine, dried over  
20    Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give 2.85 g of **48A** as a yellow solid.

B. 3-Piperidin-4-yl-1H-indole

A mixture of 400 mg (2.0 mmol) of **48A** in 100 mL of ethanol was hydrogenated at 45 psi for 17 h. The mixture was filtered through celite and concentrated to give 400 mg of **48B** as a white solid.

- 25    C. (R)-{1-[1-(1H-Indol-3-ylmethyl)-2-[4-(1H-indol-3-yl)-piperidin-1-yl]-2-oxo-ethyl]carbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

- According to General Procedure A, 100 mg (0.50 mmol) of **48B** and 195 mg (0.50 mmol) of **4C** were coupled and the product was purified by silica gel chromatography using a gradient elution of 75% ethyl acetate in hexane to  
30    87.5% ethyl acetate in hexane to give 220 mg of **48C** as a white solid.

D. 2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(1H-indol-3-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

- To 220 mg (0.39 mmol) of **48C** in 3 mL of ethanol was added 2 mL of concentrated HCl. The mixture was stirred at room temperature for 1 h and  
35    then concentrated. Methanol and a small amount of methylene chloride were added and the mixture was concentrated to give 200 mg of the title compound as a white solid.

-101-

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (1:1 mixture of rotamers) (partial) δ 8.32-8.45 (m, 1H), 7.53-7.69 (m, 1H), 7.30-7.43 (m, 3H), 6.94-7.29 (m, 6H), 5.23-5.37 (m, 1H), 4.46-4.63 (m, 1H), 3.90-4.04 (m, 1H), 1.62-1.68 (m, 6H).  
MS (CI, NH<sub>3</sub>) 471 (MH<sup>+</sup>)

5

**Example 49**

(R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(2-methyl-benzoimidazol-1-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

A. 4-(2-Methyl-benzoimidazol-1-yl)-piperidine-1-carboxylic acid tert-butyl ester

10 To 325 mg (3.00 mmol) of 2-methylbenzimidazole in 12 mL of DMSO was added 100 mg (3.00 mmol) of sodium hydride (60% oil dispersion) and the mixture was stirred at room temperature for 30 min. The mixture was heated at 70°C for 15 min and then cooled to room temperature. A solution of 520 mg (3.00 mmol) of 4-methanesulfonyloxy-piperidine-1-carboxylic acid tert-butyl  
15 ester (Yoon et al., WO9204342) in 3 mL of DMSO was added to the imidazole solution, and the mixture was stirred at room temperature for 12 h. The mixture was diluted with ethyl acetate and washed twice with a solution of saturated sodium bicarbonate and brine. The solution was dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (95:5 v/v  
20 chloroform:methanol) to give 89 mg of **49A** as a white solid.

B. 2-Methyl-1-piperidin-4-yl-1H-benzoimidazole trifluoroacetate

To 89 mg (0.28 mmol) of **49A** at 0°C was added 3 mL of trifluoroacetic acid and the mixture was stirred for 30 min. Dilution with ethyl acetate and concentration provided 92 mg of **49B** as a yellow syrup.

25 C. (R)-(1-{1-(1H-Indol-3-ylmethyl)-2-[4-(2-methyl-benzoimidazol-1-yl)-piperidin-1-yl]-2-oxo-ethyl}carbonyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 90 mg (0.42 mmol) of **49B** and 163 mg (0.42 mmol) of **4C** were coupled and the product was purified by silica gel  
30 chromatography using (75:25 v/v ethyl acetate:hexanes) followed by (95:5 v/v CH<sub>2</sub>Cl<sub>2</sub>:MeOH) to give 126 mg of **49C** as a white solid.

D. (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(2-methyl-benzoimidazol-1-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride

To 126 mg (0.21 mmol) of **49C** in 6 mL of ethanol was added 6 mL of  
35 concentrated HCl, and the mixture was stirred at room temperature for 80 min. The mixture was concentrated and the residue redissolved in a small amount of methanol. Ethyl acetate was added until the product precipitated from

-102-

solution, and the precipitated material was collected by filtration to give 85 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (1:1 mixture of rotamers) (partial) δ 7.53-7.81 (m, 4H), 7.19-7.40 (m, 2H), 7.00-7.19 (m, 3H), 2.91 (s, 1.5H), 2.81 (s, 1.5H), 1.59-1.68 (m, 6H).

5 MS (CI, NH<sub>3</sub>) 487 (MH<sup>+</sup>)

### Example 50

(R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

10 A. (1-Benzyl-piperidin-4-yl)-(2-nitro-phenyl)-amine

A mixture of 22.0 g (116 mmol) of 4-amino-1-benzylpiperidine, 16.31 g (116 mmol) of 1-fluoro-2-nitrobenzene, 19.17 g (116 mmol) of potassium iodide and 16.00 g (116 mmol) of potassium carbonate in 200 mL of DMSO was heated at 150°C for 16 h. The mixture was diluted with ethyl acetate and washed four times with water and once with brine. Drying over MgSO<sub>4</sub> and concentration provided the crude product, which was purified by silica gel chromatography (30:70 v/v ethyl acetate:hexanes) and crystallized from 10% ethyl acetate in hexane to give 27.0 g of **50A** as yellow crystals.

15 B. N-(1-Benzyl-piperidin-4-yl)-benzene-1,2-diamine

20 A mixture of 10.0 g (32.2 mmol) of **50A** and 1.00 g of 10% palladium on carbon in 50 mL of ethanol was hydrogenated at 50 psi for 3 h. The mixture was filtered through celite and concentrated to give 9.00 g of **50B** as a dark solid

25 C. 1-(1-Benzyl-piperidin-4-yl)-2-phenyl-1H-benzoimidazole

A mixture of 1.00 g (3.56 mmol) of **50B** and 1.89 g (17.8 mmol) of benzaldehyde in 4 mL of nitrobenzene was heated at 110°C for 16 h. The mixture was purified by silica gel chromatography (30:70 v/v ethyl acetate:hexanes) to give 550 mg of **50C** as a yellow foam.

30 D. 2-Phenyl-1-piperidin-4-yl-1H-benzoimidazole

To 540 mg (1.47 mmol) of **50C** in 2 mL of methylene chloride at -10°C was added 294 mg (2.06 mmol) of α-chloroethyl chloroformate and the mixture was stirred for 20 min and then concentrated. The residue was dissolved in 4 mL methanol and refluxed for 1 h. The mixture was concentrated to give 400 mg of **50D**.

35 E. (R)-(1-{1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}carbonyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester



-103-

According to General Procedure A, 70 mg (0.25 mmol) of **50D** and 98 mg (0.25 mmol) of **4C** were coupled and the product was purified by silica gel chromatography using a gradient elution of 50% ethyl acetate in hexane to 100% ethyl acetate 30 mg of **50E** was obtained as a white solid.

5 F. (R)-2-Amino-N-[1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenylbenzoimidazol-1-yl)-piperidin-1-yl]-ethyl]-isobutyramide hydrochloride

To 30 mg (0.046 mmol) of **50E** in 2 mL of ethanol was slowly added 2 mL of concentrated HCl with stirring. The mixture was stirred at room temperature for 75min and then concentrated. Crystallization from ethyl acetate/hexane provided 22 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (1:1 mixture of rotamers) (partial) δ 8.29 (d, 0.5H), 7.63-7.80 (m, 8.5H), 7.26-7.54 (m, 2.5H), 6.98-7.18 (m, 2.5H), 5.29-5.40 (m, 0.5H), 5.00-5.10 (m, 0.5H), 4.48-4.78 (m, 2H), 2.36-2.61 (m, 2H), 1.56-1.72 (m, 6H).

15 MS (CI, NH<sub>3</sub>) 549 (MH<sup>+</sup>)

**Example 51**

(R)-2-Amino-N-[2-[4-(6-fluoro-benzo[d]isoxazol-3-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide hydrochloride

A. 1-Acetyl-piperidine-4-carboxylic acid

20 A mixture of 129.2 g (1.00 mmol) of isonipecotic acid and 400 mL of acetic anhydride was refluxed for 2.5 h. The mixture was allowed to cool to room temperature and stirred for 17 h. The precipitated solids were filtered, washed with diethyl ether and dried under vacuum to give 150.1 g of **51A** as a white solid.

25 B. 1-Acetyl-piperidine-4-carbonyl chloride

To 250 mL of thionyl chloride was added 50.0 g (292 mmol) of **51A** with mechanical stirring. The mixture was stirred for 1 h and then 200 mL of petroleum ether was added. The precipitated solids were collected by filtration and washed with cold petroleum ether. The solids were dried under vacuum to give 53.97 g of **51B**.

C. 1-[4-(2,4-Difluoro-benzoyl)-piperidin-1-yl]-ethanone

To a suspension of 75.0 g (562 mmol) of aluminum chloride in 95 mL of m-difluorobenzene was added **51B** over 10 min. The mixture was refluxed for 3 h under nitrogen, and after cooling to room temperature the mixture was poured slowly onto ice. The product was extracted with methylene chloride and the combined organics were washed twice with water and once with brine. Drying over MgSO<sub>4</sub> provided 63.21 g of **51C** as a yellow solid.

-104-

D. (2,4-Difluoro-phenyl)-piperidin-4-yl-methanone hydrochloride

A mixture of 10.0 g (37.4 mmol) of **51C** in 40 mL each of concentrated HCl and glacial acetic acid was refluxed for 7 h. An additional 20 mL of concentrated HCl was added to the reaction mixture and refluxing was continued for an additional 6 h. The mixture was concentrated and the residue triturated with 2-propanol to give 8.0 g of **51D** as a white solid.

E. (2,4-Difluoro-phenyl)-piperidin-4-yl-methanone oxime

A mixture of 10.0 g (38.21 mmol) of **51D**, 2.66 g (38.2 mmol) of hydroxylamine hydrochloride, and 4.6 mL (33.0 mmol) of triethylamine in 150 mL of ethanol was stirred at room temperature for 30 min., and then refluxed for 2.5 h. During reflux, the product began precipitating from solution. After cooling to room temperature, filtration provided 6.7 g of **51E**.

F. 6-Fluoro-3-piperidin-4-yl-benzo[d]isoxazole

A mixture of 10.0 g (41.62 mmol) of **51E** and 250 mL of 50% NaOH was refluxed for 5 h. After cooling to room temperature, the mixture was diluted with toluene and washed once each with water and brine. The mixture was dried over MgSO<sub>4</sub> and concentrated to provide a residue, which was triturated with ethyl acetate and petroleum ether to give 0.56 g of **51F** as a solid.

G. (R)-{1-[2-[4-(6-Fluoro-benzo[d]isoxazol-3-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 11 mg (0.045 mmol) of **51F** and 18 mg (0.045 mmol) of **4C** were coupled and the product was purified by silica gel chromatography (95:5 v/v CH<sub>2</sub>Cl<sub>2</sub>:MeOH) to give 16 mg of **51G** as a white solid.

H. (R)-2-Amino-N-[2-[4-(6-fluoro-benzo[d]isoxazol-3-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide hydrochloride

To 16 mg (0.027 mmol) of **51G** in 2 mL of ethanol at room temperature was added 2 mL of concentrated HCl and the mixture stirred for 1 h. The mixture was concentrated and the residue crystallized from ethyl acetate/hexanes to give 11 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (1:1 mixture of rotamers) (partial) δ 8.21-8.29 (m, 1H), 7.51-7.63 (m, 2H), 7.26-7.34 (m, 2H), 6.90-7.21 (m, 5H), 5.07-5.24 (m, 1H), 4.43-4.49 (m, 1H), 3.79-3.92 (m, 1H), 2.58-2.74 (m, 2H), 1.44-1.53 (d, 6H).

MS (CI, NH<sub>3</sub>) 492 (MH<sup>+</sup>)

-105-

**Example 52**

(R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (4-hydroxy-butyl)-amide hydrochloride

A. 4-Phenyl-piperidine-1,4-dicarboxylic acid mono-tert-butyl ester

5 To a suspension of 42.84 g (114 mmol) of 4-phenyl-4-piperidine carboxylic acid 4-methylbenzenesulfonate in 250 mL of dioxane and 50 mL of water was added 200 mL of 1N NaOH, and the mixture was stirred until homogeneous. To the stirring mixture was added 27.3 g (125 mmol) of di-tert-butylidicarbonate over 15 min. Another 200 mL of 1N NaOH was added and  
10 the mixture was stirred until a pH of 10-11 was maintained. Stirring was continued for an additional 17 h at room temperature, then the solution was diluted with 350 mL of water and extracted three times with diethyl ether. The aqueous layer was acidified to pH 4-5 with glacial acetic acid, then extracted three times with ethyl acetate. The combined organics were washed three  
15 times with water and once with brine. Drying over MgSO<sub>4</sub> and concentration provided a clear oil, which was triturated with diethyl ether to give 32.3 g of **52A** as a white solid.

B. 4-Phenyl-piperidine-1,4-dicarboxylic acid benzyl ester tert-butyl ester

To 14.39 g (47 mmol) of **52A** in 200 mL of DMF was added 7.2 g (52  
20 mmol) of potassium carbonate followed by 8.22 g (48 mmol) of benzyl bromide. The mixture was stirred at room temperature for 17 h under nitrogen. The reaction mixture was diluted with 600 mL of ethyl acetate and washed five times with water and once with brine. Drying over MgSO<sub>4</sub> and concentration provided 15.95 g of **52B** as a clear oil that crystallized on standing.

25 C. 4-Phenyl-piperidine-4-carboxylic acid benzyl ester trifluoroacetate

To 4.5 g (11.4 mmol) of **52B** at 0°C was added 50 mL of cold trifluoroacetic acid and the mixture stirred for 1.5 h. The mixture was concentrated to give 5.10 g of **52C**.

30 D. (R)-1-[2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid benzyl ester

According to General Procedure A, 1.09 g (2.8 mmol) of **4C** and 885 mg (3.0 mmol) of **52C** were coupled and the product was purified by silica gel chromatography using a gradient elution of 0% to 3% methanol in methylene chloride. 1.04 g of **52D** was isolated as a light pink solid.

35 E. (R)-1-[2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid

A mixture of 1.0 g (1.5 mmol) of **52D**, and 200 mg of 20% palladium hydroxide on carbon in 30 mL of methanol was hydrogenated for 17 h at 50

-106-

psi. The mixture was filtered through celite and concentrated to give 772 mg of **52E** as an orange solid.

F. (R)-[1-[2-[4-(4-Hydroxy-butylcarbamoyl)-4-phenyl-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl]-carbamic acid tert-butyl ester

To a mixture of 66 mg (0.114 mmol) of **52E**, 9.4 mg (0.105 mmol) of 4-amino-1-butanol, and 27 mg (0.210 mmol) of diisopropylethylamine in 1.5 mL of methylene chloride was added 56 mg (0.126 mmol) of benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP) and the mixture was stirred overnight at room temperature. Concentration provided a residue which was dissolved in ethyl acetate and washed twice with 1N NaOH and once with brine. Drying over MgSO<sub>4</sub> and concentration provided the crude product, which was purified by silica gel chromatography using a gradient elution of 0% to 5% methanol in methylene chloride. 50 mg of **52F** was isolated as a white solid.

G. (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (4-hydroxy-butyl)-amide hydrochloride

According to General Procedure C, 44 mg (0.068 mmol) of **52F** was deprotected to give 30 mg of the title compound as a light pink solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.52-7.61 (m, 1H), 7.08-7.49 (m, 4H), 7.09-7.20 (m, 2H), 6.97-7.08 (m, 3H), 5.19-5.24 (m, 1H), 3.39-3.51 (m, 3H), 2.90-3.27 (m, 5H), 1.50-1.61 (m, 6H).

MS (CI, NH<sub>3</sub>) 548 (MH<sup>+</sup>)

-107-

**Example 53**

(R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid amide hydrochloride

5 A. (R)-{1-[2-(4-Carbamoyl-4-phenyl-piperidin-1-yl)-1-(1H-indol-3-ylmethyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

To a mixture of 255 mg (0.44 mmol) of **52E**, 118 mg (2.21 mmol) of ammonium chloride and 400 mg (3.09 mmol) of diisopropylethylamine in 4 mL of methylene chloride was added 235 mg (0.53 mmol) of BOP and the mixture stirred at room temperature for 24 h. The mixture was concentrated and the  
10 residue dissolved in ethyl acetate. The organic portion was washed twice with 1N NaOH and once with brine. Drying over MgSO<sub>4</sub> and concentration provided a crude product which was purified by silica gel chromatography using a gradient elution of 50% ethyl acetate in hexane to 100% ethyl acetate. 128 mg of **53A** was isolated as a white solid.

15 B. (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid amide hydrochloride

According to General Procedure C, 118 mg (0.20 mmol) of **53A** was deprotected to give 90 mg of the title compound as a pink solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.52-7.63  
20 (m, 1H), 7.19-7.47 (m, 4H), 6.98-7.17 (m, 5H), 5.13-5.25 (m, 1H), 3.82-4.21 (m, 1H), 3.98-3.27 (m, 3H), 1.50-1.65 (m, 6H).

MS (CI, NH<sub>3</sub>) 476 (MH<sup>+</sup>)

**Example 54**

25 (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid ethylamide hydrochloride

A. (R)-{1-[2-(4-Ethylcarbamoyl-4-phenyl-piperidin-1-yl)-1-(1H-indol-3-ylmethyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

To a mixture of 255 mg (0.44 mmol) of **52E**, 180 mg (2.21 mmol) of ethylamine hydrochloride and 400 mg (3.09 mmol) of diisopropylethylamine in  
30 4 mL of methylene chloride was added 235 mg (0.53 mmol) of BOP and the mixture was stirred at room temperature for 24 h. The mixture was concentrated and the residue was dissolved in ethyl acetate and washed twice with 1N NaOH and once with brine. Drying over MgSO<sub>4</sub> and concentration provided a crude product, which was purified by silica gel chromatography (1:1  
35 v/v ethyl acetate:hexanes) to give 167 mg of **54A** as a white foam.

B. (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid ethylamide hydrochloride

-108-

According to General Procedure C, 167 mg (0.28 mmol) of 54A was deprotected to give 132 mg of the title compound as a pink solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.50-7.65 (m, 1H), 6.95-7.32 (m, 9H), 5.07-5.22 (m, 1H), 4.11-4.22 (m, 1H), 3.56-3.67 (m, 1H), 2.88-3.30 (m, 5H), 2.61-2.79 (m, 1H), 1.52-1.61 (m, 6H).  
MS (CI, NH<sub>3</sub>) 504 (MH<sup>+</sup>)

### Example 55

(R)-1-Amino-cyclopropanecarboxylic acid {1-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide trifluoroacetate

10

A. (R)-(1-{1-Benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-cyclopropyl)-carbamic acid tert-butyl ester

According to General Procedure A, 20 mg (0.05 mmol) of 1B and 20 mg (0.10 mmol) of 1-tert-butoxycarbonylamino-cyclopropanecarboxylic acid (Kienzler et al., Helv. Chim. Acta (1992), 75 (4) pp 1078-84) were coupled and the product was purified by silica gel chromatography using a gradient elution of 45% ethyl acetate in hexane to 100% ethyl acetate to 2% methanol in ethyl acetate to give 10 mg of 55A.

20 B. (R)-1-Amino-cyclopropanecarboxylic acid {1-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide trifluoroacetate

To 10 mg (0.017 mmol) of 55A at 0°C was added 2 mL of cold trifluoroacetic acid and the mixture stirred for 1 h. The mixture was concentrated and then diluted with heptane and concentrated again. The product was crystallized from methylene chloride/hexane to give 7 mg of 55B as a solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) δ 7.19-7.40 (m, 5H), 6.89-7.12 (m, 4H), 5.16-5.24 (m, 1H), 4.45-4.78 (m, 4H), 4.17-4.30 (d, 1H), 3.64-3.81 (m, 2H), 3.15-3.30 (m, 1H), 2.72-2.99 (m, 1H), 2.50-2.67 (m, 0.5H), 2.37-2.45 (m, 1.5H), 1.74-1.89 (m, 2H), 1.50-1.74 (m, 2H), 1.26-1.45 (m, 2H).  
MS (CI, NH<sub>3</sub>) 478 (MH<sup>+</sup>)

### Example 56

(R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-1,2-dihydro-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

35

A. (1-Benzyl-piperidin-4-yl)-(3-nitro-pyridin-2-yl)-amine

A mixture of 1.66 g (10.5 mmol) of 2-chloro-3-nitropyridine, 2.00 g (10.5 mmol) of 4-amino-1-benzylpiperidine, 1.74 g (10.5 mmol) of potassium iodide

-109-

and 1.44 g (10.5 mmol) of potassium carbonate in 10 ml of DMSO was heated at 100°C for 17 h. After cooling to room temperature, the mixture was diluted with ethyl acetate and washed with saturated sodium bicarbonate solution and brine. Drying over MgSO<sub>4</sub> and concentration provided 1.25 g of **56A**.

5        B. N2-(1-Benzyl-piperidin-4-yl)-pyridine-2,3-diamine

A mixture of 1.25 g (4.16 mmol) of **56A** and 200 mg of 10% palladium on carbon in 150 mL of ethanol and 50 mL of ethyl acetate was hydrogenated at 50 psi for 3 h. The mixture was filtered through celite and concentrated to give 1.07 g of **56B**.

10       C. 3-(1-Benzyl-piperidin-4-yl)-1,3-dihydro-imidazo[4,5-b]pyridin-2-one

A mixture of 1.07 g (3.74 mmol) of **56B**, 2.43 g (15.0 mmol) of N,N'-carbonyldiimidazole, and 760 mg (7.52 mmol) of triethylamine in 10 mL of ethylene glycol dimethyl ether was heated at 80°C for 17 h. The mixture was diluted with ethyl acetate and washed with saturated sodium bicarbonate solution and brine. Drying over MgSO<sub>4</sub> and concentration provided a crude product which was purified by silica gel chromatography using a gradient elution of 75% ethyl acetate in hexane to 100% ethyl acetate to give 270 mg of **56C**.

20       D. 3-Piperidin-4-yl-1,3-dihydro-imidazo[4,5-b]pyridin-2-one

To a solution of 270 mg (6.88 mmol) of **56C** in 1 mL of 1,2-dichloroethane at 0°C was added 286 mg (2.0 mmol) of α-chloroethyl chloroformate. The ice bath was removed and the mixture stirred for 1 h. The mixture was concentrated and the residue dissolved in 10 mL of methanol and heated at 60°C for 3 h. The mixture was diluted with 2N NaOH and brine and extracted five times with chloroform. The organic extracts were dried over MgSO<sub>4</sub> and concentrated to give 70 mg of **56D**.

25       E. (R)-(1-(1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-1,2-dihydro-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethylcarbamoyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

30       According to General Procedure A, 70 mg (0.32 mmol) of **56D** and 125 mg (0.32 mmol) of **4C** were coupled and the product was purified by silica gel chromatography using a gradient elution of 70% ethyl acetate in hexane to 100% ethyl acetate to give 100 mg of **56E**.

35       F. (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-1,2-dihydro-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

To 100 mg (0.28 mmol) of **56E** in 3 mL of ethanol at 0°C was added 1.5 mL of concentrated HCl, the ice bath was removed and the mixture stirred for 2

-110-

h. The mixture was concentrated several times from ethanol. The residue was diluted with ethyl acetate/hexanes and the precipitated solid collected by filtration to give 52 mg of the title compound.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 300MHz) (mixture of rotamers) (partial)  $\delta$  7.97-8.06

5 (m, 1H), 7.53-7.69 (m, 2H), 7.25-7.38 (m, 2H), 7.09-7.21 (m, 3H), 5.22-5.33 (m, 1H), 4.49-4.74 (m, 2H), 1.48-1.64 (m, 6H).

MS ( $\text{Cl}$ ,  $\text{NH}_3$ ) 490 ( $\text{MH}^+$ )

### Example 57

10 (R)-2-Amino-N-[2-[4-(5-chloro-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide

A. (1-Benzyl-piperidin-4-yl)-(4-chloro-2-nitro-phenyl)-amine

A mixture of 10.0 g (53.0 mmol) of 4-amino-1-benzylpiperidine, 10.0 g (53.0 mmol) of 2,5-dichloronitrobenzene, 8.80 g (53.0 mmol) of potassium iodide, and 7.30 g (53 mmol) of potassium carbonate in 60 mL of DMSO was  
15 heated at  $140^\circ\text{C}$  for 48 h. The mixture was diluted with ethyl acetate and washed five times with water and once with brine. The mixture was dried over  $\text{MgSO}_4$  and concentrated. The residue was triturated with 30% ethyl acetate in hexane to give 8.3 g of **57A** as a red solid.

B. N1-(1-Benzyl-piperidin-4-yl)-4-chloro-benzene-1,2-diamine

20 A mixture of 8.30 g (26.28 mmol) of **57A** was hydrogenated for 3 h at 45 psi. The mixture was filtered through celite and concentrated. The product was purified by silica gel chromatography using a gradient elution of 100% ethyl acetate to 10% diethylamine in ethyl acetate to give 6.41 g of the title compound.

25 C. 1-(1-Benzyl-piperidin-4-yl)-5-chloro-1,3-dihydro-benzoimidazol-2-one

A mixture of 1.00 g (3.50 mmol) of **57B**, 2.84 g (17.5 mmol) of  $\text{N,N}'$ -carbonyldiimidazole, and 764 mg (7.0 mmol) of triethylamine in 3 mL of ethylene glycol dimethyl ether was heated at  $92^\circ\text{C}$  for 5 h. The mixture was  
30 concentrated and the product was purified by silica gel chromatography (75:25 v/v ethyl acetate:hexanes) to give 680 mg of **57C** as a pink solid.

D. 5-Chloro-1-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one

To a solution of 680 mg (2.18 mmol) of **57C** in 3 mL of methylene chloride at  $-10^\circ\text{C}$  was added 468 mg (3.27 mmol) of  $\alpha$ -chloroethyl  
35 chloroformate and the mixture stirred for 40 min. Concentration provided a dissolved residue which was in 10 mL of methanol and warmed to reflux for 1 h. The mixture was concentrated and 50 mL of chloroform was added to the residue followed by a solution of 90 mg of sodium hydroxide in 10 mL of water.



-111-

The mixture was stirred for 15 min. The layers were separated and the organic portion was washed once with brine. Drying over MgSO<sub>4</sub> and concentrations provided 510 mg of **57D** as a pink solid.

- E. (R)-{1-[2-[4-(5-Chloro-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 40 mg (0.16 mmol) of **57D** and 62 mg (0.16 mmol) of **4C** were coupled and the product was purified by silica gel chromatography eluting with 50% ethyl acetate in hexane followed by 5% methanol in methylene chloride to give 31 mg of **57E** as a white solid.

- F. (R)-2-Amino-N-[2-[4-(5-chloro-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide

To a mixture of 31 mg (0.05 mmol) of **57E** in 3 mL of ethanol was added 3 mL of concentrated HCl and the mixture was stirred at room temperature for 80 min. The mixture was concentrated, and the residue crystallized from ethyl acetate/hexane to give 26 mg of the title compound as a white solid.

- <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 8.19-8.30 (m, 1H), 7.42-7.64 (m, 1H), 6.94-7.37 (m, 8H), 5.05-5.30 (m, 1H), 4.12-4.54 (m, 3H), 2.37-2.61 (m, 2H), 1.48-1.64 (m, 6H).
- MS (CI, NH<sub>3</sub>) 523, 525 (MH<sup>+</sup>)

### Example 58

(R)-2-Amino-N-{1-benzyloxymethyl-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethyl}-isobutyramide

- A. (R)-{1-[1-Benzyloxymethyl-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

To a mixture of 42 mg (0.846 mmol) of **6C** and 10.3 mg (0.102 mmol) of triethylamine in 5 mL of methylene chloride at 0°C was added 18 mg (0.93 mmol) of toluenesulfonyl chloride and the mixture was allowed to warm to room temperature and stirred overnight. The residue was dissolved in ethyl acetate and washed once each with 1N NaOH and brine, dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (60:40 v/v ethyl acetate:hexanes) to give 19 mg of **58A** as a foam.

- B. (R)-2-Amino-N-{1-benzyloxymethyl-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethyl}-isobutyramide

According to General Procedure C, 19 mg (0.029 mmol) of **58A** was deprotected to give 12 mg of the title compound.

-112-

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (1:1 mixture of rotamers) (partial) δ 7.46-7.67 (m, 2H), 6.79-7.29 (m, 11H), 5.06-5.21 (m, 1H), 2.29-2.32 (d, 3H), 1.48-1.60 (bs, 6H).

MS (CI, NH<sub>3</sub>) 565 (MH<sup>+</sup>)

5

**Example 59**

(R)-2-Amino-N-{1-benzo[b]thiophen-3-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

A. (R)-3-Benzo[b]thiophen-3-yl-2-(2-tert-butoxycarbonylamino-2-methyl-propionylamino)-propionic acid

10

A mixture of 193 mg (0.64 mmol) of D-3-(3-benzothieryl)alanine trifluoroacetate, 180 mg (0.53 mmol) of **33A**, and 219 mg (1.70 mmol) of diisopropylethylamine in 10 mL of DMF was stirred overnight at room temperature. The mixture was acidified with 1N HCl and diluted with ethyl acetate. The aqueous phase was extracted three times with ethyl acetate and

15

the combined organics were washed five times with water, dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography, eluting with 100% chloroform, followed by 5% methanol in chloroform, followed by (9:1:0.1 v/v/v chloroform: methanol: acetic acid) to give 110 mg of **59A**.

20

B. (R)-(1-{1-Benzo[b]thiophen-3-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}carbonyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

25

According to General Procedure A, 58 mg (0.27 mmol) of 4-(2-keto-1-benzimidazoliny) piperidine and 105 mg (0.27 mmol) of **59A** were coupled and the product was purified by silica gel chromatography using a gradient elution of 20% hexane in ethyl acetate to 100% ethyl acetate. 95 mg of **59B** was isolated as a foam.

30

C. (R)-2-Amino-N-{1-benzo[b]thiophen-3-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

According to General Procedure C, 85 mg (0.14 mmol) of **59B** was deprotected to give 55 mg of the title compound as an off-white solid.

35

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (1:1 mixture of rotamers) (partial) δ 7.83-7.99 (m, 2H), 7.30-7.48 (m, 3H), 7.20-7.28 (m, 0.5H), 7.51-7.26 (m, 3H), 6.86-6.92 (m, 0.5H), 5.29-5.45 (m, 1H), 4.57-4.71 (m, 1H), 4.29-4.44 (m, 1H), 3.96-4.09 (m, 1H), 2.20-2.83 (m, 3H), 1.55-1.64 (d, 3H), 1.44-1.51 (d, 3H).

MS (CI, NH<sub>3</sub>) 506 (MH<sup>+</sup>)

-113-

**Example 60**

(R)-2-Amino-N-{1-(4-benzyloxy-benzyl)-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethyl}-isobutyramide hydrochloride

A. 1,2,3,4-Tetrahydro-isoquinolin-7-ylamine

- 5 A mixture of 500 mg (2.8 mmol) of **6A** and 500 mg of 10% palladium on carbon was hydrogenated at 45 psi overnight. The mixture was filtered through celite and concentrated to give 350 mg of **60A** as a white solid.

B. (R)-{1-[2-(7-Amino-3,4-dihydro-1H-isoquinolin-2-yl)-1-(4-benzyloxy-benzyl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

- 10 According to General Procedure A, 300 mg (0.68 mmol) of **44D** and 100 mg (0.68 mmol) of **60A** were coupled and the product was purified by silica gel chromatography (75:25 v/v ethyl acetate:hexanes) to give 230 mg of **60B** as a white solid.

- 15 C. (R)-{1-[1-(4-Benzyloxy-benzyl)-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

- A mixture of 220 mg (0.39 mmol) of **60B**, 85 mg (0.44 mmol) of p-toluenesulfonyl chloride and 55 mg (0.44 mmol) of 4-dimethylaminopyridine in 5 mL of methylene chloride was stirred overnight at room temperature. The  
20 mixture was diluted with chloroform and washed twice each with 10% HCl solution, saturated sodium bicarbonate solution and brine. Drying over MgSO<sub>4</sub> and concentration provided a crude product, which was purified by silica gel chromatography (75:25 v/v ethyl acetate:hexanes) to give 200 mg of **60C**.

- 25 D. (R)-2-Amino-N-{1-(4-benzyloxy-benzyl)-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethyl}-isobutyramide hydrochloride

- To a solution of 150 mg (0.21 mmol) of **60C** in 3 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room  
30 temperature for 1 h. The mixture was concentrated, and the residue crystallized from methylene chloride/hexane to give 112 mg of the title compound.

- <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (1:1 mixture of rotamers) (partial) δ 7.64 (d, 2H), 7.17-7.39 (m, 7H), 7.04-7.10 (m, 1H), 6.79-6.92 (m, 5H), 4.90-5.12 (m, 2H),  
35 4.29-4.57 (m, 1.5H), 4.11-4.20 (m, 0.5H), 2.82-2.97 (m, 2H), 2.34 (d, 3H), 1.46-1.52 (m, 6H).

MS (Cl, NH<sub>3</sub>) 641 (MH<sup>+</sup>)

-114-

**Example 61**

(R)-2-Amino-N-[1-(7-amino-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-isobutyramide dihydrochloride

5 A. (R)-2-Amino-N-[1-(7-amino-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-isobutyramide dihydrochloride

According to General Procedure C, 29 mg (0.056 mmol) of **36B** was deprotected to give 14 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (1:1 mixture of rotamers) (partial) δ 7.51-7.60 (m, 1H), 7.24-7.31 (d, 0.5H), 6.91-7.18 (m, 6.5H), 5.14-5.33 (m, 1H), 4.42-10 4.68 (m, 1.5H), 4.12-4.23 (m, 0.5H), 2.53-2.75 (m, 1H), 2.35-2.51 (m, 0.5H), 2.08-2.22 (m, 0.5H), 1.52-1.63 (m, 6H).

MS (CI, NH<sub>3</sub>) 420 (MH<sup>+</sup>)

**Example 62**

15 (R)-2-Amino-N-[1-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carbonyl]-4-phenyl-butyl]-isobutyramide hydrochloride

A. 2-Oxo-5,6-diphenyl-3-(3-phenyl-allyl)-morpholine-4-carboxylic acid

To a -78°C solution of 13.8 g (70.0 mmol) of cinnamyl bromide and 4.94 g (14.0 mmol) of t-butyl-(2S, 3R)-(+)-6-oxo-2,3-diphenyl-4-morpholine carboxylate in 350 mL of anhydrous THF was added

20 28 mL (28 mmol) of 1M sodium bistrimethylsilylamide in THF. The mixture was stirred at -78°C for 1.5 h and then poured into 750 mL of ethyl acetate. The mixture was washed twice with brine, dried over MgSO<sub>4</sub> and concentrated to give a yellow oil. The oil was stirred in 150 mL of hexane overnight and the precipitated solid was then collected by filtration to give 3.2 g of **62A** as a white  
25 solid.

B. 5,6-Diphenyl-3-(3-phenyl-allyl)-morpholin-2-one

According to General Procedure B, 2.97 g (6.33 mmol) of **62A** was deprotected to give an orange oil which was purified by silica gel chromatography (10:90 v/v ethyl acetate:hexane) to give 880 mg of **62B** as a  
30 white solid.

C. 2-(2-Amino-2-methyl-propionylamino)-5-phenyl-pentanoic acid

A mixture of 440 mg (1.19 mmol) of **62B** and 120 mg of palladium chloride in 20 mL of ethanol and 10 mL of THF was hydrogenated at 45 psi for 16 h. The mixture was filtered through celite and concentrated, and the residue  
35 was triturated with ether to give 240 mg of **62C** as a white solid.

D. (R)-2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-5-phenyl-pentanoic acid

-115-

A mixture of 203 mg (1.05 mmol) of **33A**, 378 mg (1.26 mmol) of **62C** and 434 mg (3.36 mmol) of diisopropylethylamine in 2 mL of DMF was stirred overnight. The mixture was diluted with ethyl acetate and extracted twice with 1N HCl. The aqueous phase was extracted once with ethyl acetate. The pooled organic extracts were washed three times with water and once with brine. The mixture was dried over MgSO<sub>4</sub> and concentrated. The residue was purified by silica gel chromatography using 80% chloroform in hexane followed by 100% chloroform followed by 10% methanol in chloroform to give 127 mg of **62D**.

5 **E. (R)-(1-Methyl-1-(1-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carbonyl]-4-phenyl-butylcarbamoyl)-ethyl)-carbamic acid tert butyl ester**

According to General Procedure A, 45 mg (0.12 mmol) of **62D** and 26 mg (0.12 mmol) of 4-(2-keto-1-benzimidazoliny) piperidine were coupled and the product purified by silica gel chromatography using a gradient of 100% methylene chloride to 3% methanol in methylene chloride to give 39 mg of **62E** as a yellow foam.

15 **F. (R)-2-Amino-N-{1-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carbonyl]-4-phenyl-butyl}-isobutyramide hydrochloride**

20 According to General Procedure C, 36 mg (0.062 mmol) of **62E** was deprotected to give 28 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 7.03-7.36 (m, 9H), 4.62-4.73 (m, 1H), 4.43-4.59 (m, 1H), 3.97-4.19 (m, 1H), 2.52-2.87 (m, 3H), 1.60 (s, 6H).

25 MS (CI, NH<sub>3</sub>) 578 (MH<sup>+</sup>)

### Example 63

**2-Amino-N-{1-(1-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride**

30 **A. 2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(1-methyl-1H-indol-3-yl)-propionic acid**

A mixture of 1.00 g (4.58 mmol) of 1-methyl-dl-tryptophan, 1.65 g (5.5 mmol) of **33A**, and 1.30 g (10.0 mmol) of diisopropylethylamine in 2 mL of DMF was heated overnight at 50°C. The mixture was diluted with ethyl acetate and washed once each with 10% HCl and brine. The solution was dried over MgSO<sub>4</sub> and concentrated, and the product was purified using a gradient of 2% to 5% methanol in methylene chloride to give 173 mg of **63A**.

-116-

B. (1-Methyl-1-((1-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl)-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 173 mg (0.42 mmol) of **63A** and 131 mg (0.42 mmol) of 4-(2-keto-1-benzimidazoliny) piperidine were coupled and the product was purified by silica gel chromatography in 100% ethyl acetate to give 210 mg of **63B**.

C. 2-Amino-N-((1-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-isobutyramide hydrochloride

According to General Procedure C, 209 mg (0.35 mmol) of **63B** was deprotected to give 166 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) (1:1 mixture of diastereomers) δ 7.67 (d, 0.5H), 7.55 (d, 0.5H), 7.31-7.39 (m, 1H), 6.99-7.26 (m, 6.5H), 6.74 (d, 0.5H), 5.12-5.33 (m, 1H), 3.78 (s, 3H), 1.49-1.63 (m, 6H). MS (CI, NH<sub>3</sub>) 503 (MH<sup>+</sup>)

#### Example 64

(R)-2-Amino-N-((1-naphthalen-1-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-isobutyramide trifluoroacetate

A. (R)-2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-naphthalen-1-yl-propionic acid

A mixture of 500 mg (1.52 mmol) of D-3-(1'-naphthyl)alanine, 547 mg (1.82 mmol) of **33A** and 0.80 mL (4.56 mmol) of diisopropylamine in 5 mL of DMF was heated at 50°C overnight. The mixture was concentrated and the residue diluted with methylene chloride. The organic portion was washed once with water, twice with 1N HCl and once with brine. The solution was dried over MgSO<sub>4</sub> and concentrated, and the product was purified by silica gel chromatography (0.1:5:95 v/v/v acetic acid:methanol:methylene chloride) to give 484 mg of **64A**.

B. (R)-((1-Methyl-1-((1-naphthalen-1-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl)-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 100 mg (0.25 mmol) of **64A** and 50 mg (0.23 mmol) of 4-(2-keto-1-benzimidazoliny)-piperidine were coupled and the product was purified by silica gel chromatography using a gradient of 100% methylene chloride to 4% methanol in methylene chloride to give 97 mg of **64B**.

-117-

C. (R)-2-Amino-N-{1-naphthalen-1-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide trifluoroacetate

To 91 mg (0.15 mmol) of **64B** at 0°C was added 2 mL of cold TFA and the mixture was stirred at 0°C for 2 h. The mixture was concentrated, then coevaporated twice each from methylene chloride and toluene to give 99 mg of the title compound as a white powder.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 8.22 (d, 0.5H), 8.17 (d, 0.5 H), 7.79-7.94 (m, 2H), 7.39-7.67 (m, 4H), 7.01-7.26 (m, 3.5H), 6.75 (d, 0.5H), 5.42-5.51 (m, 1H), 5.29-5.40 (m, 1H), 4.54-4.67 (m, 1H), 4.19-4.37 (m, 2H), 3.51-3.82 (m, 3H), 2.29-2.72 (m, 3H), 1.49-1.68 (m, 6H). MS (CI, NH<sub>3</sub>) 500 (MH<sup>+</sup>)

### Example 65

(R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[6-(morpholine-4-carbonyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide hydrochloride

A. (R)-(1-{2-(1H-Indol-3-yl)-1-[6-(morpholine-4-carbonyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}carbonyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to the method outlined in General Procedure A, 121 mg (0.22 mmol) of **26A** and 19 mg (0.33 mmol) of morpholine were coupled to give 140 mg of **65A** as a yellow oil.

B. (R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[6-(morpholine-4-carbonyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide hydrochloride

To 86 mg (0.14 mmol) of **65A** in 6 mL of ethanol was added 2.0 mL of concentrated HCl and the mixture was stirred for 1.5 h at room temperature. The mixture was concentrated to give 68 mg of the title compound as a tan solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.51-7.61 (m, 1H), 6.92-7.30 (m, 6.5H), 6.58 (d, 0.5H), 5.14-5.39 (m, 1H), 4.56-4.66 (d, 0.5H), 4.41-4.50 (d, 1H), 4.09-4.18 (d, 0.5H).

### Example 66

(R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-imidazo[4,5-c]pyridin-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide dihydrochloride

A. 1-Piperidin-4-yl-1,3-dihydro-imidazo[4,5-c]pyridin-2-one

**66A** was prepared by the same route illustrated in Example 56 using 4-chloro-3-nitro-pyridine and 4-amino-1-benzylpiperidine as starting materials. The 4-chloro-3-nitro-pyridine was prepared by adding 3.00 g (23.8 mmol) of 4-

-118-

- hydroxy-3-nitropyridine to 10 mL of phosphorous oxychloride and heating the mixture at 130°C for 3 h. The mixture was poured into ice water, neutralized with 2N NaOH and allowed to stand for 1 h. The mixture was extracted with ethyl acetate and the combined organics were washed with brine. The solution
- 5 was dried and concentrated to give 2.92 g of 4-chloro-3-nitro-pyridine.

B. (R)-(1-{1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-imidazo[4,5-c]pyridin-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

- According to General Procedure A, 90 mg (0.29 mmol) of 66A and 113
- 10 mg (0.29 mmol) of 4C were coupled and the product was purified by silica gel chromatography using an elution gradient of 1% methanol in methylene chloride to 5% methanol in methylene chloride followed by 5% methanol plus 1% acetic acid in methylene chloride to give 80 mg of 66B.

- C. (R)-(1-{1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-imidazo[4,5-c]pyridin-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester dihydrochloride
- 15

- To 80 mg (0.14 mmol) of 66B in 3 mL of ethanol was added 3 mL of concentrated HCl and the mixture was stirred at room temperature for 3 h. The mixture was concentrated and then coevaporated three times with ethanol.
- 20 The residue was crystallized from ethyl acetate/hexane to give 60 mg of the title compound.

- <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 8.41-8.57 (m, 2H), 7.87 (d, 0.5H), 7.58 (d, 0.5H), 7.21-7.39 (m, 2H), 7.07-7.13 (m, 3H), 5.24-5.34 (m, 0.5H), 5.12-5.20 (m, 1H), 4.60-4.71 (m, 1H), 1.52-1.67 (m, 6H).
- 25 MS (CI, NH<sub>3</sub>) 491 (MH<sup>+</sup>)

### Example 67

2-Amino-N-{1-biphenyl-4-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

- A. (R)-2-Amino-3-biphenyl-4-yl-propionic acid trifluoroacetate
- 30 To 1.50 g (4.4 mmol) of BOC-D-4-4'-phenylalanine at 0°C was added 25 mL of cold TFA and the mixture was stirred for 2 h. The mixture was concentrated and then heptane was added to the residue and the mixture concentrated again and dried under vacuum to give 1.59 g of 67A as a pale red solid.

- B. (R)-3-Biphenyl-4-yl-2-(2-tert-butoxycarbonylamino-2-methyl-propionylamino)-propionic acid
- 35

A mixture of 1.59 g (4.4 mmol) of 67A, 1.59 g (5.3 mmol) of 33A and 1.82 g (14.1 mmol) of diisopropylethylamine in 50 mL of DMF was stirred at



-119-

0°C for 6 days. The mixture was diluted with ethyl acetate, washed twice with 10% HCl and once with water, and twice with brine. The solution was dried over MgSO<sub>4</sub>, filtered and concentrated and the product was purified by silica gel chromatography using a gradient of 100% ethyl acetate to 10% methanol in ethyl acetate to give 900 mg of **67B**.

C. (R)-(1-(1-Biphenyl-4-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 250 mg (0.59 mmol) of **67B** and 127 mg (0.059 mmol) of 4-(2-keto-1-benzimidazoliny)-piperidine were coupled to give 270 mg of **67C** as a white solid which was used in the next step without further purification.

D. (R)-2-Amino-N-[1-biphenyl-4-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl]-isobutyramide hydrochloride

To 270 mg (0.43 mmol) of **67C** in 3 mL of ethanol was added 3 mL of concentrated HCl and the mixture was stirred at room temperature for 1.5 h. The mixture was concentrated to dryness and the residue was triturated with hexane/ethanol to give 200 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 7.52-7.70 (m, 4H), 7.28-7.51 (m, 6H), 6.91-7.11 (m, 3H), 5.25-5.33 (m, 1H), 4.67-4.80 (m, 1H), 4.43-4.58 (m, 1H), 4.16-4.27 (m, 1H), 1.53-1.66 (m, 6H). MS (CI, NH<sub>3</sub>) 527 (MH<sup>+</sup>)

### Example 68

(R)-2-Amino-N-[2-(1H-indol-3-yl)-1-(7-sulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-ethyl]-isobutyramide hydrochloride

A. (R)-(1-[2-(1H-Indol-3-yl)-1-(7-sulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-ethylcarbamoyl]-1-methyl-ethyl)-carbamic acid tert-butyl ester

The preparation of 1,2,3,4-tetrahydro-isoquinoline-7-sulfonic acid amide is described by Pendleton et al., J. Pharmacol. Exp. Ther., 208 (1979) p24.

According to General Procedure A, 83 mg (0.39 mmol) of 1,2,3,4-Tetrahydro-isoquinoline-7-sulfonic acid amide and 150 mg (0.39 mmol) of **4C** were coupled and the product was purified by silica gel chromatography (19:1 v/v chloroform:methanol) to give 206 mg of **68A**.

B. (R)-2-Amino-N-[2-(1H-indol-3-yl)-1-(7-sulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-ethyl]-isobutyramide hydrochloride

To 144 mg (0.25 mmol) of **68A** in 12 mL of ethanol was added 4 mL of concentrated HCl and the mixture was stirred at room temperature for 5 h. The

-120-

reaction mixture was concentrated to give 118 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.48-7.53 (m, 2H), 7.21-7.30 (m, 1H), 6.92-7.13 (m, 4H), 5.11-5.25 (m, 1H), 4.41-4.62 (m, 2H), 4.11-4.22 (d, 0.5H), 3.75-3.86 (m, 1H), 3.50-3.63 (m, 1H), 2.53-2.72 (m, 1H), 2.26-2.42 (m, 0.5H), 2.06-2.19 (m, 0.5H), 1.57 (m, 6H).  
MS (Cl, NH<sub>3</sub>) 519 (MH<sup>+</sup>)

### Example 69

10 2-Amino-N-{1-(6-fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-  
isobutyramide hydrochloride

A. 2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-(6-fluoro-1H-indol-3-yl)-propionic acid

15 A mixture of 150 mg (0.50 mmol) of **33A**, 100 mg (0.45 mmol) of 6-fluorotryptophan and 128 mg (1.0 mmol) of diisopropylethylamine was stirred overnight at room temperature. The mixture was acidified to pH 1 with 1N HCl, diluted with water and extracted three times with ethyl acetate. The combined organic extracts were washed four times with water, once with brine, dried over MgSO<sub>4</sub> and concentrated to give 154 mg of **69A**.

20 B. (1-{1-(6-Fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}carbonyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

25 According to General Procedure A, 79 mg (0.194 mmol) of **69A** and 42 mg (0.194 mmol) of 4-(2-keto-1-benzimidazolyl)-piperidine were coupled and the product was purified by silica gel chromatography using a gradient of 1% methanol in methylene chloride to 3% methanol in methylene chloride to give 84 mg of **69B** as a white foam.

C. 2-Amino-N-{1-(6-fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

30 According to General Procedure C, 78 mg (0.129 mmol) of **69B** was deprotected to give 54 mg of the title compound.

-121-

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) (1:1 mixture of diastereomers) δ 7.56-7.62 (m, 0.5H), 7.46-7.53 (m, 0.5H), 7.01-7.26 (m, 6H), 6.70-6.88 (m, 1H), 5.17-5.37 (m, 1H), 4.60-4.69 (m, 1H), 4.25-4.41 (m, 1H), 3.94-4.12 (m, 1H), 3.04-3.24 (m, 2H), 2.45-2.62 (m, 2H), 1.48-1.65 (m, 6H).

5 MS (CI, NH<sub>3</sub>) 519 (MH<sup>+</sup>)

### Example 70

(R)-2-Amino-N-[1-(1H-indol-3-ylmethyl)-2-(4-indol-1-yl-piperidin-1-yl)-2-oxo-ethyl]-isobutyramide hydrochloride

#### A. (1-Benzyl-piperidin-4-yl)-phenyl-amine

10 To a mixture of 10.9 g (57.6 mmol) of 1-benzyl-4-piperidone, 100.0 g (576 mmol) of sodium sulfate and 175 mL of acetic acid under nitrogen was added 7.00 g (74.9 mmol) of aniline by syringe and the mixture was stirred at room temperature for 15 min. To the stirring solution was added 61.0 g (288 mmol) of sodium triacetoxymethylborohydride and the mixture was stirred overnight.

15 The mixture was concentrated and the residue was poured onto ice and neutralized with 2N NaOH to pH 7.5. The mixture was extracted four times with chloroform and the organics washed once with brine. The solution was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was triturated with ether to give 4.2 g of **70A** as a white solid.

#### 20 B. 1-[2-(1-Benzyl-piperidin-4-ylamino)-phenyl]-2-chloro-ethanone hydrochloride

To a flame-dried flask containing 2.50 g (9.4 mmol) of **70A**, and 40 mL of dichloroethane at 0°C was added 11.3 mL (11.3 mmol) of 1M boron trichloride in methylene chloride over 5 min., followed by 1.2 mL (18.8 mmol) of  
25 chloroacetonitrile over 3 min. The mixture was warmed to room temperature and stirred for 10 min. The mixture was refluxed overnight and then allowed to cool to room temperature. Ice was added to the stirring solution followed by 10 mL of 10% HCl and the mixture was heated to 100°C for 0.5 h. The layers were separated and the aqueous phase was extracted with methylene  
30 chloride. The organic portion was dried over MgSO<sub>4</sub> and concentrated to give 1.50 g of **70B**.

#### C. 1-(1-Benzyl-piperidin-4-yl)-1H-indole

To 1.50 g (4.0 mmol) of **70B** dissolved in 11 mL of ethanol at 0°C was added 2.0 mL (4.0 mmol) of 2N NaOH and the mixture stirred for 10 min.  
35 Sodium borohydride, 80 mg (2.1 mmol), was added to the stirring solution and the mixture was stirred at 0°C for 1 h. Ice was added to the solution and the mixture was extracted three times with methylene chloride. The combined organics were washed with water and dried over MgSO<sub>4</sub> and the residue

-122-

dissolved in 1,4-dioxane and refluxed for 2 h. After cooling to room temperature, ice and a saturated aqueous sodium carbonate solution was added to the reaction mixture and the solution was extracted three times with methylene chloride and the combined organics were washed twice with water.

- 5 The solution was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography (80:20 v/v hexanes:ethyl acetate) to give 350 mg of **70C** as a white foam.

D. 4-Indol-1-yl-piperidine-1-carboxylic acid ethyl ester

- To a refluxing solution of 350 mg (1.2 mmol) of **70C** in 40 mL of dichloroethane was added 0.28 mL (3.6 mmol) of ethyl chloroformate and the mixture was refluxed for 2.5h. The mixture was concentrated to give 330 mg of **70D** as a yellow solid.

E. 1-Piperidin-4-yl-1H-indole

- A mixture of 325 mg (1.2 mmol) of **70D**, 440 mg (15.7 mmol) of sodium hydroxide, 1 mL of water and 2 mL of ethylene glycol was heated at 150°C for 17 h. After cooling to room temperature, ice and concentrated HCl were added to the mixture and the solution was stirred for 30 min. Ice and saturated aqueous sodium carbonate solution were added to the mixture until the solution was basic. The mixture was extracted twice with ether and the combined organic extracts were washed twice with water. The solution was dried over MgSO<sub>4</sub> and concentrated to give 150 mg of **70E** as a colorless oil.

F. (R)-{1-[1-(1H-Indol-3-ylmethyl)-2-(4-indol-1-yl-piperidin-1-yl)-2-oxo-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

- According to General Procedure A, 150 mg (0.75 mmol) of **70E**, and 291 mg (0.75 mmol) of **4C** were coupled to give 260 mg of **70F** as a white solid.

G. (R)-2-Amino-N-[1-(1H-indol-3-ylmethyl)-2-(4-indol-1-yl-piperidin-1-yl)-2-oxo-ethyl]-isobutyramide hydrochloride

- To 250 mg (0.44 mmol) of **70F** in 5 mL of ethanol was added 3 mL of concentrated HCl and the mixture was stirred at room temperature for 2 h. The mixture was concentrated and ethanol and hexane were added to the residue. The precipitated off-white solid was filtered and dried to give 10 mg of the title compound.

MS (Cl, NH<sub>3</sub>) 473 (MH<sup>+</sup>)

35

**Example 71**

(R)-2-Amino-N-[1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethyl]-isobutyramide dihydrochloride

A. 2-Phenyl-3-piperidin-4-yl-3H-imidazo[4,5-b]pyridine

-123-

71A was prepared according to the method outlined in Example 50 using 4-amino-1-benzylpiperidine and 2-chloro-3-nitro-pyridine as starting materials.

5 B. (R)-(1-{1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 90 mg (0.32 mmol) of 71A and 161 mg (0.41 mmol) of 4C were coupled and the product was purified by silica gel chromatography (100% ethyl acetate) to give

10 45 mg of 71B.

C. (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide dihydrochloride

To 45 mg (0.069 mmol) of 71B in 2 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 2 h. The mixture was concentrated and coevaporated twice with ethanol. Ethyl acetate and hexane were added to the residue until the product crystallized. The solid was collected by filtration to give 20 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 8.74 (d, 0.5H), 8.56 (d, 0.5H), 8.06-8.23 (m, 1H), 7.43-7.79 (m, 8H), 7.18-7.28 (m, 2H), 6.87-7.02 (m, 3H), 5.14-5.28 (m, 1H), 4.52-4.63 (m, 1H), 3.93-4.08 (m, 1H), 1.37-1.49 (d, 6H).

MS (CI, NH<sub>3</sub>) 551 (MH<sup>+</sup>)

### Example 72

25 2-Amino-N-{1-(6-fluoro-1H-indol-3ylmethyl)-2-oxo-2-[4-(2-phenyl-benzimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

A. (1-{6-Fluoro-1H-indol-3ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 101 mg (0.25 mmol) of 69A and 70 mg (0.25 mmol) of 50D were coupled and the product was purified by silica gel chromatography (90:10 v/v chloroform:methanol) to give 84 mg of 72A.

B. 2-Amino-N-{1-(6-fluoro-1H-indol-3ylmethyl)-2-oxo-2-[4-(2-phenyl-benzimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

According to General Procedure C, 76 mg (0.114 mmol) of 72A was deprotected to give 58 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) δ (mixture of rotamers) (partial) (1:1 mixture of diastereomers) 8.21-8.30 (m, 0.5H), 7.65-7.88 (m, 8.5H), 7.44-7.51 (m, 0.5H), 7.27-7.33 (m, 1H), 6.99-7.14 (m, 1.5H), 6.72-6.95 (m, 1H), 5.27-5.35

-124-

(m, 0.5H), 5.02-5.10 (m, 0.5H), 4.52-4.79 (m, 2H), 2.34-2.69 (m, 2H), 1.94-2.16 (m, 2H), 1.51-1.71 (m, 6H).  
MS (CI, NH<sub>3</sub>) 568 (MH<sup>+</sup>)

**Example 73**

5     (R)-2-Amino-N-[4-phenyl-1-[4-(2-phenyl-benzoimidazol-1-yl)-piperidine-1-carbonyl]-butyl]-isobutyramide hydrochloride

A. (R)-(1-Methyl-1-[4-phenyl-1-[4-(2-phenyl-benzoimidazol-1-yl)-piperidine-1-carbonyl]-butylcarbamoyl]-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 112 mg (0.30 mmol) of **62D** and 84  
10 mg (0.30 mmol) of **50D** were coupled and the product was purified by silica gel chromatography using a gradient elution of 1% to 10% methanol in methylene chloride to provide 55 mg of **73A** as a white foam.

B. (R)-2-Amino-N-[4-phenyl-1-[4-(2-phenyl-benzoimidazol-1-yl)-piperidine-1-carbonyl]-butyl]-isobutyramide hydrochloride

15     To 50 mg (0.08 mmol) of **73A** in 1 mL of ethanol was added 0.5 mL of concentrated HCL and the mixture was stirred at room temperature for 1 h. The reaction mixture was concentrated and coevaporated twice from ethanol. The residue was triturated twice with ether to give 38 mg of the title compound as a yellow solid.

20     <sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.78-7.98 (m, 0.5H), 7.54-7.74 (m, 6.5H), 7.29-7.42 (m, 2H), 7.02-7.29 (m, 5H), 4.58-4.80 (m, 2H), 3.80-4.17 (m, 1H), 1.56-1.71 (m, 6H).  
MS (CI, NH<sub>3</sub>) 538 (MH<sup>+</sup>)

**Example 74**

25     (R)-2-Amino-N-[2-(1H-indol-3-yl)-1-[6-(7-sulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl]-isobutyramide hydrochloride

A. (R)-(1-[2-(1H-Indol-3-yl)-1-[6-(7-sulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethylcarbamoyl]-1-methyl-ethyl)-carbamic acid tert-butyl ester  
30

According to General Procedure A, 121 mg (0.22 mmol) of **26A** and 47 mg (0.22 mmol) of 1,2,3,4-tetrahydro-isoquinoline-7-sulfonic acid amide were coupled to give 165 mg of **74A**.

B. (R)-2-Amino-N-[2-(1H-indol-3-yl)-1-[6-(7-sulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl]-isobutyramide hydrochloride  
35

To 107 mg (0.14 mmol) of **74A** in 8 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 1.5 h. An

-125-

additional 1 mL of concentrated HCl was added and the reaction mixture was stirred for 3 h. The mixture was concentrated and then coevaporated several times with ethanol to give 61 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.57-7.80 (m, 2H), 7.43-7.62 (m, 1.5H), 6.96-7.41 (m, 7.5H), 1.51-1.67 (d, 6H).  
MS (CI, NH<sub>3</sub>) 643 (MH<sup>+</sup>)

### Example 75

(R)-Piperidine-4-carboxylic acid {1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride  
10 A. {2-(1H-Indol-3-yl)-1-[4-(2-oxo-2,3-dihydro-benzoimidazol-piperidine-1-carbonyl]-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 2.82 g (13.0 mmol) of 4-(2-keto-1-benzimidazoliny)-piperidine and 3.95 g (13.0 mmol) of N-t-BOC-D-tryptophan were coupled and the product triturated with ether to give 2.5 g of 75A.

15 B. 1-{1-[2-Amino-3-(1H-indol-3-yl)-propionyl]-piperidin-4-yl}-1,3-dihydro-benzoimidazol-2-one

According to General Procedure B, 2.50 g (4.97 mmol) of 75A was deprotected to give 1.70 g of 75B.

20 C. (R)-4-{1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}carbamoyl}-piperidine-1-carboxylic acid tert-butyl ester

To a solution of 150 mg (0.372 mmol) of piperidine-1,4-dicarboxylic acid mono-tert-butyl ester, 87 mg (0.216 mmol) of 75B and 96 mg (0.744 mmol) of diisopropylethylamine in 4 mL of methylene chloride was added 197 mg  
25 (0.446 mmol) of BOP reagent and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with 20 mL of ethyl acetate and washed twice each with 10% citric acid and saturated aqueous sodium bicarbonate and once with brine. The solution was dried over MgSO<sub>4</sub> and concentrated to give a white foam which was purified by silica gel  
30 chromatography using a gradient of 100% methylene chloride followed by 1% methanol in methylene chloride to give 52 mg of 75C as a clear oil.

D. (R)-Piperidine-4-carboxylic acid {1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

35 To 20 mg (0.032 mmol) of 75C in 2 mL of ethanol was added 1 mL of concentrated HCl and the mixture was stirred at room temperature for 1 h. The solution was concentrated and coevaporated twice from ethanol, once from

-126-

methylene chloride and twice from ether to give the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 8.19 (d, 0.5H), 7.63-7.69 (m, 0.5H), 7.55-7.61 (m, 0.5H), 7.30-7.39 (m, 1.5H), 7.00-7.31 (m, 5.5H), 5.26-5.37 (m, 0.5H), 5.01-5.24 (m, 0.5H).  
MS (CI, NH<sub>3</sub>) 516 (MH<sup>+</sup>)

### Example 76

(R)-2-Amino-N-{1-(2-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide  
trifluoroacetate

10

A. (R)-2-(2-Amino-2-methyl-propionylamino)-3-(2-methyl-1H-indol-3-yl)-propionic acid

To 1.00 g (4.58 mmol) of 2-D-methyltryptophan in 37 mL of dioxane and 9.2 mL of water was added 1.4 mL (10.0 mmol) of triethylamine and 1.51 g (5.08 mmol) of **33A** and the mixture was stirred overnight at room temperature. The mixture was concentrated to remove excess dioxane and acidified to pH 5 with acetic acid. The aqueous phase was extracted three times with methylene chloride and the combined organics were washed twice with water and once with brine. The organic phase was dried over MgSO<sub>4</sub> and concentrated and the product was purified by silica gel chromatography (98:2:0.1 v/v/v CH<sub>2</sub>Cl<sub>2</sub>:MeOH:acetic acid) to give 1.98 g of **76A**.

15

B. (R)-{(1-Methyl-1-{1-(2-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}carbonyl)-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 107 mg (0.49 mmol) of 4-(2-keto-1-benzimidazol-1-yl)-piperidine and 200 mg (0.49 mmol) of **76A** were coupled and the product was purified by silica gel chromatography (94:6 v/v CH<sub>2</sub>Cl<sub>2</sub>:MeOH) to give 1.87 g of **76B**.

25

C. (R)-2-Amino-N-{1-(2-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide  
trifluoroacetate

30

To 187 mg (0.31 mmol) of **76B** at 0°C was added 2 mL of cold TFA and the mixture was stirred for 1 h at 0°C. The mixture was concentrated and coevaporated once from methylene chloride and twice from toluene to give 180 mg of the title compound as a solid.

35

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 8.32 (d, 0.5H), 7.50-7.57 (m, 0.5H), 7.44 (d, 0.5H), 5.23-5.35 (m, 0.5H), 5.18-5.20 (m, 0.5H), 2.42 (s, 1.5H), 2.32 (s, 1.5H), 1.58-1.66 (m, 6H).



-127-

MS (Cl, NH<sub>3</sub>) 503 (MH<sup>+</sup>)**Example 77**

(R)-Piperidine-4-carboxylic acid {1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

5     A. Piperidine-1,4-dicarboxylic acid tert-butyl ester 2,5-dioxo-pyrrolidin-1-yl

To 15.0 g (65.4 mmol) of piperidine-1,4-dicarboxylic acid mono-tert-butyl ester was added 150 mL of methylene chloride and 20 mL of DMF. To the stirring solution was added 9.30 g (78.5 mmol) of N-hydroxysuccinimide  
10 followed by 15.0 g (78.5 mmol) of DEC, and the mixture was stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate and washed twice with 10% citric acid, four times with a saturated aqueous sodium bicarbonate and once each with water and brine. The solution was dried over MgSO<sub>4</sub> and concentrated to give 20.4 g of **77A** as a white solid.

15     B. (R)-4-[1-Carboxy-2-(1H-indol-3-yl)-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

A mixture of 2.04 g (6.28 mmol) of **77A**, 2.00 g (6.28 mmol) of D-tryptophan trifluoroacetate and 1.16 g (11.5 mmol) of triethylamine in 120 mL of dioxane and 30 mL of water was stirred at room temperature for 17 h. The  
20 reaction mixture was concentrated to remove excess dioxane and then was diluted with water. The aqueous portion was extracted four times with chloroform and concentrated. The crude product was purified by silica gel chromatography (9:1 v/v CHCl<sub>3</sub>:MeOH) followed by (36:4:1 v/v/v CHCl<sub>3</sub>:MeOH:acetic acid) to give 0.87 g of **77B**.

25     C. (R)-4-[1-(1H-Indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl]-piperidine-1-carboxylic acid

According to General Procedure A, 100 mg (0.36 mmol) of **50D** and 150 mg (0.36 mmol) of **77B** were coupled and the product was purified by silica gel chromatography using a gradient elution of 75% ethyl acetate in  
30 hexane to 100% ethyl acetate and 144 mg of **77C** was isolated as a white solid.

D. (R)-Piperidine-4-carboxylic acid {1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

To 144 mg (0.21 mmol) of **77C** in 2 mL of ethanol was added 4 mL of  
35 concentrated HCl and the mixture was stirred at room temperature for 30 min. The solution was concentrated and then coevaporated several times from ethanol. The residue was crystallized from methanol/ethyl acetate and the

-128-

precipitated product was collected by filtration to give 140 mg of the title compound as a white solid.

MS (Cl, NH<sub>3</sub>) 576 (MH<sup>+</sup>)

### Example 78

- 5 (R)-Piperidine-4-carboxylic acid {1-naphthalen-1-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

A. 4-(1-Carboxy-2-naphthalen-1-yl-ethylamino)-piperidine-1-carboxylic acid tert-butyl ester

- 10 To 200 mg (0.607 mmol) of piperidine-1,4-dicarboxylic acid mono-tert-butyl ester in 5 mL of dioxane and 1.2 mL of water was added 0.186 mL (1.33 mmol) of triethylamine and 198 mg (0.607 mmol) of D-3-(1'-Naphthyl)alanine and the mixture was stirred overnight at room temperature. Excess solvent was removed by concentration and the residue was diluted with water and acidified to pH 5 with acetic acid. The solution was extracted three times with methylene chloride, and the combined organics were washed twice with water and once with brine. The organic portion was dried over MgSO<sub>4</sub> and the product was purified by silica gel chromatography (96:4:0.1 v/v/v methylene chloride:methanol:acetic acid) to give 132 mg of **78A**.

- 20 B. (R)-4-{1-Naphthalen-1-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 132 mg (0.32 mmol) of **78A** and 70 mg (0.32 mmol) of 4-(2-keto-1-benzimidazoliny)-piperidine were coupled and the product was purified by silica gel chromatography (96:4 v/v methylene chloride:methanol) to give 132 mg of **78B**.

C. (R)-Piperidine-4-carboxylic acid {1-naphthalen-1-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

- 30 To 132 mg (0.21 mmol) of **78B** dissolved in 2.5 mL of ethanol was added 0.25 mL of concentrated HCl at 0°C. The ice bath was removed and the solution stirred for 1 h. The mixture was coevaporated from methanol and then methylene chloride to give 102 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 8.23 (d, 0.5H), 8.18 (d, 0.5H), 7.79-7.92 (m, 2H), 7.46-7.68 (m, 3H), 7.37-7.44 (m, 1H), 7.14-7.27 (m, 1H), 7.02-7.13 (m, 2.5H), 6.73 (d, 0.5H), 5.38-5.48 (m, 0.5H), 5.26-5.36 (m, 0.5H).

MS (Cl, NH<sub>3</sub>) 527 (MH<sup>+</sup>)

-129-

**Example 79**

(R)-2-Amino-N-{1-[(1H-indol-3-ylmethyl)-2-[7-(morpholine-4-sulfonyl)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethyl]-isobutyramide hydrochloride

5 A. 2,2,2-Trifluoro-1-[7-(morpholine-4-sulfonyl)-3,4-dihydro-1H-isoquinolin-2-yl]-ethanone

2-Trifluoroacetyl-1,2,3,4-tetrahydroisoquinoline-7-sulfonyl chloride can be prepared according to the method outlined by Pendleton et al., J. Pharmacol. Exp. Ther., 208 (1979) p24. A mixture of 250 mg (0.76 mmol) of 2-Trifluoroacetyl-1,2,3,4-tetrahydro-isoquinoline-7-sulfonyl chloride, 86 mg (0.99  
10 mmol) of morpholine and 0.3 mL of anhydrous pyridine in 10 mL of acetone was refluxed for 1 h. The mixture was concentrated and the residue dissolved in chloroform and washed once with water. The organic portion was dried over MgSO<sub>4</sub> and concentrated. The residue was purified by silica gel chromatography (6:4 v/v hexane:ethyl acetate) to give 289 mg of **79A**.

15 B. 7-(Morpholine-4-sulfonyl)-1,2,3,4-tetrahydro-isoquinoline

A mixture of 218 mg (0.58 mmol) of **79A** and 90 mg (0.65 mmol) of potassium carbonate in 10 mL of methanol was stirred at room temperature for 3 h. The mixture was concentrated and the residue was purified by silica gel chromatography (18:2:1 v/v/v ethyl acetate:methanol:triethylamine) to give 163  
20 mg of **79B**.

C. (R)-[1-{1-[(1H-Indol-3-ylmethyl)-2-[7-(morpholine-4-sulfonyl)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethylcarbamoyl]-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 136 mg (0.35 mmol) of **4C** and 100  
25 mg (0.35 mmol) of **79B** were coupled and the product was purified by silica gel chromatography (39:1 v/v chloroform:methanol) to give 194 mg of **79C**.

D. (R)-2-Amino-N-{1-[(1H-indol-3-ylmethyl)-2-[7-(morpholine-4-sulfonyl)-3,4-dihydro-1H-isoquinolin-2-yl]-2-oxo-ethyl]-isobutyramide hydrochloride

To 141 mg (0.22 mmol) of **79C** in 12 mL of ethanol was added 5 mL of  
30 concentrated HCl and the mixture was stirred at room temperature for 3 h. The reaction mixture was concentrated to give 123 mg of the title compound as an off-white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.98-8.07 (bs, 1H), 7.43-7.61 (m, 2.5H), 6.91-7.37 (m, 5H), 5.14-5.29 (m, 1H), 4.48-4.70  
35 (m, 1H), 3.66-3.78 (m, 4H), 3.29-3.46 (m, 4H), 2.84-3.07 (m, 7H), 1.56-1.67 (m, 6H).

MS (Cl, NH<sub>3</sub>) 554 (MH<sup>+</sup>)

-130-

**Example 80**

(R)-Piperidine-4-carboxylic acid [2-(3,4-dihydro-1H-isoquinolin-2-yl)-1-naphthalen-2-ylmethyl-2-oxo-ethyl]-amide hydrochloride

5     A. (R)-4-(1-Carboxy-2-naphthalen-2-yl-ethylcarbamoyl)-piperidine-1-carboxylic acid tert-butyl ester

A mixture of 3.08 g (9.4 mmol) of 77A, 2.03 g (9.4 mmol) of D-3-(2-Naphthyl)-alanine and 2.9 g (28.7 mmol) of triethylamine in 40 mL of dioxane and 10 mL of water was stirred for 72 h at room temperature. Excess dioxane was removed by concentration and the remaining aqueous solution was  
10 diluted with 200 mL of water and acidified to pH 5 with 10% citric acid. The aqueous portion was extracted twice with ethyl acetate. The combined organics were washed twice with saturated aqueous sodium bicarbonate. The orange oil was separated from the aqueous extracts and acidified with 10% acetic acid. The aqueous portion was extracted three times with ethyl acetate  
15 and the combined organics were washed three times with water, once with brine, dried over MgSO<sub>4</sub> and concentrated to give 3.7 g of a yellow oil which was taken up in 200 mL of ether. A white crystalline solid precipitated from solution and was collected by filtration to give 3.31 g of 80A.

20     B. (R)-4-[2-(3,4-Dihydro-1H-isoquinolin-2-yl)-1-naphthalen-2-ylmethyl-2-oxo-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 95 mg (0.22 mmol) of 80A and 29.6 mg (0.22 mmol) of 1,2,3,4-tetrahydroisoquinoline were coupled and the resulting oil triturated with ethyl acetate/hexane to give 70 mg of 80B as a white solid.

25     C. (R)-Piperidine-4-carboxylic acid [2-(3,4-dihydro-1H-isoquinolin-2-yl)-1-naphthalen-2-ylmethyl-2-oxo-ethyl]-amide hydrochloride

To 70 mg (0.13 mmol) of 80B in 6 mL of ethanol was added 2 mL of concentrated HCl at room temperature and the mixture was stirred for 1.5 h. The mixture was concentrated and coevaporated twice from methylene  
30 chloride to give 50 mg of the title compound as a white foam.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.62-7.80 (m, 4H), 7.34-7.48 (m, 3H), 7.05-7.19 (m, 1.5H), 6.82-7.00 (m, 2.5H), 5.18-5.21 (m, 1H), 4.51-4.73 (m, 1.5H), 4.29-4.39 (m, 1H).

MS (CI, NH<sub>3</sub>) 442 (MH<sup>+</sup>)

-131-

**Example 81**

(R)-Piperidine-4-carboxylic acid {2-[4-(2-cyclopropyl-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethyl}-amide hydrochloride

A. 1-(1-Benzyl-piperidin-4-yl)-2-cyclopropyl-1H-benzoimidazole

- 5 A mixture of 530 mg (1.90 mmol) of **50B** and 651 mg (7.56 mmol) of cyclopropanecarboxylic acid were heated at 160°C for 16h. After cooling to room temperature, 10 mL of ethanol was added to the reaction mixture, followed by the slow addition of 20 mL of 2N NaOH in 90 mL of ethanol. The mixture was stirred for 30 min at room temperature and then concentrated. The
- 10 residue was dissolved in ethyl acetate and washed once each with saturated aqueous sodium bicarbonate and brine. The solution was dried over MgSO<sub>4</sub> and concentrated, and the product was purified by silica gel chromatography using an elution gradient of 75% ethyl acetate in hexane to 100% ethyl acetate to give 340 mg of **81A**.

15 B. 2-Cyclopropyl-1-piperidin-4-yl-1H-benzoimidazole hydrochloride

- To 340 mg (1.03 mmol) of **81A** in 3 mL of methylene chloride at -10°C was added 220 mg (1.54 mmol) of α-chloroethylchloroformate and the mixture was stirred at -10°C for 50 min. The mixture was concentrated and the residue dissolved in methanol and heated at 70°C for 1 h. The mixture was
- 20 concentrated and dried in vacuo to give 230 mg of **81B** as a light purple solid.

C. (R)-4-{2-[4-(2-Cyclopropyl-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethylcarbamoyl}-piperidine-1-carboxylic acid tert-butyl ester

- According to General Procedure A, 80 mg (0.29 mmol) of **81B** and 123
- 25 mg (0.29 mmol) of **80A** were coupled and the product was purified by silica gel chromatography (75:25 v/v ethyl acetate:hexane) to give 50 mg of **81C** as a red solid.

-132-

D. (R)-Piperidine-4-carboxylic acid {2-[4-(2-cyclopropyl-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethyl}-amide hydrochloride

To 42 mg (0.065 mmol) of **81C** in 2 mL of ethanol was added 2 mL of concentrated HCl at room temperature and the mixture was stirred for 30 min.

5 The mixture was concentrated to give 40 mg of the title compound as a white solid.

MS (Cl, NH<sub>3</sub>) 550 (MH<sup>+</sup>)

### Example 82

10 (R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

A. (R)-4-{1-Naphthalen-2-ylmethyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-piperidine-1-carboxylic acid

According to General Procedure A, 95 mg (0.34 mmol) of **50D** and 100 mg (0.23 mmol) of **80A** were coupled and the product was purified by silica gel  
15 chromatography using a gradient elution of 55% ethyl acetate in hexane to 100% ethyl acetate to give 97 mg of **82A** as a white solid.

B. (R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

20 To 97 mg (0.14 mmol) of **82A** in 4 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 2 h. The mixture was concentrated to give 70 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 7.77-8.00 (m, 12H), 7.32-7.77 (m, 4H), 5.18-5.38 (m, 1H), 4.68-4.89 (m, 2H), 4.10-4.36 (m,  
25 1H).

MS (Cl, NH<sub>3</sub>) 587 (MH<sup>+</sup>)

### Example 83

(R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[5-(toluene-4-sulfonylamino)-1,3-dihydro-isoindol-2-yl]-ethyl}-amide hydrochloride

30 A. (R)-4-[2-(5-Amino-1,3-dihydro-isoindol-2-yl)-1-naphthalen-2-ylmethyl-2-oxo-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 47 mg (0.35 mmol) of **43C** and 150 mg (0.35 mmol) of **80A** were coupled to give 150 mg of **83A**.

B. (R)-4-{1-Naphthalen-2-ylmethyl-2-oxo-2-[5-(toluene-4-sulfonylamino)-1,3-dihydro-isoindol-2-yl]-ethylcarbamoyl}-piperidine-1-carboxylic acid tert-butyl ester

35 A mixture of 150 mg (0.28 mmol) of **83A**, 61 mg (0.32 mmol) of p-toluenesulfonyl chloride and 39 mg (0.32 mmol) of 4-dimethylaminopyridine

-133-

was stirred at room temperature overnight. The mixture was diluted with methylene chloride and washed twice each with 10% HCl, saturated aqueous sodium bicarbonate, and brine, dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography using a gradient elution of 75% ethyl acetate in hexane to 2% methanol in ethyl acetate to give 50 mg of **83B**.

C. (R)-Piperidine-4-carboxylic acid [1-naphthalen-2-ylmethyl-2-oxo-2-[5-(toluene-4-sulfonylamino)-1,3-dihydro-isoindol-2-yl]-ethyl]-amide hydrochloride

To 50 mg (0.07 mmol) of **83B** in 4 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 3 h. The mixture was concentrated and the residue was crystallized from ethanol/hexane to give 25 mg of the title compound.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 250MHz) (mixture of rotamers) (partial) δ 8.56 (d, 1H), 7.72-7.89 (m, 5H), 7.57-7.68 (m, 2H), 7.39-7.52 (m, 5H), 7.24-7.31 (m, 2H), 6.91-7.16 (m, 3H), 4.49-4.73 (m, 2H), 4.36-4.47 (m, 1H), 2.47-2.68 (m, 2H), 2.39 (s, 3H).

MS (CI, NH<sub>3</sub>) 598 (MH<sup>+</sup>)

#### Example 84

(R)-Piperidine-4-carboxylic acid [1-(5-benzoylamino-1,3-dihydro-isoindole-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-amide hydrochloride

A. (R)-4-[1-(5-Amino-1,3-dihydro-isoindole-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 40 mg (0.30 mmol) of **43C** and 125 mg (0.30 mmol) of **77B** were coupled to give 170 mg of **84A**.

B. (R)-4-[1-(5-Benzoylamino-1,3-dihydro-isoindole-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

A mixture of 170 mg (0.32 mmol) of **84A**, 52 mg (0.37 mmol) of benzoyl chloride and 45 mg (0.37 mmol) of 4-dimethylaminopyridine in 15 mL of methylene chloride was stirred overnight at room temperature. The reaction mixture was diluted with methylene chloride and washed twice each with 10% HCl, saturated aqueous sodium bicarbonate and brine. The organic portion was dried over MgSO<sub>4</sub> and concentrated. The product was purified by silica gel chromatography using an elution gradient of 90% ethyl acetate in hexane to 2% methanol in ethyl acetate to give 100 mg of **84B**.

C. (R)-Piperidine-4-carboxylic acid [1-(5-benzoylamino-1,3-dihydro-isoindole-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-amide hydrochloride

-134-

To 100 mg (0.16 mmol) of **84B** in 4 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 1 h. The mixture was diluted with ethanol and concentrated. The residue was crystallized from ethanol/hexane to give 50 mg of the title compound.

5  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 250MHz) (mixture of rotamers) (partial)  $\delta$  7.91-7.99 (m, 2H), 7.48-7.70 (m, 6H), 7.03-7.38 (m, 5H), 4.68-4.80 (m, 1H), 4.45-4.59 (m, 1H), 4.03-4.16 (dd, 1H), 2.98-3.13 (m, 2H), 2.59-2.76 (m, 1H), 1.76-2.10 (m, 4H).

MS (CI,  $\text{NH}_3$ ) 537 ( $\text{MH}^+$ )

10

### Example 85

(R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

A. (R)-4-{1-Naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-  
15 benzimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl}-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 225 mg (0.53 mmol) of **80A** and 117 mg (0.54 mmol) of 4-(2-keto-1-benzimidazol-1-yl)piperidine were coupled and the product was purified by silica gel chromatography using an elution  
20 gradient of 1% to 3% methanol in methylene chloride to give 178 mg of **85A**.

B. (R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

To 165 mg (0.26 mmol) of **85A** in 3 mL of ethanol was added 1.5 mL of  
25 concentrated HCl and the mixture was stirred at room temperature for 3 h. The mixture was concentrated to give 117 mg of the title compound as a solid.

$^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 300MHz) (mixture of rotamers) (partial)  $\delta$  7.70-7.89 (m, 4H), 7.37-7.50 (m, 3H), 7.22-7.29 (m, 0.5H), 6.90-7.04 (m, 3H), 6.68 (d, 0.5H), 5.25-5.36 (m, 1H), 4.67-4.74 (m, 1H), 4.38-4.50 (m, 1H), 4.11-4.32 (m,  
30 1H).

MS (CI,  $\text{NH}_3$ ) 527 ( $\text{MH}^+$ )

### Example 86

(R)-Piperidine-4-carboxylic acid [1-(7-ethylsulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-amide hydrochloride

A. 2-Trifluoroacetyl-1,2,3,4-tetrahydro-isoquinoline-7-sulfonic acid  
2-Trifluoroacetyl-1,2,3,4-tetrahydro-isoquinoline-7-sulfonyl chloride can be prepared according to the method outlined by Pendleton et al., J. Pharmacol. Exp. Ther., 208 (1979) p24. A mixture of 295 mg (0.90 mmol) of 2-  
35



-135-

Trifluoroacetyl-1,2,3,4-tetrahydro-isoquinoline-7-sulfonyl chloride, 95 mg (1.17 mmol) of ethylamine hydrochloride and 0.3 mL (2.15 mmol) of triethylamine in 10 mL of acetone was refluxed for 1.5 h. The reaction mixture was filtered and concentrated and the product was purified by silica gel chromatography (6:4 v/v hexane:ethyl acetate) to give 283 mg of **86A**.

B. 1,2,3,4-Tetrahydro-isoquinoline-7-sulfonic acid ethylamide

To 225 mg (0.67 mmol) of **86A** and 104 mg (0.75 mmol) of potassium carbonate in 10 mL of methanol and 0.5 mL of water was stirred at room temperature for 3 h. The mixture was concentrated to give 335 mg of **86B**.

C. (R)-4-[1-(7-Ethylsulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 145 mg (0.35 mmol) of **86B** and 85 mg (0.35 mmol) of **77B** were coupled and the product was purified by silica gel chromatography (39:1 v/v chloroform:methanol) to give 68 mg of **86C**.

D. (R)-Piperidine-4-carboxylic acid [1-(7-ethylsulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-amide hydrochloride

To 68 mg (0.11 mmol) of **86C** in 6 mL of ethanol was added 2.5 mL of concentrated HCl and the mixture was stirred at room temperature for 5 h. The reaction mixture was concentrated to give 62 mg of the title compounds as a tan solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.96-8.02 (bs, 0.5H), 7.62-7.53 (m, 0.5H), 7.52-7.61 (m, 2H), 6.91-7.37 (m, 5H), 5.09-5.20 (m, 1H), 4.42-4.70 (m, 1H), 1.00-1.11 (m, 3H).

MS (CI, NH<sub>3</sub>) 538 (MH<sup>+</sup>)

**Example 87**

(R)-2-Amino-N-[1-(7-ethylsulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-isobutyramide hydrochloride

A. (R)-{1-[1-(7-Ethylsulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-1-methyl-ethyl}-carbamic acid tert-butyl ester

According to General Procedure A, 136 mg (0.35 mmol) of **4C** and 85 mg (0.35 mmol) of **86B** were coupled and the product was purified by silica gel chromatography (39:1 v/v chloroform:methanol) followed by (9:1 v/v chloroform:methanol) to give 73 mg of **87A**.

B. (R)-2-Amino-N-[1-(7-ethylsulfamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethyl]-isobutyramide hydrochloride

To 62 mg (0.10 mmol) of **87A** in 6 mL of ethanol was added 2.5 mL of concentrated HCl and the mixture was stirred at room temperature for 5 h and then concentrated to give 56 mg of the title compound as an off-white solid.

-136-

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial)  $\delta$  8.28-8.41 (bs, 0.5H), 7.42-7.59 (m, 2H), 6.85-7.07 (m, 5H), 5.19-5.25 (m, 1H), 4.39-4.68 (m, 2H), 2.71-2.87 (m, 2H), 1.40-1.63 (m, 6H), 0.92-1.06 (m, 3H).  
MS (CI, NH<sub>3</sub>) 513 (MH<sup>+</sup>)

5

**Example 88**

(R)-2-[3-(1H-Indol-3-yl)-2-[(piperidine-4-carbonyl)-amino]-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride

A. 3,4-Dihydro-1H-isoquinoline-2,6-dicarboxylic acid 2-tert-butyl ester

To 1.20 g (4.1 mmol) of **5C** in 36 mL of methanol was added 0.62 g (4.5 mmol) of potassium carbonate and the mixture was heated at reflux for 7 h. The product was purified by silica gel chromatography (9:1 v/v chloroform:methanol) to give 1.49 g of **88A**.

B. 6-Ethylcarbamoyl-3,4-dihydro-1H-isoquinoline-2-carboxylic acid tert-butyl ester

According to General Procedure A, 650 mg (2.34 mmol) of **88A** and 191 mg (2.34 mmol) of ethylamine hydrochloride were coupled and the product was purified by silica gel chromatography (39:1 v/v chloroform:methanol) followed by (1:1 v/v ethyl acetate:methanol) to give 566 mg of **88B**.

20 C. 1,2,3,4-Tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride

To 493 mg (1.62 mmol) of **88B** in 30 mL of ethanol was added 12.5 mL of concentrated HCl and the mixture was stirred at room temperature for 5 h. The mixture was concentrated to give 384 mg of **88C** as a white solid.

25 D. (R)-4-[1-(6-Ethylcarbamoyl-3,4-dihydro-1H-isoquinoline-2-carbonyl)-2-(1H-indol-3-yl)-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 99 mg (0.41 mmol) of **88C** and 170 mg (0.41 mmol) of **77B** were coupled and the product was purified by silica gel chromatography (39:1 v/v chloroform:methanol) followed by (9:1 v/v chloroform:methanol) to give 107 mg of **88D**.

30

-137-

E. (R)-2-{3-(1H-Indol-3-yl)-2-[(piperidine-4-carbonyl)-amino]-propionyl}-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride

To 66 mg (0.11 mmol) of **88D** in 6 mL of ethanol was added 2.5 mL of concentrated HCl and the mixture was stirred at room temperature for 5 h. The mixture was concentrated to give 56 mg of the title compound as an off-white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.52-7.60 (m, 1.5H), 7.38-7.45 (m, 1H), 7.24 (d, 0.5H), 5.11-5.24 (m, 1H), 4.40-4.69 (m, 1.5H), 4.04-4.13 (m, 0.5H), 1.13-1.24 (m, 3H).

MS (CI, NH<sub>3</sub>) 502 (MH<sup>+</sup>)

### Example 89

(R)-2-{3-Naphthalen-2-yl-2-[(piperidine-4-carbonyl)-amino]-propionyl}-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride

A. 4-[2-(6-Ethylcarbamoyl-3,4-dihydro-1H-isoquinolin-2-yl)-1-naphthalen-2-ylmethyl-2-oxo-ethylcarbamoyl]-piperidine-1-carboxylic acid

According to General Procedure A, 96 mg (0.40 mmol) of **88C** and 170 mg (0.40 mmol) of **80A** were coupled and the product was purified by silica gel chromatography (39:1 v/v chloroform:methanol) to give 88 mg of **89A**.

B. (R)-2-{3-Naphthalen-2-yl-2-[(piperidine-4-carbonyl)-amino]-propionyl}-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride

To 61 mg (0.10 mmol) of **89B** in 6 mL of ethanol was added 2.5 mL of concentrated HCl and the mixture was stirred at room temperature for 5 h. The reaction mixture was concentrated to give 58 mg of the title compound as an off-white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.46-7.76 (m, 4.5H), 7.09-7.43 (m, 5.5H), 5.08-5.32 (m, 1H), 1.09-1.24 (m, 3H).

MS (CI, NH<sub>3</sub>) 513 (MH<sup>+</sup>)

### Example 90

(R)-2-Dimethylamino-N-{1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

A. (R)-2-Dimethylamino-N-{1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

To a solution of 92 mg (0.17 mmol) of **35D** in 2 mL of methanol was added 127 mg of powdered 3A sieves, 12 mg (0.40 mmol) of 37% w/w formaldehyde, 27 mg (0.43 mmol) of sodium cyanoborohydride and 103 mg (1.72 mmol) of acetic acid. The mixture was stirred for 60 h at room temperature and then filtered through celite and concentrated. The residue

-138-

was dissolved in ethyl acetate and washed three times with 1N NaOH, once with water and once with brine, dried over MgSO<sub>4</sub> and concentrated. The resulting oil was triturated to give 39 mg of the title compound as a white solid.

<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 7.68-7.89 (m, 4H), 7.42-7.54 (m, 4H), 6.90-7.08 (m, 3H), 5.32-5.47 (m, 1H), 4.68-4.77 (m, 1H), 4.33-4.51 (m, 1H), 4.16-4.28 (m, 1H), 2.64 (d, 3H), 2.34 (d, 3H).  
MS (CI, NH<sub>3</sub>) 529 (MH<sup>+</sup>)

### Example 91

(R)-3-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-3-methyl-butyramide hydrochloride

A. (R)-2-Amino-3-(1H-indol-3-yl)-propionic acid methyl ester

To 9.65 g (30.3 mmol) of 4A in 40 mL of ethanol was added 20 mL of concentrated HCl and the mixture was stirred at room temperature for 2.5 h. The mixture was concentrated and the residue was diluted with ethyl acetate.  
The mixture was washed twice with saturated aqueous sodium bicarbonate and once with brine, dried over MgSO<sub>4</sub> and concentrated to give 5.12 g of 91A as a yellow oil.

B. (R)-2-(3-tert-Butoxycarbonylamino-3-methyl-butyrylamino)-3-(1H-indol-3-yl)-propionic acid

According to General Procedure A, 2.01 g (9.2 mmol) of 91A and 2.00 g (9.2 mmol) of 3-tert-Butoxycarbonylamino-3-methyl-butyric acid were coupled to give (R)-2-(3-tert-Butoxycarbonylamino-3-methyl-butyrylamino)-3-(1H-indol-3-yl)-propionic acid methyl ester. The crude ester was hydrolyzed according to General Procedure D to give 3.53 g of 91B as a white solid.

C. (R)-2-(2-(1H-Indol-3-yl)-1-[4-(2-phenyl-benzoimidazol-1-yl)-piperidine-1-carbonyl]-ethylcarbamoyl)-1,1-dimethyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 350 mg (1.3 mmol) of 91B and 510 mg (1.3 mmol) of 50D were coupled and the product was purified by silica gel chromatography using a gradient of 60% ethyl acetate in hexane to 100% ethyl acetate to give 180 mg of 91C.

D. (R)-3-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-3-methyl-butyramide hydrochloride

To 100 mg (0.27 mmol) of 91C in 3 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 2 h. The mixture was concentrated and precipitated from ethanol/hexane to give 125 mg of the title compound as a white solid.

MS (CI, NH<sub>3</sub>) 564 (MH<sup>+</sup>)

-139-

**Example 92**

(R)-Piperidine-4-carboxylic acid {2-[4-(3-methyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethyl}-amide hydrochloride

5       A. 4-(2-Oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carboxylic acid tert-butyl ester

To a mixture of 1.00 g (4.6 mmol) of 4-(2-keto-1-benzimidazoliny)-piperidine and 1.00 g (4.6 mmol) of 4-dimethylaminopyridine in 20 mL of methylene chloride at 0°C was added 80 mL of di-tert-butyl dicarbonate in 80 mL of methylene chloride dropwise over 30 min. The mixture was allowed to warm to room temperature and stirred overnight. The mixture was washed three times each with 10% HCl, saturated aqueous sodium bicarbonate and brine, dried over MgSO<sub>4</sub> and concentrated. The residue was purified by silica gel chromatography using an elution gradient of 50% ethyl acetate in hexane to 100% ethyl acetate to give 1.16 g of **92A**.

15       B. 4-(3-Methyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidine-1-carboxylic acid tert-butyl ester

To 390 mg (1.2 mmol) of **92A** in 3 mL of DMF was added 47 mg (1.2 mmol) of sodium hydride (60% dispersion in mineral oil), followed by 520 mg (3.7 mmol) of iodomethane and the mixture was stirred for 72 h at room temperature. The mixture was diluted with ethyl acetate and filtered. The filtrate was washed twice with brine and concentrated to give 400 mg of **92B**.

20       C. 1-Methyl-3-piperidin-4-yl-1,3-dihydro-benzoimidazol-2-one hydrochloride

To 400 mg (1.2 mmol) of **92B** in 6 mL of ethanol was added 3 mL of concentrated HCl and the mixture was stirred at room temperature for 1.5 h. The mixture was concentrated to give 330 mg of **92C** as a white solid.

25       D. (R)-4-{2-[4-(3-Methyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethylcarbamoyl}-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 100 mg (0.37 mmol) of **92C** and 159 mg (0.37 mmol) of **80A** were coupled and the product was purified by silica gel chromatography using an elution gradient of 100% ethyl acetate to 4% methanol in ethyl acetate to give 120 mg of **92D**.

30       E. (R)-Piperidine-4-carboxylic acid {2-[4-(3-methyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethyl}-amide hydrochloride

-140-

To 120 mg (0.19 mmol) of **92D** in 5 mL of ethanol was added 2 mL of concentrated HCl and the mixture was stirred at room temperature for 1.5 h. The mixture was diluted with ethanol and concentrated, and the residue was crystallized from ethanol/hexane to give 45 mg of the title compound.

5  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 250MHz) (mixture of rotamers) (partial)  $\delta$  7.79-7.99 (m, 4H), 7.43-7.78 (m, 3.5H), 7.30-7.39 (m, 0.5H), 7.01-7.19 (m, 3H), 5.26-5.39 (m, 1H), 4.68-4.79 (m, 1H), 4.41-4.60 (m, 1H), 4.13-4.32 (m, 1H).  
MS (CI,  $\text{NH}_3$ ) 540 ( $\text{MH}^+$ )

### Example 93

10 (R)-2-[3-Naphthalen-2-yl-2-[(piperidine-4-carbonyl)-amino]-propionyl]-2,3-dihydro-1H-isoindole-5-carboxylic acid ethylamide hydrochloride

A. 3,4-Dimethyl-benzoic acid ethyl ester

A mixture of 10.00 g (66.6 mmol) of 3,4-dimethylbenzoic acid and 3 mL of concentrated sulfuric acid in 200 mL of ethanol was heated at 50°C for 2 h,  
15 and 65°C for 6 h. The mixture was concentrated and the residue was diluted with ethyl acetate and washed twice each with brine, saturated aqueous sodium bicarbonate, and brine. The solution was dried over  $\text{MgSO}_4$  and concentrated to give 12.0 g of **93A**.

B. 3,4-Bis-bromomethyl-benzoic acid ethyl ester

20 A mixture of 12.0 g (67.3 mmol) of **93A**, 26.10 g (146.6 mmol) of N-bromosuccinimide and 130 mg (0.79 mmol) of 2,2'-azobis(isobutyronitrile) in 130 mL of carbon tetrachloride was heated at reflux for 17 h. After cooling to room temperature, the mixture was filtered and concentrated to give 22.0 g of **93B** as a yellow oil.

-141-

C 2,3-Dihydro-1H-isoindole-5-carboxylic acid ethyl ester

To 22.0 g (65.4 mmol) of **93B** in 150 mL of benzene was added a solution of 7.87 g (73.4 mmol) of benzylamine and 15.27 g (151 mmol) of triethylamine in 50 mL of benzene dropwise over 30min, and the mixture was then refluxed for 18 h. After standing at room temperature for 24 h, the mixture was filtered and the filtrate washed once each saturated aqueous sodium bicarbonate and water. The solution was dried over MgSO<sub>4</sub> and concentrated. The crude 2-Benzyl-2,3-dihydro-1H-isoindole-5-carboxylic acid ethyl ester was dissolved in 100 mL of ethanol and 20 mL of water and 2.00 g of 10% palladium on carbon was added. The mixture was hydrogenated for 18 h at 45 psi, filtered and concentrated and the residue was purified by silica gel chromatography using an elution gradient of 100% ethyl acetate to 10% diethylamine in ethyl acetate to give 220 mg of **93C**.

D. 2-{2-[(1-tert-Butoxycarbonyl-piperidine-4-carbonyl)-amino]-3-naphthalen-2-yl-propionyl}-2,3-dihydro-1H-isoindole-5-carboxylic acid ethyl ester

According to General Procedure A, 200 mg (1.05 mmol) of **93C** and 446 mg (1.05 mmol) of **80A** were coupled to give 460 mg of **93D** as a brown solid.

E. (R)-2-{2-[(1-tert-Butoxycarbonyl-piperidine-4-carbonyl)-amino]-3-naphthalen-2-yl-propionyl}-2,3-dihydro-1H-isoindole-5-carboxylic acid

To 410 mg (0.68 mmol) of **93D** in 50 mL of ethanol containing 1.05 mL of 2N NaOH was stirred overnight at room temperature. The mixture was concentrated and the residue partitioned between chloroform and 10% HCl. The layers were separated and the aqueous portion was extracted twice with chloroform. The combined organics were concentrated to give 350 mg of **93E**.

F. 4-[2-(5-Ethylcarbamoyl-1,3-dihydro-isoindol-2-yl)-1-naphthalen-2-ylmethyl-2-oxo-ethylcarbamoyl]-piperidine-1-carboxylic acid tert-butyl ester

According to General Procedure A, 350 mg (0.61 mmol) of **93E** and 50 mg (0.61 mmol) of ethylamine hydrochloride were coupled and the product was purified by silica gel chromatography using a gradient of 100% ethyl acetate to 4% methanol in ethyl acetate to give 40 mg of **93F**.

G. (R)-2-{3-Naphthalen-2-yl-2-[(piperidine-4-carbonyl)-amino]-propionyl}-2,3-dihydro-1H-isoindole-5-carboxylic acid ethylamide hydrochloride

-142-

To 40 mg (0.07 mmol) of **93F** in 5 mL of ethanol was added 2 mL of concentrated HCl, and the mixture was stirred at room temperature for 2 h. The mixture was concentrated and the residue crystallized from ethanol/hexane to give 16 mg of the title compound as a white solid.

- 5  $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 250MHz) (mixture of rotamers) (partial)  $\delta$  8.63 (d, 1H), 7.71-7.90 (m, 6H), 7.39-7.65 (m, 4H), 5.01-5.20 (m, 2H), 4.67-4.93 (q, 2H), 4.48-4.59 (m, 1H).  
MS (Cl,  $\text{NH}_3$ ) 499 ( $\text{MH}^+$ )

**Example 94**

- 10 (R)-1-Methyl-piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride  
A. (R)-1-Methyl-piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride

- 15 To 40 mg (0.06 mmol) of **82B** in 4 mL of methanol was added 40 mL (0.64 mmol) of acetic acid, 0.16 mL (0.16 mmol) of 1M sodium cyanoborohydride in THF, 12 mL (0.15 mmol) of formaldehyde and 40 mg of **3A** molecular sieves and the mixture was stirred for 72 h at room temperature. The mixture was concentrated and the residue dissolved in ethyl acetate,  
20 washed twice with 2N NaOH and once with brine, dried over  $\text{MgSO}_4$  and concentrated. The solid was recrystallized from ethanol/hexane to give 24 mg of the title compound.

- $^1\text{H}$  NMR ( $\text{CD}_3\text{OD}$ , 250MHz) (mixture of rotamers) (partial)  $\delta$  7.81-7.98 (m, 4H), 7.40-7.76 (m, 10H), 7.27-7.39 (m, 2H), 5.24-5.38 (m, 1H), 2.37 (s, 3H).  
25 MS (Cl,  $\text{NH}_3$ ) 600 ( $\text{MH}^+$ )

**Example 95**

(R)-2-Amino-N-{1-benzyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride

- A. (R)-2-(2-tert-Butoxycarbonylamino-2-methyl-propionylamino)-3-phenyl-propionic acid

- 30 To a mixture of 500 mg (3.03 mmol) of D-phenylalanine and 910 mg (3.03 mmol) of **33A** in 10 mL of water and 40 mL of dioxane was added 1.26 mL (9.08 mmol) of triethylamine and the mixture was stirred at room temperature for 30 min., and then heated at 30°C for 16 h. The mixture was  
35 diluted with chloroform and the aqueous portion acidified to pH 4 with acetic acid. The layers were separated and the organic phase was washed three times with brine, dried over  $\text{MgSO}_4$  and concentrated to give 488 mg of **95A**.



-143-

B. (R)-(1-(1-Benzyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethylcarbamoyl)-1-methyl-ethyl)-carbamic acid tert-butyl ester

According to General Procedure A, 76 mg (0.22 mmol) of **95A** and 60 mg (0.22 mmol) of **50D** were coupled and the product was purified by silica gel chromatography (75:25 v/v ethyl acetate:hexanes) to give 120 mg of **95B**.

C. (R)-2-Amino-N-(1-benzyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl)-isobutyramide hydrochloride

To 120 mg (0.20 mmol) of **95B** in 5 mL of ethanol was added 5 mL of concentrated HCl and the mixture was stirred at room temperature for 40 min. The mixture was diluted with ethanol and concentrated and the residue was crystallized from methanol/hexanes to give 65 mg of the title compound as a white solid.


<sup>1</sup>H NMR (CD<sub>3</sub>OD, 300MHz) (mixture of rotamers) (partial) δ 8.24 (d, 0.5H), 7.64-7.90 (m, 8.5H), 7.21-7.43 (m, 5H), 5.17-5.27 (m, 0.33H), 5.02-5.09 (m, 0.67H), 4.18-4.26 (m, 0.33H), 3.94-4.06 (m, 0.67H), 1.71 (s, 2H), 1.56 (s, 3H), 1.48 (s, 1H).

MS (CI, NH<sub>3</sub>) 510 (MH<sup>+</sup>)

-144-

## CLAIMS

1. A compound of the formula:



10

wherein

Z is  $-\text{COCR}^1\text{R}^2\text{LCOANR}^4\text{R}^5$ ;


L is  $\text{NR}^6$ , O or  $\text{CH}_2$ ;

15 W is hydrogen or in combination with X is a benzo fusion in which W and X are linked to form a phenyl ring optionally substituted with one to three substituents independently selected from  $\text{R}^{3a}$ ,  $\text{T-R}^{3b}$  and  $\text{R}^{12}$ ;


Y is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>4</sub>-C<sub>10</sub> cycloalkyl or aryl or phenyl, each optionally substituted with one to three substituents independently selected from  $\text{R}^{3a}$ ,  $\text{R}^{3b}$  and  $\text{R}^{12}$ ;

20

X is  $\text{OR}^2$ ,



30



35


or X in combination with W is a benzofusion in which W and X are linked to form a phenyl ring optionally substituted with one to three substituents independently selected from  $\text{R}^{3a}$ ,  $\text{T-R}^{3b}$  and  $\text{R}^{12}$ ;

$\text{R}^1$  is C<sub>1</sub>-C<sub>10</sub> alkyl, aryl, aryl (C<sub>1</sub>-C<sub>6</sub> alkyl) and C<sub>3</sub>-C<sub>7</sub> cycloalkyl (C<sub>1</sub>-C<sub>6</sub> alkyl) or C<sub>1</sub>-C<sub>5</sub> alkyl-K-C<sub>1</sub>-C<sub>5</sub> alkyl, aryl (C<sub>0</sub>-C<sub>5</sub> alkyl)-K-(C<sub>1</sub>-C<sub>5</sub> alkyl), (C<sub>3</sub>-C<sub>7</sub>) cycloalkyl(C<sub>1</sub>-C<sub>5</sub>alkyl)-K-(C<sub>1</sub>-C<sub>5</sub> alkyl) where K is O, S(O)<sub>m</sub>,

-145-

- $N(R^2)C(O)$ ,  $C(O)N(R^2)$ ,  $OC(O)$ ,  $C(O)O$ ,  $-CR^2=CR^2-$  or  $-C\equiv C-$  where the aryl groups are defined below and  $R^2$  and the alkyl groups may be further substituted by 1-5 halogens,  $S(O)_mR^{2a}$ , 1 to 3 of  $OR^{2a}$  or  $C(O)OR^{2a}$  and the aryl groups may be further substituted independently by phenyl, phenoxy,
- 5 arylalkyloxy, halophenyl, 1 to 3 of  $C_1$ - $C_6$  alkyl, 1 to 3 of halogen, 1 to 2 of  $OR^2$ , methylenedioxy,  $S(O)_mR^2$ , 1 to 2 of  $CF_3$ ,  $OCF_3$ , nitro,  $N(R^2)(R^2)$ ,  $N(R^2)C(O)(R^2)$ ,  $C(O)OR^2$ ,  $C(O)N(R^2)(R^2)$ ,  $SO_2N(R^2)(R^2)$ ,  $N(R^2)SO_2$  aryl or  $N(R^2)SO_2R^2$ ;
- $R^{2c}$  is hydrogen,  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_7$  cycloalkyl, and may be joined with
- 10  $R^1$  to form a  $C_3$ - $C_8$  ring optionally including oxygen, sulfur or  $NR^{2a}$ ;
- $R^2$  is hydrogen,  $C_1$ - $C_6$  alkyl,  $C_3$ - $C_7$  cycloalkyl;
- $R^{2a}$  is hydrogen or  $C_1$ - $C_6$  alkyl;
- $R^{2b}$  is hydrogen  $C_1$ - $C_8$  alkyl,  $C_1$ - $C_8$  halogenated alkyl,  $C_3$ - $C_8$  cycloalkyl, alkylaryl or aryl;
- 15  $R^{3a}$  is H, F, Cl, Br, I,  $CH_3$ ,  $OCH_3$  or  $CF_3$ ;
- $R^{12}$  is H, F, Cl, Br, I,  $CH_3$ ,  $OCH_3$  or  $CF_3$ ;
- T is a bond or is phenyl or a 5 or 6-membered heterocycle containing 1-3 hetero atoms selected independently from nitrogen, sulfur or oxygen, each optionally substituted independently with one to three substituents selected
- 20 from F, Cl, Br, I,  $CH_3$ ,  $OCH_3$ ,  $OCF_3$  and  $CF_3$ ;
- $R^{3b}$  is hydrogen,  $CONR^8R^9$ ,  $SO_2NR^8R^9$ ,  $COOH$ ,  $COO(C_1-C_6)alkyl$ ,  $NR^2SO_2R^9$ ,  $NR^2CONR^8R^9$ ,  $NR^2SO_2NR^8R^9$ ,  $NR^2C(O)R^9$ , imidazolyl, thiazolyl or tetrazolyl;
- $R^4$  and  $R^5$  are independently hydrogen,  $C_1$ - $C_6$  alkyl, substituted  $C_1$ - $C_6$
- 25 alkyl where the substituents independently may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3  $C_1$ - $C_{10}$  alkanoyloxy, 1 to 3  $C_1$ - $C_6$  alkoxy, phenyl, phenoxy, 2-furyl,  $C_1$ - $C_6$  alkoxy carbonyl,  $S(O)_m(C_1-C_6 alkyl)$ ; or  $R^4$  and  $R^5$  can be taken together to form  $-(CH_2)_rL_a(CH_2)_s-$  where  $L_a$  is  $C(R^2)_2$ ,  $S(O)_m$  or  $N(R^2)$ , r and s are independently 1 to 3 and  $R^2$  is as defined above;
- 30  $R^6$  is hydrogen or  $C_1$ - $C_6$  alkyl, and may be joined with  $R^{2c}$  to form a  $C_3$ - $C_8$  ring;
- $R^{50}$  is 4-morpholino, 4-(1-methylpiperazinyl),  $C_3$ - $C_7$  cycloalkyl or  $C_1$ - $C_6$  alkyl each optionally substituted with one to three substituents individually selected from F, OH,  $OCH_3$ ,  $OCF_3$ ,  $CF_3$  and  $C_3$ - $C_7$  cycloalkyl;
- 35 M is  $-C(O)-$  or  $-SO_2-$ ;
- $A^1$  is a bond,  $C_1$ - $C_6$  alkylene,  $C_1$ - $C_6$  haloalkylenyl or  $C_1$ - $C_6$  hydroxyalkylenyl;
- A is a bond or is

-146-



5

where x and y are independently 0-3;

$\text{Z}^1$  is  $\text{N-R}^2$  or O; or  $\text{Z}^1$  is a bond;

$\text{R}^7$  and  $\text{R}^{7a}$  are independently hydrogen, C1-C6 alkyl, trifluoromethyl, phenyl, substituted C1-C6 alkyl where the substituents are imidazolyl, phenyl, indolyl, p-hydroxyphenyl,  $\text{OR}^2$ ,  $\text{S(O)}_m\text{R}^2$ ,  $\text{C(O)OR}^2$ , C3-C7 cycloalkyl,  $\text{N(R}^2\text{)(R}^2\text{)}$ ,  $\text{C(O)N(R}^2\text{)(R}^2\text{)}$ ; or  $\text{R}^7$  and  $\text{R}^{7a}$  can independently be joined to one or both of  $\text{R}^4$  and  $\text{R}^5$  groups to form alkylene bridges between the terminal nitrogen and the alkyl portion of the  $\text{R}^7$  and  $\text{R}^{7a}$  groups, wherein the bridge contains 1 to 5 carbon atoms; or  $\text{R}^7$  and  $\text{R}^{7a}$  may be joined to form a 3- to 7-membered ring.

$\text{R}^9$  is hydrogen, C1-C6 alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl each optionally substituted with one to three substituents selected from Cl, F,  $\text{CH}_3$ ,  $\text{OCH}_3$ ,  $\text{OCF}_3$  and  $\text{CF}_3$ ;

$\text{R}^8$  is hydrogen, C1-C6 alkyl, substituted C1-C6 alkyl where the substituents may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3 C1-C10 alkanoyloxy, 1 to 3 C1-C6 alkoxy, phenyl, phenoxy, C1-C6 alkoxycarbonyl,  $\text{S(O)}_m\text{(C1-C6 alkyl)}$ ; or  $\text{R}^8$  and  $\text{R}^9$  can be taken together to form  $\text{---(CH}_2\text{)}_r\text{L}_a\text{(CH}_2\text{)}_s\text{---}$  where  $\text{L}_a$  is  $\text{C(R}^2\text{)}_2$ , O,  $\text{S(O)}_m$  or  $\text{N(R}^2\text{)}$ , r and s are independently 1 to 3.

$\text{F}^1$ , B, D and E are carbon, nitrogen, one of which is joined to  $\text{A}^1$  and each of the remaining of  $\text{F}^1$ , B, D and E may be optionally substituted with  $\text{R}^{aa}$ ,  $\text{R}^{bb}$ ; or  $\text{F}^1$ , B, D and E, if not joined to  $\text{A}^1$  may also be sulfur, oxygen or carbonyl;  $\text{F}^1$ , B, D and E may form a saturated or unsaturated ring; and one of  $\text{F}^1$ , B, D or E may be optionally missing to afford a saturated or unsaturated five-membered ring;

$\text{R}^{aa}$  is H, C1-C8 alkyl optionally substituted with one to three halogens; or phenyl optionally substituted with one to three substituents independently selected from halogen,  $\text{CH}_3$ ,  $\text{OCH}_3$ ,  $\text{OCF}_3$ ,  $\text{CF}_3$  and C1-C8 alkyl aryl;

$\text{R}^{bb}$  is C1-C8 alkyl, C1-C8 alkoxy, C1-C8 thioalkyl, each optionally substituted with one to three substituents independently selected from halogen,  $\text{CH}_3$ ,  $\text{OCH}_3$ ,  $\text{OCF}_3$ ,  $\text{CF}_3$ ; C3-C8 cycloalkyl; phenyl optionally substituted with one to three substituents independently selected from halo,  $\text{CH}_3$ ,  $\text{OCH}_3$ ,  $\text{OCF}_3$  or  $\text{CF}_3$ ;  $\text{---O-C1-C8 alkyl}$ ; or  $\text{---S-C1-C8 alkyl}$ .

-147-

G, H<sup>1</sup>, I and J are carbon, nitrogen, sulfur or oxygen atoms which form an aromatic ring, a partially saturated ring or a saturated ring; one of G, H<sup>1</sup>, I or J may be optionally missing to afford a five-membered ring;

aryl is phenyl, naphthyl or a 5- or 6- membered ring with 1 to 3

- 5 heteroatoms independently selected from oxygen, sulfur and nitrogen; or a bicyclic ring system consisting of a 5 or 6 membered heterocyclic ring with 1 to 3 heteroatoms independently selected from nitrogen, sulfur or oxygen, fused to a phenyl ring, each aryl ring being optionally and independently substituted with up to three substituents selected from R<sup>3a</sup>, R<sup>3b</sup> and R<sup>12</sup>;

10 m is 0 to 2;

n is 0 to 2;

q is 0 to 3;

and pharmaceutically acceptable salts and individual diastereomers thereof.

- 15 2. A compound of claim 1 wherein

Z is -COCR<sup>1</sup>R<sup>2c</sup>NHCOANR<sup>4</sup>R<sup>5</sup>

A is -CR<sup>7</sup>R<sup>7a</sup>(CH<sub>2</sub>)<sub>Y</sub>; Y is 0 to 3;

and R<sup>2c</sup> is H or CH<sub>3</sub>;

R<sup>7</sup> is C<sub>1</sub>-C<sub>3</sub> alkyl;

- 20 R<sup>7a</sup> is H or C<sub>1</sub>-C<sub>3</sub> alkyl;

R<sup>4</sup> is hydrogen or C<sub>1</sub>-C<sub>3</sub> alkyl; or R<sup>4</sup> and R<sup>7</sup> are combined to form an alkylene bridge;

R<sup>5</sup> is hydrogen or C<sub>1</sub>-C<sub>3</sub> alkyl optionally substituted with one or two hydroxyl groups.


- 25 3. A compound of claim 2 wherein

R<sup>1</sup> is selected from the group consisting of 1-indolyl-CH<sub>2</sub>-; 2-indolyl-CH<sub>2</sub>-; 3-indolyl-CH<sub>2</sub>-; 1-naphthyl-CH<sub>2</sub>-; 2-naphthyl-CH<sub>2</sub>-; 1-benzimidazolyl-CH<sub>2</sub>-; 2-benzimidazolyl-CH<sub>2</sub>-; phenyl-C<sub>1</sub>-C<sub>4</sub> alkyl-; 2-, 3- or 4-pyridyl-C<sub>1</sub>-C<sub>4</sub> alkyl-; thienyl-C<sub>1</sub>-C<sub>4</sub> alkyl-; phenyl-(CH<sub>2</sub>)<sub>n</sub>-O-CH<sub>2</sub>- where n is zero to three; 30 phenyl-CH<sub>2</sub>O-phenyl-CH<sub>2</sub>-; and 3-benzothieryl-CH<sub>2</sub>-; or any of the above groups substituted in the aryl portion with one to three F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub> substituents.

4. A compound of claim 3 wherein R<sup>2c</sup> is hydrogen; R<sup>7</sup> and R<sup>7a</sup> are methyl and R<sup>4</sup> and R<sup>5</sup> are hydrogen and R<sup>1</sup> is C<sub>6</sub>H<sub>5</sub>CH<sub>2</sub>-O-CH<sub>2</sub>-; 1-naphthyl-

- 35 CH<sub>2</sub>-, 2-naphthyl-CH<sub>2</sub>-, phenylpropyl or,

-148-




5

and the aryl portion of  $R^1$  is optionally substituted with fluorine,  $CH_3$  or  $CF_3$ .

5. A compound of claim 3 which is of the formula

10



15

wherein  $n$  is zero or one;  $m$  is one or two;

$R^{3a}$  is H, F, Cl, Br, I,  $CH_3$ ,  $OCH_3$  or  $CF_3$ ;

20  $R^{12}$  is H, F, Cl, Br, I,  $CH_3$ ,  $OCH_3$  or  $CF_3$ ;

$T$  is a bond or is an aryl group selected from phenyl, pyridyl, pyrimidyl, thienyl, pyrrolyl, pyrazolyl, oxazolyl, isoxazolyl, imidazolyl, thiazolyl and tetrazolyl; each optionally substituted with one to three F,  $CH_3$ , Cl,  $OR^8$ ,  $OCH_3$ ,  $OCF_3$  or  $CF_3$ ;

25  $R^{3b}$  is hydrogen,  $CONR^8R^9$ ,  $SO_2NR^8R^9$ ,  $COOH$ ,  $COO(C_1-C_6)alkyl$ ,  $NHSO_2R^9$ ,  $NHC(O)NR^8R^9$ ,  $NHSO_2NR^8R^9$ ,  $NHC(O)R^9$ ,  $NR^8R^9$ , imidazolyl, thiazolyl or tetrazolyl;

$R^9$  is hydrogen,  $C_1-C_6$  alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl each optionally substituted with one to three substituents selected from Cl, F, 30  $CH_3$ ,  $OCH_3$ ,  $OCF_3$  and  $CF_3$ ;

$R^8$  is hydrogen,  $C_1-C_6$  alkyl,  $C_3-C_7$  cycloalkyl, substituted  $C_1-C_6$  alkyl where the substituents may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3  $C_1-C_{10}$  alkanoyloxy, 1 to 3  $C_1-C_6$  alkoxy, phenyl, phenoxy,  $C_1-C_6$  alkoxy carbonyl, or  $S(O)_2(C_1-C_6 alkyl)$ ; or  $R^8$  and  $R^9$  can be taken together to form 35  $-(CH_2)_rL_a(CH_2)_s-$  where  $L_a$  is  $C(R^2)_2$ , oxo,  $S(O)_2$  or  $N(R^2)$ , and  $r$  and  $s$  are independently 1 to 3.

6. A compound of claim 5 wherein:  
 $m$  is one;

-149-

R<sup>12</sup> is hydrogen;

R<sup>3a</sup> is hydrogen, F or Cl;

T is phenyl, thienyl, thiazolyl, oxazolyl, isoxazolyl or pyrazolyl, each optionally substituted with one to three of substituents, selected from F, OH,

5 OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub> and CH<sub>3</sub>; and

R<sup>3b</sup> is hydrogen, C(O)NR<sup>8</sup>R<sup>9</sup>, NHC(O)NR<sup>8</sup>R<sup>9</sup>, NHS(O)<sub>2</sub>R<sup>9</sup>, or NHC(O)R<sup>9</sup>.

7. A compound of claim 5 wherein:

m is one; and

10 T is a bond.

8. A compound of claim 7 wherein

n is one;

R<sup>3a</sup> and R<sup>12</sup> are hydrogen.

R<sup>3b</sup> is CONR<sup>8</sup>R<sup>9</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, COOH, COO(C<sub>1</sub>-C<sub>6</sub>)alkyl, NHSO<sub>2</sub>R<sup>9</sup>,  
15 NHSO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, NHC(O)NR<sup>8</sup>R<sup>9</sup>, NHC(O)R<sup>9</sup>, NR<sup>8</sup>R<sup>9</sup> or OR<sup>8</sup>;

R<sup>9</sup> is hydrogen, phenyl, or thienyl optionally substituted with one to three substituents selected from F, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; or R<sup>9</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one to three substituents selected from F, OH, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; or R<sup>8</sup>; and

20 R<sup>8</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl or C<sub>3</sub>-C<sub>7</sub> cycloalkyl where the substituents may be 1 to 5 halo or 1 to 3 hydroxy; or R<sup>8</sup> and R<sup>9</sup> can be taken together to form -(CH<sub>2</sub>)<sub>r</sub>La(CH<sub>2</sub>)<sub>s</sub>- wherein La is C(R<sup>2</sup>)<sub>2</sub>, O, s(O)<sub>m</sub> or N(R<sup>2</sup>) where r and s are independently 1 to 3.

9. A compound of claim 7 wherein

25 n is zero;

m is one;

R<sup>3a</sup> is hydrogen;

R<sup>12</sup> is hydrogen;


R<sup>3b</sup> is CONR<sup>8</sup>R<sup>9</sup>, NHC(O)NR<sup>8</sup>R<sup>9</sup>, NHS(O)<sub>2</sub>R<sup>9</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>,  
30 NHSO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, NHCOR<sup>9</sup> or OR<sup>8</sup>;

R<sup>9</sup> is phenyl, or thienyl optionally substituted with one to three substituents selected from F, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; or R<sup>9</sup> is C<sub>1</sub>-C<sub>6</sub> alkyl optionally substituted with one to three substituents selected from F, OH, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; and

35 R<sup>8</sup> is hydrogen, C<sub>1</sub>-C<sub>6</sub> alkyl, or C<sub>3</sub>-C<sub>7</sub> cycloalkyl each optionally substituted with 1 to 5 halo or 1 to 3 hydroxy; or R<sup>8</sup> and R<sup>9</sup> can be taken together to form -(CH<sub>2</sub>)<sub>n</sub>La(CH<sub>2</sub>)<sub>s</sub>- wherein La is C(R<sup>2</sup>)<sub>2</sub>, O, s(O)<sub>m</sub> or N(R<sup>2</sup>) where r and s are independently 1 to 3.

-150-

10. A compound of claim 3 wherein:  
 n is one;  
 m is one;  
 W is hydrogen;  
 5 Y is hydrogen or methyl;  
 X is  $R^{50}-M-\overset{\overset{|}{N}}{\text{---}}\text{Aryl}$ ;  
 M is -C(O)-;  
 $R^{50}$  is 4-morpholino, 4-(1-methylpiperazinyl), C3-C7 cycloalkyl or C1-C6 alkyl each optionally substituted with one to three substituents individually selected from F, OH, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; and  
 10 Aryl is phenyl, pyridyl, thienyl, pyrimidyl or thiazolyl, each optionally substituted with one to three substituents individually selected from F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>.
11. A compound of claim 3 wherein:  
 n and m are one;  
 Y is phenyl, pyridyl, pyrimidyl, thienyl, or thiazolyl, each optionally substituted with one to three substituents independently selected from F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>;  
 20 X is -C(O)-NR<sup>8</sup>R<sup>9</sup>;  
 $R^9$  is hydrogen, C1-C6 alkyl, phenyl, thiazolyl, imidazolyl, furyl or thienyl each optionally substituted with one to three substituents selected from Cl, F, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> and CF<sub>3</sub>; and  
 $R^8$  is hydrogen, C1-C6 alkyl, substituted C1-C6 alkyl where the  
 25 substituents may be 1 to 5 halo, 1 to 3 hydroxy, 1 to 3 C1-C10 alkanoyloxy, 1 to 3 C1-C6 alkoxy, phenyl, phenoxy, C1-C6 alkoxy carbonyl, S(O)<sub>m</sub>(C1-C6 alkyl); or  $R^8$  and  $R^9$  can be taken together to form -(CH<sub>2</sub>)<sub>r</sub>L<sub>a</sub>(CH<sub>2</sub>)<sub>s</sub>- where L<sub>a</sub> is C(R<sup>2</sup>)<sub>2</sub>, oxo, S(O)<sub>m</sub> or N(R<sup>2</sup>), and r and s are independently 1 to 3.
12. A compound of claim 3 wherein Y is hydrogen and X is:




wherein n and m are one and A<sup>1</sup> is a bond.



-151-

13. A compound of claim 12 wherein X is



wherein R<sup>3a</sup> is hydrogen, F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>;

R<sup>12</sup> is hydrogen, F, Cl, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>;

R<sup>aa</sup> is hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl optionally substituted with one to three  
 20 halogens; or phenyl optionally substituted with one to three substituents  
 independently selected from halogen, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub> and C<sub>1</sub>-C<sub>8</sub>  
 alkyl aryl; and

R<sup>bb</sup> is hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl, optionally substituted with one to three  
 substituents independently selected from halogen, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub>, CF<sub>3</sub>;  
 25 C<sub>3</sub>-C<sub>8</sub> cycloalkyl; phenyl optionally substituted with one to three substituents  
 independently selected from halo, CH<sub>3</sub>, OCH<sub>3</sub>, OCF<sub>3</sub> or CF<sub>3</sub>; -O-C<sub>1</sub>-C<sub>8</sub> alkyl;  
 or -S-C<sub>1</sub>-C<sub>8</sub> alkyl.

14. A compound of claim 13 wherein  
 G, H<sup>1</sup>, I and J are carbon.

30 15. A compound of claim 13 wherein  
 G is nitrogen; and  
 H<sup>1</sup>, I and J are carbon.

16. A compound of claim 13 wherein  
 H<sup>1</sup> is nitrogen; and  
 35 G, I and J are carbon.

17. A compound of claim 13 wherein  
 I is nitrogen; and  
 G, H<sup>1</sup> and J are carbon.


-152-

18. A compound of claim 13 wherein J is nitrogen; and G, H<sup>1</sup> and I are carbon.

19. A compound of claim 3 wherein Z is


5

10



20. A compound of claim 3 wherein Z is


15



- 20 wherein each n is independently zero or one.


21. A compound of claim 19 wherein R<sup>1</sup> is

25



22. A compound of claim 20 wherein R<sup>1</sup> is

30



35


23. A compound of claim 19 wherein R<sup>1</sup> is -CH<sub>2</sub>OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, 1-naphthyl-CH<sub>2</sub>-, or 2-naphthyl-CH<sub>2</sub>-.

-153-

24. A compound of claim 20 wherein R<sup>1</sup> is -CH<sub>2</sub>OCH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>C<sub>6</sub>H<sub>5</sub>, 1-naphthyl-CH<sub>2</sub>-, or 2-naphthyl-CH<sub>2</sub>-.

25. A compound of claim 5 wherein Z is

5



10

26. A compound of claim 25 wherein n is one.

27. A compound of claim 25 wherein n is zero.

28. A compound of claim 5 selected from the group consisting of:

(R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[6-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide hydrochloride;

15 (R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid ethylamide hydrochloride;

(R)-2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinoline-6-carboxylic acid phenylamide

20 hydrochloride;

(R)-2-Amino-N-(2-(1H-indol-3-yl)-1-[6-[2-(3-methyl-ureido)-phenyl]-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl)-isobutyramide hydrochloride;

(R)-2-Amino-N-{2-(1H-indol-3-yl)-1-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinoline-2-carbonyl]-ethyl}-isobutyramide;

25 (R)-N-{2-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-1,2,3,4-tetrahydro-isoquinolin-7-yl}-benzamide;

(R)-2-Amino-N-{1-benzyloxymethyl-2-oxo-2-[7-(toluene-4-sulfonylamino)-3,4-dihydro-1H-isoquinolin-2-yl]-ethyl}-isobutyramide hydrochloride; and

30 (R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[5-(toluene-4-sulfonylamino)-1,3-dihydro-isoindol-2-yl]-ethyl}-amide hydrochloride.

29. A compound of claim 10 which is

2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(phenyl-propionyl-amino)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride or

35 (R)-N-[1-[4-(Acetyl-phenyl-amino)-piperidine-1-carbonyl]-2-(1H-indol-3-yl)-ethyl]-2-amino-isobutyramide

30. A compound of claim 11 selected from the group consisting of:

-154-

- (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride;  
 2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide;  
 5 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-phenyl-4-(pyrrolidine-1-carbonyl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride; and  
 (R)-1-[2-(2-Amino-2-methyl-propionylamino)-3-(1H-indol-3-yl)-propionyl]-4-phenyl-piperidine-4-carboxylic acid (4-hydroxy-butyl)-amide hydrochloride.
- 10 31. A compound of claim 12 selected from the group consisting of:  
 2-Amino-N-{1-(R)-benzyloxymethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;  
 2-Amino-N-{1-(R)-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide;  
 15 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(1H-indol-3-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride;  
 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-[4-(2-methyl-benzoimidazol-1-yl)-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide hydrochloride;  
 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;  
 20 (R)-2-Amino-N-[2-[4-(6-fluoro-benzo[d]isoxazol-3-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide hydrochloride;  
 (R)-2-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-1,2-dihydro-imidazo[4,5-b]pyridin-3-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;  
 25 (R)-2-Amino-N-[2-[4-(5-chloro-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-(1H-indol-3-ylmethyl)-2-oxo-ethyl]-isobutyramide;  
 (R)-2-Amino-N-{1-benzo[b]thiophen-3-ylmethyl-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;  
 2-Amino-N-{1-(5-fluoro-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;  
 30 2-Amino-N-{1-(1-methyl-1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-isobutyramide hydrochloride;  
 (R)-Piperidine-4-carboxylic acid {1-naphthalen-2-ylmethyl-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-amide hydrochloride;  
 35 (R)-3-Amino-N-{1-(1H-indol-3-ylmethyl)-2-oxo-2-[4-(2-phenyl-benzoimidazol-1-yl)-piperidin-1-yl]-ethyl}-3-methyl-butyramide hydrochloride;  
 and

-155-

(R)-Piperidine-4-carboxylic acid {2-[4-(3-methyl-2-oxo-2,3-dihydro-benzoimidazol-1-yl)-piperidin-1-yl]-1-naphthalen-2-ylmethyl-2-oxo-ethyl)-amide hydrochloride.

32. The compound of claim 11 which is

5 2-Amino-N-{1-(R)-benzyloxymethyl-2-[4-(morpholine-4-carbonyl)-4-phenyl-piperidin-1-yl]-2-oxo-ethyl}-isobutyramide.

33. A method for increasing levels of endogenous growth hormone in a human or an animal which comprises administering to such human or animal an effective amount of a compound of Claim 1.

10 34. A composition useful for increasing the endogenous production or release of growth hormone in a human or an animal which comprises an inert carrier and an effective amount of a compound of Claim 1.

35. A composition useful for increasing the endogenous production or release of growth hormone in a human or an animal which comprises an  
15 effective amount of a compound of Claim 1 and a growth hormone secretagogues selected from GHRP-6, Hexarelin, GHRP-1, growth hormone releasing factor (GRF), IGF-1 or IGF-2, or B-HT920.

36. A method of treating or preventing osteoporosis which comprises administering to a mammal in need of such treatment or prevention an amount  
20 of a compound of claim 1 which is effective in treating or preventing osteoporosis.

37. A method for treating or preventing diseases or conditions which may be treated or prevented by growth hormone which comprises administering to a mammal in need of such treatment or prevention an amount  
25 of a compound of claim 1 which is effective in promoting release of said growth hormone.

38. A method for the treatment of osteoporosis which comprises administering to a patient with osteoporosis a combination of a bisphosphonate compound such as alendronate, and a compound of Claim 1.

# INTERNATIONAL SEARCH REPORT

International Application No <b>PC 1/IB 95/00333</b>					
<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 6    C07K5/06    C07D401/12    C07D401/04    C07D401/14    C07D413/06 C07D211/56    C07D211/64    C07D207/14    C07D401/06    A61K31/455 A61K38/05    C07D209/44    C07K5/02					
According to International Patent Classification (IPC) or to both national classification and IPC					
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 6    C07K    C07D    A61K					
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched					
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)					
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>					
Category *	Citation of document, with indication, where appropriate, of the relevant passages				Relevant to claim No.
A	WO,A,94 13696 (MERCK & CO. ) 23 June 1994 cited in the application see the whole document				1-38
E	WO,A,95 13069 (MERCK & CO.) 18 May 1995 see the whole document				1-38
<input type="checkbox"/> Further documents are listed in the continuation of box C.					
<input checked="" type="checkbox"/> Patent family members are listed in annex.					
* Special categories of cited documents :					
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*A* document member of the same patent family</p> </div> </div>					
Date of the actual completion of the international search  <div style="text-align: center; font-size: 1.2em;">22 January 1996</div>			Date of mailing of the international search report  <div style="text-align: center; font-size: 1.2em;">31.01.96</div>		
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. ( + 31-70 ) 340-2040, Tx. 31 651 epo nl, Fax ( + 31-70 ) 340-3016			Authorized officer  <div style="text-align: center; font-size: 1.2em;">Masturzo, P</div>		

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 95/00333

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 33, 36-38  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 33 and 36-38 refer to methods of treatment of the human body, the search was carried out and based on the alleged effects of the compounds.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 95/00333

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9413696	23-06-94	AU-B- 5232093	23-06-94
		CA-A- 2110670	12-06-94
		CN-A- 1092071	14-09-94
		EP-A- 0615977	21-09-94
		FI-A- 952863	09-06-95
		JP-A- 6263737	20-09-94
		NO-A- 952295	10-08-95
		PL-A- 309331	02-10-95
		SI-A- 9300646	30-09-94
		AU-B- 5232193	23-06-94
		CA-A- 2110672	12-06-94
		CN-A- 1092767	28-09-94
		EP-A- 0662481	12-07-95
		FI-A- 952862	09-06-95
		JP-A- 7097367	11-04-95
		NO-A- 952294	10-08-95
		PL-A- 309332	02-10-95
		SI-A- 9300647	30-09-94
		WO-A- 9419367	01-09-94
WO-A-9513069	18-05-95	AU-B- 1172995	29-05-95